



US009181220B2

(12) **United States Patent**  
**Singh et al.**

(10) **Patent No.:** **US 9,181,220 B2**  
(45) **Date of Patent:** **\*Nov. 10, 2015**

(54) **N-SUBSTITUTED-  
HETEROCYCLOALKYLOXYBENZAMIDE  
COMPOUNDS AND METHODS OF USE**

(71) Applicant: **Rigel Pharmaceuticals, Inc.**, South San Francisco, CA (US)

(72) Inventors: **Rajinder Singh**, Belmont, CA (US); **Jiaxin Yu**, Foster City, CA (US); **Hui Hong**, Foster City, CA (US); **Sambaiah Thota**, Fremont, CA (US); **Xiang Xu**, Foster City, CA (US); **Ihab S. Darwish**, San Carlos, CA (US)

(73) Assignee: **Rigel Pharmaceuticals, Inc.**, South San Francisco, CA (US)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/189,638**

(22) Filed: **Feb. 25, 2014**

(65) **Prior Publication Data**

US 2014/0179738 A1 Jun. 26, 2014

**Related U.S. Application Data**

(63) Continuation of application No. 13/185,030, filed on Jul. 18, 2011, now Pat. No. 8,697,727, which is a continuation of application No. 11/963,742, filed on Dec. 21, 2007, now Pat. No. 8,012,955.

(60) Provisional application No. 60/882,312, filed on Dec. 28, 2006, provisional application No. 60/988,719, filed on Nov. 16, 2007.

(51) **Int. Cl.**

**C07D 211/58** (2006.01)  
**C07D 211/96** (2006.01)  
**C07D 401/12** (2006.01)  
**C07D 401/14** (2006.01)  
**C07D 405/14** (2006.01)  
**C07D 409/14** (2006.01)

(52) **U.S. Cl.**

CPC ..... **C07D 401/12** (2013.01); **C07D 211/58** (2013.01); **C07D 211/96** (2013.01); **C07D 401/14** (2013.01); **C07D 405/14** (2013.01); **C07D 409/14** (2013.01)

(58) **Field of Classification Search**

CPC .. **C07D 211/58**; **C07D 211/94**; **C07D 211/96**;  
**C07D 401/12**; **C07D 401/14**; **C07D 405/14**;  
**C07D 409/14**; **A61K 31/444**

See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,332,732 A 7/1994 Scott et al.  
5,438,064 A 8/1995 Mobilio et al.  
5,665,719 A 9/1997 Bock et al.  
5,958,945 A 9/1999 Imbert et al.  
5,965,591 A 10/1999 Kojima et al.  
6,096,771 A 8/2000 Kojima et al.  
6,472,394 B1 10/2002 McKittrick et al.  
6,492,355 B1 12/2002 Alcaraz et al.  
6,821,965 B1 11/2004 Brown et al.  
7,001,900 B2 2/2006 Jacobsen et al.  
7,208,491 B2 4/2007 Fertig et al.  
7,273,868 B2 9/2007 Yamada et al.  
8,119,809 B2 2/2012 Hong et al.  
8,129,390 B2 3/2012 Darwish et al.  
8,314,107 B2 11/2012 Singh et al.  
2003/0216582 A1 11/2003 Nikolaides et al.  
2004/0082542 A1 4/2004 Mjalli et al.  
2005/0038081 A1 2/2005 Brown et al.  
2005/0107399 A1 5/2005 Boman et al.  
2005/0131018 A1 6/2005 Sendzik  
2005/0182040 A1 8/2005 Imazaki et al.  
2005/0187261 A1 8/2005 Verner et al.

(Continued)

**FOREIGN PATENT DOCUMENTS**

EP 0158380 A1 10/1985  
EP 0254322 A1 1/1988

(Continued)

**OTHER PUBLICATIONS**

Tudose et al., "Synthesis and Properties of dinitrobenzamido-TEMPO derivatives", ARKIVOC vi (2005), 225-237.


(Continued)

*Primary Examiner* — Sarah Pihonak

(74) *Attorney, Agent, or Firm* — McDonnell Boehnen Hulbert & Berghoff LLP; Travis Young

(57) **ABSTRACT**

The present invention provides N-substituted-heterocycloalkyloxybenzamide compounds, as well as pharmaceutical compositions and methods of use. One embodiment of the invention is a compound having the structure



in which R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, T, n, w and x are as described herein. In certain embodiments of the invention, a compound of the present invention activates the AMPK pathway, and can be used to treat metabolism-related disorders and conditions.

**24 Claims, 5 Drawing Sheets**

(56)

**References Cited**

## U.S. PATENT DOCUMENTS

2005/0234057 A1 10/2005 Lee et al.  
 2005/0282864 A1 12/2005 McArthur et al.  
 2006/0009449 A1 1/2006 McArthur et al.  
 2006/0084679 A1 4/2006 McArthur et al.  
 2006/0167053 A1 7/2006 Lino et al.  
 2006/0205772 A1 9/2006 Coppola et al.  
 2006/0270653 A1 11/2006 Drutu et al.  
 2007/0021386 A1 1/2007 Mjalli et al.  
 2007/0123515 A1 5/2007 Nettekoven et al.  
 2007/0123525 A1 5/2007 Nettekoven et al.  
 2007/0123526 A1 5/2007 Nettekoven et al.

## FOREIGN PATENT DOCUMENTS

WO WO 96/06837 A1 3/1996  
 WO WO 99/59959 A1 11/1999  
 WO WO0012074 3/2000  
 WO WO 01/64639 A2 9/2001  
 WO WO 02/34718 A1 5/2002  
 WO 03/015774 2/2003  
 WO WO 03/022856 A1 3/2003  
 WO WO03018586 3/2003  
 WO WO03022856 3/2003  
 WO WO 03/028641 A2 4/2003  
 WO WO03070732 8/2003  
 WO WO 03/072578 A1 9/2003  
 WO WO 03/097047 A1 11/2003  
 WO WO 2004/005295 A1 1/2004  
 WO WO 2004/022060 A2 3/2004  
 WO WO2004054974 7/2004  
 WO WO 2004/065351 A1 8/2004  
 WO WO 2004/099204 A1 11/2004  
 WO WO2004111003 12/2004  
 WO WO 2005/014533 A2 2/2005  
 WO WO2005061442 7/2005  
 WO WO2005116000 12/2005  
 WO WO 2006/009054 A1 1/2006

WO WO 2006/020879 A1 2/2006  
 WO WO 2006/056848 A1 6/2006  
 WO WO 2006/066924 A2 6/2006  
 WO WO 2006/067139 A1 6/2006  
 WO WO 2006/091862 A2 8/2006  
 WO WO 2006/108965 A2 10/2006  
 WO WO 2007/003962 A2 1/2007  
 WO WO 2007/057329 A1 5/2007  
 WO WO 2007/065948 A1 6/2007  
 WO WO 2007/073935 A1 7/2007  
 WO WO2007075688 7/2007  
 WO WO2007098086 8/2007  
 WO WO2007/143824 12/2007  
 WO WO 2007/149929 A1 12/2007  
 WO WO2007143823 12/2007  
 WO WO2008017685 2/2008  
 WO WO2008133975 11/2008

## OTHER PUBLICATIONS

Bell et al, "Development of Orally Active Oxytocin Antagonists: Studies on 1-(1-{4-[1-(2-Methyl-1-oxidopyridin-3-ylmethyl)piperidin-4-yloxy]-2-methoxybenzoyl}piperidin-4-yl)-1,4-dihydrobenz[d][1,3]oxazin-2-one (L-372,662) and Related Pyridines", *Journal of Medicinal Chemistry* (1998), 2146-2163, vol. 41, No. 12.  
 Database Registry [Online] Chemical Abstract Service, Columbus, Ohio, USA; Dec. 19, 2007, XP002474730; retrieved from STN; CAS RF: 958847-90-0.  
 Copending U.S. Appl. No. 13/185,030, filed Jul. 18, 2011.  
 Database Registry [Online] Chemical Abstract Service, Columbus, Ohio, USA; Dec. 20, 2007, XP002474728; retrieved from STN; CAS RF: 958939-37-2.  
 International Search Report and Written Opinion for PCT/US2007/088742, Apr. 28, 2008.  
 Fan et al, *Bioorganic & Medicinal Chemistry Letters*, 1997, Elsevier Science, vol. 7, No. 24, pp. 3107-3112.  
 Kitamura et al, *Chem. Pharm. Bull.*, 2001, Pharmaceutical Society of Japan, vol. 49, issue 3, pp. 268-277.  
 West, *Solid State Chemistry and its Applications*, 1987, John Wiley & Sons, pp. 358 & 365.




FIG. 1




FIG. 2




FIG. 3




FIG. 4





FIG. 5

1

**N-SUBSTITUTED-  
HETEROCYCLOALKYLOXYBENZAMIDE  
COMPOUNDS AND METHODS OF USE**

**CROSS REFERENCE TO RELATED  
APPLICATIONS**

This application is a continuation of U.S. patent application Ser. No. 13/185,030, filed Jul. 18, 2011, which is a continuation of U.S. patent application Ser. No. 11/963,742, filed Dec. 21, 2007, now U.S. Pat. No. 8,012,955, which claims priority under 35 U.S.C. §119(e) to U.S. Provisional Patent Application Ser. Nos. 60/882,312, filed Dec. 28, 2006, and 60/988,719, filed Nov. 16, 2007, each of which is hereby incorporated herein by reference.

**BACKGROUND OF THE INVENTION**

**1. Field of the Invention**

The invention relates generally to compounds, pharmaceutical compositions and methods of use of the compounds and compositions containing them. The invention relates more particularly to N-substituted-heterocycloalkyloxybenzamide compounds and pharmaceutical compositions thereof, and to methods of treating and preventing disease states such as type II diabetes, atherosclerosis and cardiovascular disease using N-substituted-heterocycloalkyloxybenzamide compounds.

**2. Technical Background**

Adiponectin is a protein hormone exclusively expressed in and secreted from adipose tissue and is the most abundant adipose-specific protein. Adiponectin has been implicated in the modulation of glucose and lipid metabolism in insulin-sensitive tissues. Decreased circulating adiponectin levels have been demonstrated in some insulin-resistant states, such as obesity and type 2 diabetes mellitus and also in patients with coronary artery disease, atherosclerosis and hypertension. Adiponectin levels are positively correlated with insulin sensitivity, HDL (high density lipoprotein) levels and insulin stimulated glucose disposal and inversely correlated with adiposity and glucose, insulin and triglyceride levels. Thiazolidinedione drugs, which enhance insulin sensitivity through activation of the peroxisome proliferator-activated receptor- $\gamma$ , increase endogenous adiponectin production in humans.


Adiponectin binds its receptors in liver and skeletal muscle and thereby activates the 5'-AMP-activated protein kinase (AMPK) pathway. Adiponectin receptors 1 and 2 are membrane-bound proteins found in skeletal muscle and liver tissue. Being a multi-substrate enzyme, AMPK regulates a variety of metabolic processes, such as glucose transport, glycolysis and lipid metabolism. It acts as a sensor of cellular energy homeostasis and is activated in response to certain hormones and muscle contraction as well as to intracellular metabolic stress signals such as exercise, ischemia, hypoxia and nutrient deprivation. Once activated, AMPK switches on catabolic pathways (such as fatty acid oxidation and glycolysis) and switches off ATP-consuming pathways (such as lipogenesis). Adiponectin improves insulin sensitivity by directly stimulating glucose uptake in adipocytes and muscle and by increasing fatty acid oxidation in liver and muscle, resulting in reduced circulating fatty acid levels and reduced intracellular triglyceride contents. Moreover, adiponectin decreases glycogen concentration by reducing the activity of glycogen synthase. Adiponectin also plays a protective role against inflammation and atherosclerosis. It suppresses the expression of adhesion molecules in vascular endothelial cells and cytokine production from macrophages, thus inhib-

2

iting the inflammatory processes that occur during the early phases of atherosclerosis. What is needed are compounds, pharmaceutical compositions and methods of using them to treat disease states associated with circulating adiponectin levels, such as type II diabetes, atherosclerosis and cardiovascular disease.

**SUMMARY OF THE INVENTION**

A first aspect of the invention is a compound having structural formula (I)



(I)

or a pharmaceutically acceptable salt, solvate, hydrate, or N-oxide thereof, wherein

R<sup>1</sup> is H;

R<sup>2</sup> is —Hca;

each R<sup>3</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN;


w is 0, 1, 2, or 3;

n is 0, 12 or 3;

each R<sup>4</sup> is independently selected from (C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; (C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>4</sup> on the same carbon optionally combine to form oxo;

x is 0, 1, 2, 3 or 4;

T is (C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>; or



in which

Q is (C<sub>0</sub>-C<sub>3</sub> alkyl)-, in which each carbon of the (C<sub>0</sub>-C<sub>3</sub> alkyl)- is optionally and independently substituted with one or two R<sup>16</sup>;

each R<sup>16</sup> is independently selected from (C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; (C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>16</sup> on the same carbon optionally combine to form oxo;

the ring system denoted by "A" is heteroaryl, aryl, cycloalkyl or heterocycloalkyl;

each R<sup>5</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca,

3

—(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN; and

y is 0, 1, 2, 3 or 4;

in which

each L is independently selected from —NR<sup>9</sup>C(O)O—, —NR<sup>9</sup>C(O)—NR<sup>9</sup>—, —NR<sup>9</sup>C(O)S—, —NR<sup>9</sup>C(O)—, —NR<sup>9</sup>C(S)O—, —NR<sup>9</sup>C(S)—NR<sup>9</sup>—, —NR<sup>9</sup>C(S)S—, —NR<sup>9</sup>C(S)—, —OC(O)NR<sup>9</sup>—, —SC(O)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —OC(S)NR<sup>9</sup>—, —SC(S)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —S(O)<sub>0-2</sub>—, —C(O)O—, —OC(O)—, —C(S)O—, —OC(S)—, —C(O)S—, —SC(O)—, —C(S)S—, —SC(S)—, —OC(O)O—, —SC(O)O—, —OC(O)S—, —SC(S)O—, —OC(S)S—, —NR<sup>9</sup>C(NR<sup>2</sup>)NR<sup>9</sup>—, —NR<sup>9</sup>SO<sub>2</sub>—, —SO<sub>2</sub>NR<sup>9</sup>— and —NR<sup>9</sup>SO<sub>2</sub>NR<sup>9</sup>—,

each R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>10</sup> is independently selected from H, —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-O-(C<sub>0</sub>-C<sub>6</sub> alkyl) and —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>-(C<sub>0</sub>-C<sub>6</sub> alkyl), each R<sup>9</sup> is independently selected from —H, —(C<sub>1</sub>-C<sub>4</sub> alkyl) and —C(O)—(C<sub>1</sub>-C<sub>4</sub> alkyl),

each Ar is an optionally substituted aryl,

each Het is an optionally substituted heteroaryl,

each Cak is an optionally substituted cycloalkyl,

each Hca is an optionally substituted heterocycloalkyl, and

each alkyl is optionally substituted,

provided that the compound is not

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-methoxybenzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-(furan-2-ylmethyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(pyridin-4-ylmethyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(3-phenylpropyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(methylsulfonyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide;


N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide; or

N-(1-benzylpiperidin-4-yl)-3-(1-benzylpiperidin-4-yloxy)benzamide.

Another aspect of the invention is a pharmaceutical composition comprising at least one pharmaceutically acceptable carrier, diluent or excipient; and a compound having structural formula (I)

4



(I)

or a pharmaceutically acceptable salt, solvate, hydrate, or N-oxide thereof, wherein

R<sup>1</sup> is H;

R<sup>2</sup> is —Hca;

each R<sup>3</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN;


w is 0, 1, 2 or 3;

n is 0, 1, 2 or 3;

each R<sup>4</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>4</sup> on the same carbon optionally combine to form oxo;

x is 0, 1, 2, 3 or 4;

T is —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>; or



in which

Q is —(C<sub>0</sub>-C<sub>3</sub> alkyl)-, in which each carbon of the (C<sub>0</sub>-C<sub>3</sub> alkyl) is optionally and independently substituted with one or two R<sup>16</sup>;

each R<sup>16</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>16</sup> on the same carbon optionally combine to form oxo;

the ring system denoted by "A" is heteroaryl, aryl, cycloalkyl or heterocycloalkyl;

each R<sup>5</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN; and

y is 0, 1, 2, 3 or 4;

in which

each L is independently selected from —NR<sup>9</sup>C(O)O—, —NR<sup>9</sup>C(O)—NR<sup>9</sup>—, —NR<sup>9</sup>C(O)S—, —NR<sup>9</sup>C(O)—, —NR<sup>9</sup>C(S)O—, —NR<sup>9</sup>C(S)—NR<sup>9</sup>—, —NR<sup>9</sup>C(S)S—, —NR<sup>9</sup>C(S)—, —OC(O)NR<sup>9</sup>—, —SC(O)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —OC(S)NR<sup>9</sup>—, —SC(S)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —S(O)<sub>0-2</sub>—, —C(O)O—, —OC(O)—, —C(S)O—, —OC(S)—, —C(O)S—, —SC(O)—,

5

—C(S)S—, —SC(S)—, —OC(O)O—, —SC(O)O—,  
—OC(O)S—, —SC(S)O—, —OC(S)S—, —NR<sup>9</sup>C  
(NR<sup>2</sup>)NR<sup>9</sup>—, —NR<sup>9</sup>SO<sub>2</sub>—, —SO<sub>2</sub>NR<sup>9</sup>— and  
—NR<sup>9</sup>SO<sub>2</sub>NR<sup>9</sup>—,

each R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>10</sup> is independently selected from H,  
—(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Ar, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-(C<sub>0</sub>-C<sub>6</sub> alkyl),  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>9</sup>(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-  
O-(C<sub>0</sub>-C<sub>6</sub> alkyl) and —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>-(C<sub>0</sub>-C<sub>6</sub>  
alkyl),

each R<sup>9</sup> is independently selected from —H, —(C<sub>1</sub>-C<sub>4</sub>  
alkyl) and —C(O)—(C<sub>1</sub>-C<sub>4</sub> alkyl),

each Ar is an optionally substituted aryl,

each Het is an optionally substituted heteroaryl,

each Cak is an optionally substituted cycloalkyl,

each Hca is an optionally substituted heterocycloalkyl, and  
each alkyl is optionally substituted.

Another aspect of the invention is a method for activating  
the AMPK pathway in a cell, the method comprising contact-  
ing the cell with an effective amount of a compound, phar-  
maceutically acceptable salt, prodrug, solvate, hydrate or  
N-oxide or composition described above.

Another aspect of the invention is a method for increasing  
fatty acid oxidation in a cell, the method comprising contact-  
ing the cell with an effective amount of a compound, phar-  
maceutically acceptable salt, prodrug, solvate, hydrate or  
N-oxide or composition described above.

Another aspect of the invention is a method for decreasing  
glycogen concentration in a cell, the method comprising con-  
tacting the cell with an effective amount of a compound,  
pharmaceutically acceptable salt, prodrug, solvate, hydrate or  
N-oxide or composition described above.

Another aspect of the invention is a method for reducing  
triglyceride levels in a subject, the method comprising admin-  
istering to the subject an effective amount of a compound,  
pharmaceutically acceptable salt, prodrug, solvate, hydrate or  
N-oxide or composition described above.

Another aspect of the invention is a method for treating  
type II diabetes in a subject, the method comprising admin-  
istering to the subject an effective amount of a compound,  
pharmaceutically acceptable salt, prodrug, solvate, hydrate or  
N-oxide or composition described above.

Another aspect of the invention is a method for treating or  
preventing atherosclerosis or cardiovascular disease in a sub-  
ject, the method comprising administering to the subject an  
effective amount of a compound, pharmaceutically accept-  
able salt, prodrug, solvate, hydrate or N-oxide or composition  
described above.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is a plot of AMPK activity vs. concentration for both  
glutathione S-transferase and its fusion protein with globular  
adiponectin (gAd);

FIG. 2 is a plot of AMPK activity vs. concentration for both  
gAd and its polyhistidine-tagged analog;

FIG. 3 is a diagram of data for compounds 1-4 in a western  
blot assay for AMPK activity;


FIG. 4 is a set of plots showing reduction of glycogen  
content in HepG2 cells by compounds 1-4; and

FIG. 5 is a bar graph showing glucose uptake data for  
compounds 1-4 relative to adiponectin.

#### DETAILED DESCRIPTION OF THE INVENTION

A first aspect of the invention provides compounds having  
structural formula (I):

6



(I)

and pharmaceutically acceptable salts, solvates, hydrates, and  
N-oxides thereof, in which

R<sup>1</sup> is H;

R<sup>2</sup> is —Hca;

each R<sup>3</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl),  
—(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-  
C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen,  
—NO<sub>2</sub> and —CN;


w is 0, 1, 2, or 3;

n is 0, 1 2 or 3;

each R<sup>4</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl),  
—(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-  
C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen,  
—NO<sub>2</sub> and —CN, and two R<sup>4</sup> on the same carbon  
optionally combine to form oxo;

x is 0, 1, 2, 3 or 4;

T is —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>; or



in which

Q is —(C<sub>0</sub>-C<sub>3</sub> alkyl)-, in which each carbon of the —(C<sub>0</sub>-  
C<sub>3</sub> alkyl)- is optionally and independently substituted  
with one or two R<sup>16</sup>; each R<sup>16</sup> is independently selected  
from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-  
S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN; and

the ring system denoted by “A” is heteroaryl, aryl,  
cycloalkyl or heterocycloalkyl;

each R<sup>5</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl),  
—(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-  
C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen,  
—NO<sub>2</sub> and —CN; and

y is 0, 1, 2, 3 or 4;

in which

each L is independently selected from —NR<sup>9</sup>C(O)O—,  
—NR<sup>9</sup>C(O)—NR<sup>9</sup>—, —NR<sup>9</sup>C(O)S—, —NR<sup>9</sup>C(O)—,  
—NR<sup>9</sup>C(S)O—, —NR<sup>9</sup>C(S)—NR<sup>9</sup>—, —NR<sup>9</sup>C(S)  
S—, —NR<sup>9</sup>C(S)—, —OC(O)NR<sup>9</sup>—, —SC(O)NR<sup>9</sup>—,  
—C(S)NR<sup>9</sup>—, —OC(S)NR<sup>9</sup>—, —SC(S)NR<sup>9</sup>—,  
—C(S)NR<sup>9</sup>—, —S(O)<sub>0-2</sub>—, —C(O)O—, —OC(O)—,  
—C(S)O—, —OC(S)—, —C(O)S—, —SC(O)—,  
—C(S)S—, —SC(S)—, —OC(O)O—, —SC(O)O—,

7

—OC(O)S—, —SC(S)O—, —OC(S)S—, —NR<sup>9</sup>C  
(NR<sup>2</sup>)NR<sup>9</sup>—, —NR<sup>9</sup>SO<sub>2</sub>—, —SO<sub>2</sub>NR<sup>9</sup>— and  
—NR<sup>9</sup>SO<sub>2</sub>NR<sup>9</sup>—,

each R<sup>6</sup>, R<sup>7</sup>, R<sup>8</sup> and R<sup>10</sup> is independently selected from H,  
—(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub>  
alkyl)-Ar, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak,  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-(C<sub>0</sub>-C<sub>6</sub> alkyl),  
—(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>9</sup>(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-  
O-(C<sub>0</sub>-C<sub>6</sub> alkyl) and —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>-(C<sub>0</sub>-C<sub>6</sub>  
alkyl),

each R<sup>9</sup> is independently selected from —H, —(C<sub>1</sub>-C<sub>4</sub>  
alkyl) and —C(O)—(C<sub>1</sub>-C<sub>4</sub> alkyl),

each Ar is an optionally substituted aryl,

each Het is an optionally substituted heteroaryl,

each Cak is an optionally substituted cycloalkyl,

each Hca is an optionally substituted heterocycloalkyl, and  
each alkyl is optionally substituted,

provided that the compound is not

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-methoxybenzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-(furan-2-ylmethyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(pyridin-4-ylmethyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(3-phenylpropyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(methylsulfonyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide;


N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide; or

N-(1-benzylpiperidin-4-yl)-3-(1-benzylpiperidin-4-yloxy)benzamide.

In one embodiment according to the first aspect of the invention, the compound is a compound as described above with reference to structural formula (I), but is not 3-fluoro-N-(1-phenylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide; or -(1-(4-cyanobenzyl)piperidin-4-yloxy)-N-(piperidin-4-yl)benzamide.

In one embodiment according to the first aspect of the invention, two R<sup>4</sup>s combine to form an oxo. The oxo can be bound, for example, at the position alpha to the nitrogen of the azacycloalkyl.

In certain embodiments according to the first aspect of the invention, T is



In these embodiments of the invention, Q is —(C<sub>0</sub>-C<sub>3</sub> alkyl)—, in which each carbon of the (C<sub>0</sub>-C<sub>3</sub> alkyl) is optionally and independently substituted with one or two R<sup>16</sup>, in which each R<sup>16</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub>


8

alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>16</sup> on the same carbon optionally combine to form an oxo. Q can be, for example, an unsubstituted (C<sub>1</sub>-C<sub>3</sub> alkyl). In other embodiments of the invention, Q is a (C<sub>1</sub>-C<sub>3</sub> alkyl) having as its only substitution a single oxo group. For example, in certain embodiments of the invention Q is —CH<sub>2</sub>—; a single bond; or —C(O)— or —CH(CH<sub>3</sub>)—.

The number of substituents on the ring system denoted by “A”, y, in these embodiments is 0, 1, 2, 3 or 4. For example, in some embodiments of the invention, y is 0, 1, 2 or 3, for example 0, or 1. In one embodiment of the invention, y is not zero and at least one R<sup>5</sup> is halo, cyano, trifluoromethyl or trifluoromethoxy.

The ring system denoted by “A” is heteroaryl, aryl, cycloalkyl or heterocycloalkyl. For example, in one embodiment of the invention, the ring system denoted by “A” is an aryl or a heteroaryl. In one embodiment of the invention, the ring system denoted by “A” is an aryl. The ring system denoted by “A” can be, for example, a monocyclic aryl or heteroaryl.

For example, in one embodiment of the invention the ring system denoted by “A” is an aryl, such as a phenyl. In one embodiment of the invention, y is 1 and R<sup>5</sup> is attached to the phenyl para to Q. In another embodiment of the invention, y is 1 and R<sup>5</sup> is selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl, carboxylate, carboxamide and nitro. R<sup>5</sup> can be, for example, —Cl, —F, cyano, trifluoromethyl or trifluoromethoxy. In another embodiment of the invention, the




moiety is a 3,4-dihalophenyl.

In another embodiment of the invention, the ring system denoted by “A” is a heteroaryl. For example, in certain embodiments of the invention, the ring system denoted by “A” is a pyridinyl, a thienyl, or a furanyl.

In one embodiment according to the first aspect of the invention, the compound has structural formula (II):


(II)



in which the variables are defined as described above with reference to formula (I).

In another embodiment according to the first aspect of the invention, the compound has the structural formula (III):


9



(III)

in which the variables are defined as described above with reference to formula (I).


For example, compounds according to certain embodiments of the first aspect of the invention have structural formula (IV):



(IV)

in which the variables are defined as described above with reference to formula (I).

In other embodiments according to the first aspect of the invention, compounds of the present invention have structural formula (V):




(V)

in which the variables are defined as described above with reference to formula (I).

In certain embodiments of the invention, n is 1 or 2. For example, in one embodiment according to the first aspect of the invention, n is 2.

In one embodiment according to the first aspect of the invention, the compound has structural formula (VI):




(VI)

in which the variables are defined as described above with reference to formula (I).

In another embodiment according to the first aspect of the invention, the compound has the structural formula (VII):


10



(VII)

in which the variables are defined as described above with reference to formula (I).


For example, compounds according to the first aspect of the invention can have structural formula (VIII):



(VIII)

in which the variables are defined as described above with reference to formula (I).

In other embodiments according to the first aspect of the invention, compounds of the present invention have structural formula (IX):



(IX)

in which the variables are defined as described above with reference to formula (I).

According to the first aspect of the invention, R<sup>1</sup> is —H and R<sup>2</sup> is —Hca. In certain embodiments according to the first aspect of the invention, R<sup>2</sup> is substituted with (C<sub>0</sub>-C<sub>3</sub> alkyl)-Het or (C<sub>0</sub>-C<sub>3</sub> alkyl)-Ar. In one embodiment according to the first aspect of the invention, R<sup>2</sup> is -(optionally-substituted azetidiny), -(optionally-substituted pyrrolidiny), -(optionally-substituted piperidiny), or -(optionally-substituted azepany). For example, R<sup>2</sup> can be -(optionally substituted piperidiny) or -(optionally substituted pyrrolidiny).

In one embodiment according to the first aspect of the invention, R<sup>1</sup> is —H and R<sup>2</sup> is -(optionally-substituted azetid-3-yl), -(optionally substituted piperidin-4-yl), -(optionally substituted pyrrolidin-3-yl) or -(optionally-substituted azepan-4-yl). For example, in one embodiment according to the first aspect of the invention, R<sup>2</sup> is -(optionally substituted piperidin-4-yl). In another embodiment according to the first aspect of the invention, R<sup>2</sup> is -(optionally substituted pyrrolidin-3-yl).

In certain embodiments of the invention, the azetidiny, pyrrolidiny, piperidiny and azepany R<sup>2</sup> moieties described above are substituted at their 1-positions. For example, in one embodiment of the invention R<sup>2</sup> is substituted at its 1-position with (C<sub>0</sub>-C<sub>3</sub> alkyl)-Ar or (C<sub>0</sub>-C<sub>3</sub> alkyl)-Het. For example, in

## 11


one embodiment of the invention the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety is substituted at its 1-position with an optionally substituted benzyl or an optionally substituted phenyl. In another embodiment of the invention, the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety is substituted at its 1-position with a benzyl substituted with an electron withdrawing group; or with a pyridinylmethyl substituted with an electron withdrawing group. For example, the benzyl or pyridinylmethyl can be substituted with an electron withdrawing group selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl groups, carboxylate groups, carboxamide groups, cyano groups, sulfonate groups, and nitro groups. In other embodiments of the invention, the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety is substituted at its 1-position with an unsubstituted benzyl or an unsubstituted phenyl.

In other embodiments of the invention, the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety is substituted at its 1-position with an optionally substituted pyridinylmethyl, an optionally substituted furanylmethyl or an optionally substituted thienylmethyl. For example, the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety can be substituted with an unsubstituted pyridinylmethyl, an unsubstituted furanylmethyl, or an unsubstituted thienylmethyl.

In other embodiments of the invention, the azetidiny, pyrrolidiny, piperidiny or azepany R<sup>2</sup> moiety is substituted at its 1-position with —CO—O—(C<sub>0</sub>-C<sub>6</sub> alkyl), —CO—Het, —CO—Ar or —SO<sub>2</sub>—(C<sub>0</sub>-C<sub>6</sub> alkyl).

According to the first aspect of the invention, the number of substituents on the central phenyl ring, w, is 0, 1, 2, 3 or 4. For example, in one embodiment of the invention, w is 0, 1 or 2. In another embodiment according to the first aspect of the invention, w is 0. In other embodiments according to the first aspect of the invention, w is at least 1, and at least one R<sup>3</sup> is selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl, carboxylate, carboxamide and nitro.


In certain embodiments according to the first aspect of the invention, R<sup>3</sup> is selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl, carboxylate, carboxamide and nitro. R<sup>3</sup> can be, for example, —Cl or —F. For example, compounds according to these embodiments of the invention can have structural formula (X):



in which the remaining variables are defined as described above with reference to formula (I).

Certain other compounds according to these embodiments of the invention have structural formula (XI):


## 12



in which the remaining variables are defined as described above with reference to formula (I).

According to the first aspect of the invention, the number of substituents on the ethereal azacycloalkane ring, x, is 0, 1, 2, 3 or 4. In one embodiment according to the first aspect of the invention, x is 0, 1, 2 or 3. For example, x can be 0, or can be 1 or 2.

Compounds according to one embodiment of the first aspect of the invention have the structural formula (XII):




in which Q and G are each independently a bond, —CH<sub>2</sub>—, —C(H)(R<sup>16</sup>)— or —C(R<sup>16</sup>)<sub>2</sub>—; v is 0, 1, 2, 3 or 4; each R<sup>15</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl)-Ar; —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Het, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Cak, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-Hca, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN, and two R<sup>15</sup> on the same carbon optionally combine to form oxo; and R<sup>17</sup> is Het or Ar, and all other variables are defined as described above with reference to formula (I). In one embodiment of the invention, v is 0. In one embodiment of the invention, Q is a single bond. In another embodiment of the invention, G is —CH<sub>2</sub>— or —CO—. For example, in one embodiment of the invention, Q is a single bond and G is —CH<sub>2</sub>— or —CO—. The ethereal linkage of the piperidine to the benzamide can be at any aryl carbon. For example, the ether can be substituted at the 3-position or the 4-position of the benzamide. In one embodiment of the invention, two R<sup>15</sup>s combine to form an oxo, which can be bound, for example, at a position alpha to the piperidine nitrogen. As described above, in certain embodiments of the invention the ring system denoted by “A” is aryl or heteroaryl. In one embodiment of the invention, the ring system denoted by “A” is substituted with one or more electron-withdrawing groups. In another embodiment of the invention, R<sup>17</sup> is substituted with one or more electron-withdrawing groups.

Compounds according to certain embodiments of the first aspect of the invention have the structural formula (XIII):


13

14




in which Q is  $-\text{CH}_2-$  or a single bond;  $\text{R}^3$  is halo;  $\text{R}^H$  is H, halo, cyano, or a carboxylate; and  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

Compounds according to other embodiments according to the first aspect of the invention have structural formula (XIV):




in which Q is  $-\text{CH}_2-$  or a single bond;  $\text{R}^H$  is H, halo, cyano, or a carboxylate; and  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

Compounds according to certain embodiments according to the first aspect of the invention have the structural formula (XV):



in which Q is  $-\text{CH}_2-$  or a single bond;  $\text{R}^3$  is halo; and  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

Compounds according to other embodiments according to the first aspect of the invention have the structural formula (XVI):



in which Q is  $-\text{CH}_2-$  or a single bond; and  $\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

Examples of compounds of the present invention according to structural formula (I) include those listed below in Table 1. These compounds can be made according to the general scheme described below, for example using a procedure similar to that described below in Example 1.

65

TABLE 1

Cpd Name	Structure
14 N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-chlorobenzamide	
15 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-((2,3-dihydrobenzo[b][1,4]dioxin-6-yl)methyl)piperidin-4-yloxy)benzamide	
16 N-(1-benzylpiperidin-4-yl)-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-chlorobenzamide	
17 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
18 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)benzoyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
19 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-fluorobenzyl)piperidin-4-yloxy)benzamide	
20 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)benzamide	
21 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(1-phenylethyl)piperidin-4-yloxy)benzamide	
22 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-fluorobenzoyl)piperidin-4-yloxy)benzamide	
23 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
24 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(2-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
25 3-fluoro-N-(1-phenylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
26 tert-butyl 4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamido)piperidine-1-carboxylate	
27 3-fluoro-N-(1-(4-fluorobenzyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
28 3-fluoro-N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
29 3-fluoro-N-(1-(pyridin-3-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd	Name	Structure
30	3-fluoro-N-(1-pivaloylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
31	3-fluoro-N-(1-(4-fluorobenzoyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
32	3-fluoro-N-(1-(pyridin-2-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
33	methyl 4-((4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamido)piperidin-1-yl)methyl)benzoate	
34	3-fluoro-N-(1-(isopropylsulfonyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
35 4-((4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamido)piperidin-1-yl)methyl)benzoic acid	
36 4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-fluoro-N-(1-phenylpiperidin-4-yl)benzamide	
37 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-4-ylmethyl)piperidin-4-yloxy)benzamide	
38 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-3-ylmethyl)piperidin-4-yloxy)benzamide	
39 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-2-ylmethyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
40 3-fluoro-N-(1-isonicotinoylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
41 N-(1-benzylpiperidin-4-yl)-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
42 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide	
43 N-(1-(4-cyanobenzyl)piperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
44 4-(1-(4-cyanophenyl)piperidin-4-yloxy)-3-fluoro-N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)benzamide	
45 N-(1-(4-cyanobenzyl)piperidin-4-yl)-4-(1-(4-cyanophenyl)piperidin-4-yloxy)-3-fluorobenzamide	

TABLE 1-continued

Cpd Name	Structure
46 N-(1-benzylpiperidin-4-yl)-3-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
47 N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
48 N-(1-benzylpiperidin-4-yl)-3-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide	
49 N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide	
50 N-(1-benzylpiperidin-4-yl)-3-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide	
51 N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide	
52 N-(1-benzylpiperidin-4-yl)-3-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
53 N-(1-benzylpiperidin-4-yl)-3-(1-(pyridin-4-yl)piperidin-4-yloxy)benzamide	
54 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide	
55 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
56 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
57 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
58 3-chloro-N-(1-methylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
59 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
60 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
61 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(fluorobenzyl)piperidin-4-yloxy)benzamide	
62 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide	
63 N-(1-benzylpiperidin-4-yl)-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
64 N-(1-benzylpiperidin-4-yl)-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
65 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)benzamide	
66 3-chloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-N-(1-(4-fluorobenzyl)piperidin-4-yl)benzamide	
67 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide	
68 3-chloro-N-(1-(4-fluorobenzyl)piperidin-4-yl)-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
69 N-(1-benzylpiperidin-4-yl)-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
70 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
71 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
72 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
73 N-(1-(4-chlorobenzyl)piperidin-4-yl)-3-fluoro-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide	

TABLE 1-continued

Cpd Name	Structure
74 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)benzamide	
75 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)benzamide	
76 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)benzamide	
77 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide	
78 tert-butyl 4-(4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamido)piperidine-1-carboxylate	

TABLE 1-continued

Cpd Name	Structure
79 4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-N-(piperidin-4-yl)benzamide	
80 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
81 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(4-cyanophenyl)piperidin-4-yloxy)-3-fluorobenzamide	
82 N-(1-benzylpiperidin-4-yl)-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide	
83 4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-N-(1-(piperidin-4-ylmethyl)piperidin-4-yl)benzamide	

TABLE 1-continued

Cpd Name	Structure
84 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)benzamide	
85 4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-N-(1-(pyridin-2-ylmethyl)piperidin-4-yl)benzamide	
86 N-(1-benzylpiperidin-4-yl)-3-(1-(pyridin-2-yl)piperidin-4-yloxy)benzamide	
87 N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-3-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide	
88 N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-3-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide	
89 N-(1-benzylpiperidin-4-yl)-3-(1-(3-cyanobenzyl)piperidin-4-yloxy)benzamide	
90 tert-butyl 4-(4-(1-benzylpiperidin-4-ylcarbamoyl)-2-chlorophenoxy)piperidin-1-carboxylate	

TABLE 1-continued

Cpd Name	Structure
91 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-pivaloylpiperidin-4-yloxy)benzamide	
92 tert-butyl 4-(4-(1-benzylpiperidin-4-ylcarbamoyl)-2-fluorophenoxy)piperidine-1-carboxylate	
93 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(piperidin-4-yloxy)benzamide	

For simplicity, chemical moieties are defined and referred to throughout primarily as univalent chemical moieties (e.g., alkyl, aryl, etc.). Nevertheless, such terms are also used to convey corresponding multivalent moieties under the appropriate structural circumstances clear to those skilled in the art. For example, while an “alkyl” moiety generally refers to a monovalent radical (e.g.  $\text{CH}_3\text{—CH}_2\text{—}$ ), in certain circumstances a bivalent linking moiety can be “alkyl,” in which case those skilled in the art will understand the alkyl to be a divalent radical (e.g.,  $\text{—CH}_2\text{—CH}_2\text{—}$ ), which is equivalent to the term “alkylene.” (Similarly, in circumstances in which a divalent moiety is required and is stated as being “aryl,” those skilled in the art will understand that the term “aryl” refers to the corresponding divalent moiety, arylene). All atoms are understood to have their normal number of valences for bond formation (i.e., 4 for carbon, 3 for N, 2 for O, and 2, 4, or 6 for S, depending on the oxidation state of the S). Nitrogens in compounds of the invention can be hypervalent, e.g., an N-oxide or tetrasubstituted ammonium salt. On occasion a moiety may be defined, for example, as (A)<sub>a</sub>-B-, wherein a is 0 or 1. In such instances, when a is 0 the moiety is B- and when a is 1 the moiety is A-B-.

As used herein, the term “alkyl” includes alkyl, alkenyl and alkynyl groups of a designed number of carbon atoms, desirably from 1 to about 12 carbons (i.e., inclusive of 1 and 12). The term “C<sub>m</sub>-C<sub>n</sub> alkyl” means an alkyl group having from m to n carbon atoms (i.e., inclusive of m and n). The term “C<sub>m</sub>-C<sub>n</sub> alkyl” means an alkyl group having from m to n carbon atoms. For example, “C<sub>1</sub>-C<sub>6</sub> alkyl” is an alkyl group having from one to six carbon atoms. Alkyl and alkyl groups may be straight or branched and depending on context, may be a monovalent radical or a divalent radical (i.e., an alkylene group). In the case of an alkyl or alkyl group having zero carbon atoms (i.e., “C<sub>0</sub> alkyl”), the group is simply a single covalent bond if it is a divalent radical or is a hydrogen atom if it is a monovalent radical. For example, the moiety “—(C<sub>0</sub>-C<sub>6</sub> alkyl)-Ar” signifies connection of an optionally substituted aryl through a single bond or an alkylene bridge having from 1 to 6 carbons. Examples of “alkyl” include, for example, methyl, ethyl, propyl, isopropyl, butyl, iso-, sec- and tert-butyl, pentyl, hexyl, heptyl, 3-ethylbutyl, 3-hexenyl

and propargyl. If the number of carbon atoms is not specified, the subject “alkyl” or “alkyl” moiety has from 1 to 12 carbons.

The term “fluoroalkyl” is an alkyl group substituted with one or more fluorine atoms. Examples of “fluoroalkyl” include fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, hexafluoroisopropyl and the like.

The term “aryl” represents an aromatic carbocyclic ring system having a single ring (e.g., phenyl) which is optionally fused to other aromatic hydrocarbon rings or non-aromatic hydrocarbon rings. “Aryl” includes ring systems having multiple condensed rings and in which at least one is aromatic, (e.g., 1,2,3,4-tetrahydronaphthyl, naphthyl). Examples of aryl groups include phenyl, 1-naphthyl, 2-naphthyl, indenyl, indanyl, dihydronaphthyl, fluorenyl, tetralinyl, 2,3-dihydrobenzofuranyl and 6,7,8,9-tetrahydro-5H-benzo[a]cycloheptenyl. The aryl groups herein are unsubstituted or, when specified as “optionally substituted”, can unless stated otherwise be substituted in one or more substitutable positions with various groups, as described below.

The term “heteroaryl” refers to an aromatic ring system containing at least one heteroatom selected from nitrogen, oxygen and sulfur in an aromatic ring. The heteroaryl may be fused to one or more cycloalkyl or heterocycloalkyl rings. Examples of heteroaryl groups include, for example, pyridyl, pyrimidinyl, quinolinyl, benzothienyl, indolyl, indolinyl, pyridazinyl, pyrazinyl, isoindolyl, isoquinolyl, quinazoliny, quinoxaliny, phthalazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, indoliziny, indazolyl, benzothiazolyl, benzimidazolyl, benzofuranyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, benzo[1,4]oxazinyl, triazolyl, tetrazolyl, isothiazolyl, naphthyridinyl, isochromanyl, chromanyl, tetrahydroisoquinoliny, isoindoliny, isobenzotetrahydrofuranly, isobenzotetrahydrothienyl, isobenzothienyl, benzoxazolyl, pyridopyridinyl, benzotetrahydrofuranly, benzotetrahydrothienyl, purinyl, benzodioxolyl, triazinyl, pteridinyl, benzothiazolyl, imidazopyridinyl, imidazothiazolyl, dihydrobenzisoxazinyl, benzisoxazinyl, benzoxazinyl, dihydrobenzisothiazinyl, benzopyranyl, benzothioapyranyl, chromonyl, chromanonyl, pyridinyl-N-oxide, tetrahydroquinolinyl, dihydroquinolinyl, dihydroquinolinonyl, dihydroisoquinolinonyl, dihydrocoumarinyl, dihydroisocoumarin-

nyl, isoindolinonyl, benzodioxanyl, benzoxazolinonyl, pyrrolyl N-oxide, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, quinolinyl N-oxide, indolyl N-oxide, indolinyl N-oxide, isoquinolyl N-oxide, quinazoliny N-oxide, quinoxaliny N-oxide, phthalazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, indoliziny N-oxide, indazolyl N-oxide, benzothiazolyl N-oxide, benzimidazolyl N-oxide, pyrrolyl N-oxide, oxadiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, tetrazolyl N-oxide, benzothiopyranyl S-oxide, benzothiopyranyl S,S-dioxide. Preferred heteroaryl groups include pyridyl, pyrimidyl, quinolinyl, indolyl, pyrrolyl, furanyl, thienyl and imidazolyl, pyrazolyl, indazolyl, thiazolyl and benzothiazolyl. In certain embodiments of the invention, each heteroaryl is selected from pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, imidazolyl, isoxazolyl, pyrazolyl, oxazolyl, thiazolyl, furanyl, thienyl, pyrrolyl, oxadiazolyl, thiadiazolyl, triazolyl, tetrazolyl, isothiazolyl, pyridinyl-N-oxide, pyrrolyl N-oxide, pyrimidinyl N-oxide, pyridazinyl N-oxide, pyrazinyl N-oxide, imidazolyl N-oxide, isoxazolyl N-oxide, oxazolyl N-oxide, thiazolyl N-oxide, thiadiazolyl N-oxide, triazolyl N-oxide, and tetrazolyl N-oxide. Preferred heteroaryl groups include pyridyl, pyrimidyl, quinolinyl, indolyl, pyrrolyl, furanyl, thienyl, imidazolyl, pyrazolyl, indazolyl, thiazolyl and benzothiazolyl. The heteroaryl groups herein are unsubstituted or, when specified as "optionally substituted", can unless stated otherwise be substituted in one or more substitutable positions with various groups, as described below.

The term "heterocycloalkyl" refers to a non-aromatic ring or ring system containing at least one heteroatom that is preferably selected from nitrogen, oxygen and sulfur, wherein said heteroatom is in a non-aromatic ring. The heterocycloalkyl may be saturated (i.e., a heterocycloalkyl) or partially unsaturated (i.e., a heterocycloalkenyl). The heterocycloalkyl ring is optionally fused to other heterocycloalkyl rings and/or non-aromatic hydrocarbon rings and/or phenyl rings. In certain embodiments of the invention, the heterocycloalkyl groups have from 3 to 7 members in a single ring. In other embodiments of the invention, heterocycloalkyl groups have 5 or 6 members in a single ring. Examples of heterocycloalkyl groups include, for example, azabicyclo[2.2.2]octyl (in each case also "quinuclidinyl" or a quinuclidine derivative), azabicyclo[3.2.1]octyl, morpholinyl, thiomorpholinyl, thiomorpholinyl S-oxide, thiomorpholinyl S,S-dioxide, 2-oxazolidonyl, piperazinyl, homopiperazinyl, piperazinonyl, pyrrolidinyl, azepanyl, azetidiny, pyrrolinyl, tetrahydropyranyl, piperidinyl, tetrahydrofuranyl, tetrahydrothienyl, 3,4-dihydroisoquinolin-2(1H)-yl, isoindolindionyl, homopiperidinyl, homomorpholinyl, homothiomorpholinyl, homothiomorpholinyl S,S-dioxide, oxazolidinonyl, dihydropyrazolyl, dihydropyrrolyl, dihydropyrazinyl, dihydropyridinyl, dihydropyrimidinyl, dihydrofuryl, dihydropyranyl, imidazolidonyl, tetrahydrothienyl S-oxide, tetrahydrothienyl S,S-dioxide and homothiomorpholinyl S-oxide. Especially desirable heterocycloalkyl groups include morpholinyl, 3,4-dihydroisoquinolin-2(1H)-yl, tetrahydropyranyl, piperidinyl, aza-bicyclo[2.2.2]octyl,  $\gamma$ -butyrolactonyl (i.e., an oxo-substituted tetrahydrofuranyl),  $\gamma$ -butyrolactamyl (i.e., an oxo-substituted pyrrolidine), pyrrolidinyl, piperazinyl, azepanyl, azetidiny, thiomorpholinyl, thiomorpholinyl S,S-dioxide, 2-oxazolidonyl, imidazolidonyl, isoindolindionyl, piperazinonyl. The heterocycloalkyl groups herein are unsubstituted or, when specified as "optionally substituted", can unless stated otherwise be substituted in one or more substitutable positions with various groups, as described below.

The term "cycloalkyl" refers to a non-aromatic carbocyclic ring or ring system, which may be saturated (i.e., a cycloalkyl) or partially unsaturated (i.e., a cycloalkenyl). The cycloalkyl ring optionally fused to or otherwise attached (e.g., bridged systems) to other cycloalkyl rings. Preferred cycloalkyl groups have from 3 to 7 members in a single ring. More preferred cycloalkyl groups have 5 or 6 members in a single ring. Examples of cycloalkyl groups include, for example, cyclohexyl, cyclopentyl, cyclobutyl, cyclopropyl, tetrahydronaphthyl and bicyclo[2.2.1]heptane. The cycloalkyl groups herein are unsubstituted or, when specified as "optionally substituted", may be substituted in one or more substitutable positions with various groups.

The term "oxa" means a divalent oxygen radical in a chain, sometimes designated as —O—.

The term "oxo" means a doubly bonded oxygen, sometimes designated as =O or for example in describing a carbonyl "C(O)" may be used to show an oxo substituted carbon.

The term "electron withdrawing group" means a group that withdraws electron density from the structure to which it is attached than would a similarly-attached hydrogen atom. For example, electron withdrawing groups can be selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl groups (e.g., —C(O)—H, —C(O)-alkyl), carboxylate groups (e.g., carboxylic acids and esters), carboxamide groups, cyano groups, sulfonate groups (including sulfonic acid and sulfonic esters), and nitro groups.

The term "substituted," when used to modify a specified group or radical, means that one or more hydrogen atoms of the specified group or radical are each, independently of one another, replaced with the same or different substituent groups as defined below.

Substituent groups for substituting for hydrogens on saturated carbon atoms in the specified group or radical are, unless otherwise specified, —R<sup>60</sup>, halo, —O<sup>-</sup>M<sup>+</sup>, =O, —OR<sup>70</sup>, —SR<sup>70</sup>, —S<sup>-</sup>M<sup>+</sup>, =S, —NR<sup>80</sup>R<sup>80</sup>, =NR<sup>70</sup>, =N—OR<sup>70</sup>, trihalomethyl, —CF<sub>3</sub>, —CN, —OCN, —SCN, —NO, —NO<sub>2</sub>, —N<sub>2</sub>, —N<sub>3</sub>, —SO<sub>2</sub>R<sup>70</sup>, —SO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —SO<sub>2</sub>OR<sup>70</sup>, —OSO<sub>2</sub>R<sup>70</sup>, —OSO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, OSO<sub>2</sub>OR<sup>70</sup>, —P(O)(O<sup>-</sup>)<sub>2</sub>(M<sup>+</sup>)<sub>2</sub>, —P(O)(OR<sup>70</sup>)O<sup>-</sup>M<sup>+</sup>, —P(O)(OR<sup>70</sup>)<sub>2</sub>, —C(O)R<sup>70</sup>, —C(S)R<sup>70</sup>, —C(NR<sup>70</sup>)R<sup>70</sup>, —C(O)O<sup>-</sup>M<sup>+</sup>, —C(O)OR<sup>70</sup>, —C(S)OR<sup>70</sup>, —C(O)NR<sup>80</sup>R<sup>80</sup>, —C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>, —OC(O)R<sup>70</sup>, —OC(S)R<sup>70</sup>, —OC(O)O<sup>-</sup>M<sup>+</sup>, —OC(O)OR<sup>70</sup>, —OC(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)R<sup>70</sup>, —NR<sup>70</sup>C(S)R<sup>70</sup>, —NR<sup>70</sup>CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —NR<sup>70</sup>CO<sub>2</sub>R<sup>70</sup>, —NR<sup>70</sup>C(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)NR<sup>80</sup>R<sup>80</sup>, —NR<sup>70</sup>C(NR<sup>70</sup>)R<sup>70</sup> and —NR<sup>70</sup>C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>. Each R<sup>60</sup> is independently selected from the group consisting of alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, heterocycloalkylalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl and heteroarylalkyl, each of which is optionally substituted with 1, 2, 3, 4 or 5 groups selected from the group consisting of halo, —O<sup>-</sup>M<sup>+</sup>, =O, —OR<sup>71</sup>, —SR<sup>71</sup>, —S<sup>-</sup>M<sup>+</sup>, =S, —NR<sup>81</sup>R<sup>81</sup>, =NR<sup>71</sup>, =N—OR<sup>71</sup>, trihalomethyl, —CF<sub>3</sub>, —CN, —OCN, —SCN, —NO, —NO<sub>2</sub>, =N<sub>2</sub>, —N<sub>3</sub>, —SO<sub>2</sub>R<sup>71</sup>, —SO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —SO<sub>2</sub>OR<sup>71</sup>, —OSO<sub>2</sub>R<sup>71</sup>, —OSO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, OSO<sub>2</sub>OR<sup>71</sup>, —P(O)(O<sup>-</sup>)<sub>2</sub>(M<sup>+</sup>)<sub>2</sub>, —P(O)(OR<sup>71</sup>)O<sup>-</sup>M<sup>+</sup>, —P(O)(OR<sup>71</sup>)<sub>2</sub>, —C(O)R<sup>71</sup>, —C(S)R<sup>71</sup>, —C(NR<sup>71</sup>)R<sup>71</sup>, —C(O)O<sup>-</sup>M<sup>+</sup>, —C(O)OR<sup>71</sup>, —C(S)OR<sup>71</sup>, —C(O)NR<sup>81</sup>R<sup>81</sup>, —C(NR<sup>71</sup>)NR<sup>81</sup>R<sup>81</sup>, —OC(O)R<sup>71</sup>, —OC(S)R<sup>71</sup>, —OC(O)O<sup>-</sup>M<sup>+</sup>, —OC(O)OR<sup>71</sup>, —OC(S)OR<sup>71</sup>, —NR<sup>71</sup>C(O)R<sup>71</sup>, —NR<sup>71</sup>C(S)R<sup>71</sup>, —NR<sup>71</sup>CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —NR<sup>71</sup>CO<sub>2</sub>R<sup>71</sup>, —NR<sup>71</sup>C(S)OR<sup>71</sup>, —NR<sup>71</sup>C(O)NR<sup>81</sup>R<sup>81</sup>, —NR<sup>71</sup>C(NR<sup>71</sup>)R<sup>71</sup> and —NR<sup>71</sup>C(NR<sup>71</sup>)NR<sup>81</sup>R<sup>81</sup>. Each R<sup>70</sup> is independently hydrogen or R<sup>60</sup>; each R<sup>80</sup> is independently R<sup>70</sup> or alternatively, two R<sup>80</sup>'s, taken together with the nitrogen atom to which they are

47

bonded, form a 5-, 6- or 7-membered heterocycloalkyl which may optionally include from 1 to 4 of the same or different additional heteroatoms selected from the group consisting of O, N and S, of which N may have —H or C<sub>1</sub>-C<sub>3</sub> alkyl substitution; and each M is a counter ion with a net single positive charge. Each R<sup>71</sup> is independently hydrogen or R<sup>61</sup>, in which R<sup>61</sup> is alkyl, heteroalkyl, cycloalkyl, heterocycloalkyl, heterocycloalkylalkyl, cycloalkylalkyl, aryl, arylalkyl, heteroaryl and heteroarylalkyl, each of which is optionally substituted with 1, 2, 3, 4 or 5 groups selected from the group consisting of halo, —O<sup>-</sup>M<sup>+</sup>, =O, —OR<sup>72</sup>, —SR<sup>72</sup>, —S<sup>-</sup>M<sup>+</sup>, =S, —NR<sup>82</sup>R<sup>82</sup>, —NR<sup>72</sup>, —N—OR<sup>72</sup>, trihalomethyl, —CF<sub>3</sub>, —OCN, —SCN, —NO, —NO<sub>2</sub>, =N<sub>2</sub>, —N<sub>3</sub>, —SO<sub>2</sub>R<sup>71</sup>, —SO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —SO<sub>2</sub>OR<sup>72</sup>, —OSO<sub>2</sub>R<sup>72</sup>, —OSO<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —OSO<sub>2</sub>OR<sup>72</sup>, —P(O)(O<sup>-</sup>)<sub>2</sub>(M<sup>+</sup>), —P(O)(OR<sup>72</sup>)O<sup>-</sup>M<sup>+</sup>, —P(O)(OR<sup>72</sup>)<sub>2</sub>, —C(O)R<sup>72</sup>, —C(S)R<sup>72</sup>, —C(NR<sup>72</sup>)R<sup>72</sup>, —C(O)O<sup>-</sup>M<sup>+</sup>-C(O)OR<sup>72</sup>, —C(S)OR<sup>72</sup>, —C(O)NR<sup>82</sup>R<sup>82</sup>, —C(NR<sup>72</sup>)NR<sup>82</sup>R<sup>82</sup>, —OC(O)R<sup>72</sup>, —OC(S)R<sup>72</sup>, —OC(O)O<sup>-</sup>M<sup>+</sup>, —OC(O)OR<sup>72</sup>, —OC(S)OR<sup>72</sup>, —NR<sup>72</sup>C(O)R<sup>72</sup>, —NR<sup>72</sup>C(S)R<sup>72</sup>, —NR<sup>72</sup>CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —NR<sup>72</sup>CO<sub>2</sub>R<sup>72</sup>, —NR<sup>72</sup>C(S)OR<sup>72</sup>, —NR<sup>72</sup>C(O)NR<sup>82</sup>R<sup>82</sup>, —NR<sup>72</sup>C(NR<sup>72</sup>)R<sup>72</sup> and —NR<sup>72</sup>C(NR<sup>72</sup>)NR<sup>82</sup>R<sup>82</sup>; and each R<sup>81</sup> is independently R<sup>71</sup> or alternatively, two R<sup>81</sup>s, taken together with the nitrogen atom to which they are bonded, form a 5-, 6- or 7-membered heterocycloalkyl which may optionally include from 1 to 4 of the same or different additional heteroatoms selected from the group consisting of O, N and S, of which N may have —H or C<sub>1</sub>-C<sub>3</sub> alkyl substitution. Each M<sup>+</sup> may independently be, for example, an alkali ion, such as K<sup>+</sup>, Na<sup>+</sup>, Li<sup>+</sup>; an ammonium ion, such as or an alkaline earth ion, such as [Ca<sup>2+</sup>]<sub>0.5</sub>, [Mg<sup>2+</sup>]<sub>0.5</sub>, or [Ba<sup>2+</sup>]<sub>0.5</sub> ("subscript 0.5 means e.g. that one of the counter ions for such divalent alkali earth ions can be an ionized form of a compound of the invention and the other a typical counter ion such as chloride, or two ionized compounds of the invention can serve as counter ions for such divalent alkali earth ions, or a doubly ionized compound of the invention can serve as the counter ion for such divalent alkali earth ions). As specific examples, —NR<sup>80</sup>R<sup>80</sup> is meant to include —NH<sub>2</sub>, —NH-alkyl, N-pyrrolidinyl, N-piperazinyl, 4-methyl-piperazin-1-yl and N-morpholinyl.

Substituent groups for hydrogens on unsaturated carbon atoms in "substituted" alkene, alkyne, aryl and heteroaryl groups are, unless otherwise specified, —R<sup>60</sup>, halo, —O<sup>-</sup>M<sup>+</sup>, —OR<sup>70</sup>, —SR<sup>70</sup>, —S<sup>-</sup>M<sup>+</sup>, —NR<sup>80</sup>R<sup>80</sup>, trihalomethyl, —CF<sub>3</sub>, —CN, —OCN, —SCN, —NO, —NO<sub>2</sub>, —N<sub>3</sub>, —SO<sub>2</sub>R<sup>70</sup>, —SO<sub>3</sub><sup>-</sup>M<sup>+</sup>, —SO<sub>3</sub>R<sup>70</sup>, —OSO<sub>2</sub>R<sup>70</sup>, —OSO<sub>3</sub><sup>-</sup>M<sup>+</sup>, —OSO<sub>3</sub>R<sup>70</sup>, —PO<sub>3</sub><sup>-2</sup>(M<sup>+</sup>)<sub>2</sub>, —P(O)(OR<sup>70</sup>)O<sup>-</sup>M<sup>+</sup>, —P(O)(OR<sup>70</sup>)<sub>2</sub>, —C(O)R<sup>70</sup>, —C(S)R<sup>70</sup>, —C(NR<sup>70</sup>)R<sup>70</sup>, —CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —CO<sub>2</sub>R<sup>70</sup>, —C(S)OR<sup>70</sup>, —C(O)NR<sup>80</sup>R<sup>80</sup>, —C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>, —OC(O)R<sup>70</sup>, —OC(S)R<sup>70</sup>, —CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —OCO<sub>2</sub>R<sup>70</sup>, —OC(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)R<sup>70</sup>, —NR<sup>70</sup>C(S)R<sup>70</sup>, —NR<sup>70</sup>CO<sub>2</sub><sup>-</sup>M<sup>+</sup>, —NR<sup>70</sup>CO<sub>2</sub>R<sup>70</sup>, —NR<sup>70</sup>C(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)NR<sup>80</sup>R<sup>80</sup>, —NR<sup>70</sup>C(NR<sup>70</sup>)R<sup>70</sup> and —NR<sup>70</sup>C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>, where R<sup>60</sup>, R<sup>70</sup>, R<sup>80</sup> and M<sup>+</sup> are as previously defined.

Substituent groups for hydrogens on nitrogen atoms in "substituted" heteroalkyl and cycloheteroalkyl groups are, unless otherwise specified, —R<sup>60</sup>, —O<sup>-</sup>M<sup>+</sup>, —OR<sup>70</sup>, —SR<sup>70</sup>, —S<sup>-</sup>M<sup>+</sup>, —NR<sup>80</sup>R<sup>80</sup>, trihalomethyl, —CF<sub>3</sub>, —CN,

48

—NO, —NO<sub>2</sub>, —S(O)<sub>2</sub>R<sup>70</sup>, —S(O)<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —S(O)<sub>2</sub>OR<sup>70</sup>, —OS(O)<sub>2</sub>R<sup>70</sup>, —OS(O)<sub>2</sub>O<sup>-</sup>M<sup>+</sup>, —OS(O)<sub>2</sub>OR<sup>70</sup>, —P(O)(O<sup>-</sup>)<sub>2</sub>(M<sup>+</sup>)<sub>2</sub>, —P(O)(OR<sup>70</sup>)O<sup>-</sup>M<sup>+</sup>, —P(O)(OR<sup>70</sup>)(OR<sup>70</sup>), —C(O)R<sup>70</sup>, —C(S)R<sup>70</sup>, —C(NR<sup>70</sup>)R<sup>70</sup>, —C(O)OR<sup>70</sup>, —C(S)OR<sup>70</sup>, —C(O)NR<sup>80</sup>R<sup>80</sup>, —C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>, —OC(O)R<sup>70</sup>, —OC(S)R<sup>70</sup>, —OC(O)OR<sup>70</sup>, —OC(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)R<sup>70</sup>, —NR<sup>70</sup>C(S)R<sup>70</sup>, —NR<sup>70</sup>C(O)OR<sup>70</sup>, —NR<sup>70</sup>C(S)OR<sup>70</sup>, —NR<sup>70</sup>C(O)NR<sup>80</sup>R<sup>80</sup>, —NR<sup>70</sup>C(NR<sup>70</sup>)R<sup>70</sup> and —NR<sup>70</sup>C(NR<sup>70</sup>)NR<sup>80</sup>R<sup>80</sup>, where R<sup>60</sup>, R<sup>70</sup>, R<sup>80</sup> and M<sup>+</sup> are as previously defined.

In a preferred embodiment, a group that is substituted has 1, 2, 3, or 4 substituents, 1, 2, or 3 substituents, 1 or 2 substituents, or 1 substituent.


The term "pharmaceutically acceptable salts" or "a pharmaceutically acceptable salt thereof" refer to salts prepared from pharmaceutically acceptable non-toxic acids or bases including inorganic acids and bases and organic acids and bases. If the compound used in the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids. Such salts may be, for example, acid addition salts of at least one of the following acids: benzenesulfonic acid, citric acid, α-glucoheptonic acid, D-gluconic acid, glycolic acid, lactic acid, malic acid, malonic acid, mandelic acid, phosphoric acid, propanoic acid, succinic acid, sulfuric acid, tartaric acid (d, l, or dl), tosic acid (toluenesulfonic acid), valeric acid, palmitic acid, pamoic acid, sebacic acid, stearic acid, lauric acid, acetic acid, adipic acid, carbonic acid, 4-chlorobenzenesulfonic acid, ethanedisulfonic acid, ethylsuccinic acid, fumaric acid, galactaric acid (mucic acid), D-glucuronic acid, 2-oxo-glutaric acid, glycerophosphoric acid, hippuric acid, isethionic acid (ethanolsulfonic acid), lactobionic acid, maleic acid, 1,5-naphthalene-disulfonic acid, 2-naphthalene-sulfonic acid, pivalic acid, terephthalic acid, thiocyanic acid, cholic acid, n-dodecyl sulfate, 3-hydroxy-2-naphthoic acid, 1-hydroxy-2-naphthoic acid, oleic acid, undecylenic acid, ascorbic acid, (+)-camphoric acid, d-camphorsulfonic acid, dichloroacetic acid, ethanesulfonic acid, formic acid, hydriodic acid, hydrobromic acid, hydrochloric acid, methanesulfonic acid, nicotinic acid, nitric acid, orotic acid, oxalic acid, picric acid, L-pyroglytamic acid, saccharine, salicylic acid, gentisic acid, and/or 4-acetamidobenzoic acid.

"Prodrug" refers to a derivative of an active compound (drug) that requires a transformation under the conditions of use, such as within the body, to release the active drug. Prodrugs are frequently, but not necessarily, pharmacologically inactive until converted into the active drug. Prodrugs are typically obtained by masking a functional group in the drug believed to be in part required for activity with a progroup (defined below) to form a promoiety which undergoes a transformation, such as cleavage, under the specified conditions of use to release the functional group, and hence the active drug. The cleavage of the promoiety can proceed spontaneously, such as by way of a hydrolysis reaction, or it can be catalyzed or induced by another agent, such as by an enzyme, by light, by acid, or by a change of or exposure to a physical or environmental parameter, such as a change of temperature. The agent can be endogenous to the conditions of use, such as an enzyme present in the cells to which the prodrug is administered or the acidic conditions of the stomach, or it can be supplied exogenously. A wide variety of progroups, as well as the resultant promoieties, suitable for masking functional groups in the active drugs to yield prodrugs are well-known in the art. For example, a hydroxyl functional group can be masked as a sulfonate, ester or carbonate promoiety, which can be hydrolyzed in vivo to provide the hydroxyl group. An amino functional group can be masked as an amide, carbam-

49

ate, imine, urea, phosphanyl, phosphoryl or sulfonyl promoieties, which can be hydrolyzed *in vivo* to provide the amino group. A carboxyl group can be masked as an ester (including silyl esters and thioesters), amide or hydrazide promoieties, which can be hydrolyzed *in vivo* to provide the carboxyl group. Other specific examples of suitable progroups and their respective promoieties will be apparent to those of skill in the art.

The compounds of the present invention can be made using procedures familiar to the person of ordinary skill in the art and as described herein. For example, compounds of structural formulae (I)-(XVI) can be prepared according to Scheme 1, below:



Referring to Scheme 1, bromides 1, for example, can be coupled with 4-hydroxypiperidines 2 to provide 1-substituted 4-hydroxypiperidine 3. Substituted 4-hydroxypiperidine 3 can be subjected to Mitsunobu conditions, e.g., with appropriate phenols to give coupled products 4. The ester group of 4 is saponified, for example, converted to the corresponding carboxylic acid and coupled with an appropriate amine to yield amine to give compounds of structural formula (I).

One of ordinary skill in the art can adapt the reaction sequence of Scheme 1 to fit the desired target molecule. For example, a benzyl bromide can be used as a starting material to afford compounds of structural formula (I) in which the ring system denoted by "A" is a phenyl. Similarly, a (heteroaryl)methyl bromide may be used as a starting material to afford compounds of structural formula (I) in which the ring system denoted by "A" is a heteroaryl. Alternatively, reductive amination of an aryl or heteroaryl aldehyde, for example, with the nitrogen of azacycloalkyl 2 would also afford 3. In

50

certain situations one of ordinary skill in the art will use different reagents to effect one or more of the individual steps or to use protected versions of certain of the  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$  and  $R^5$  substituents. An example of the synthesis of a compound of the present invention is provided below in Example 1.

The compounds of structural formulae (I)-(XVI) may be administered, for example, orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing one or more pharmaceutically acceptable carriers, diluents or excipients. The term parenteral as used herein includes percutaneous, subcutaneous, intravascular (e.g., intravenous), intramuscular, or intrathecal injection or infusion techniques and the like.

Another aspect of the invention is a pharmaceutical formulation comprising a pharmaceutically acceptable carrier, diluent or excipient, and compound as described above with reference to structural formulae (I)-(XVI). In one embodiment according to this aspect of the invention, the compound is as described above with reference to structural formulae (I)-(XVI), but is not N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide.

In the pharmaceutical compositions according to this aspect of the invention, one or more compounds of structural formulae (I)-(XVI) may be present in association with one or more pharmaceutically acceptable carriers, diluents or excipients, and, if desired, other active ingredients. The pharmaceutical compositions containing compounds of structural formulae (I)-(XVI) may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs. In one embodiment according to this aspect of the invention, the compound is as described above with reference to structural formulae (I)-(XVI), but is not N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide.

Compositions intended for oral use can be prepared according to any suitable method for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preservative agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients that are suitable for the manufacture of tablets. These excipients can be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets can be uncoated or they can be coated by known techniques. In some cases such coatings can be prepared by suitable techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate can be employed.

Formulations for oral use can also be presented as hard gelatin capsules, wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein

the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Formulations for oral use can also be presented as lozenges.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients can be suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents such as a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions can be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents or suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, can also be present.

Pharmaceutical compositions of the invention can also be in the form of oil-in-water emulsions. The oily phase can be a vegetable oil or a mineral oil or mixtures of these. Suitable

emulsifying agents can be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions can also contain sweetening and flavoring agents.

Syrups and elixirs can be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol, glucose or sucrose. Such formulations can also contain a demulcent, a preservative, flavoring, and coloring agents. The pharmaceutical compositions can be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension can be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents that have been mentioned above. The sterile injectable preparation can also be a sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that can be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils can be employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Compounds of structural formulae (I)-(XVI) can also be administered in the form of suppositories, e.g., for rectal administration of the drug. These compositions can be prepared by mixing the compound with a suitable non-irritating excipient that is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Compounds of structural formula (I)-(XVI) can also be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Compounds suitable for use in the pharmaceutical compositions according to this aspect of the invention include compounds of Table 1, above, as well as compounds of Table 2, below. These compounds can be made according to the general scheme described above, for example using a procedure similar to that described below in Example 1.

TABLE 2

Cpd Name	Structure
1 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yl)oxybenzamide	

TABLE 2-continued


Cpd Name	Structure
2 N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-methoxybenzamide	
3 N-(1-benzylpiperidin-4-yl)-4-(1-(furan-2-ylmethyl)piperidin-4-yloxy)benzamide	
4 N-(1-benzylpiperidin-4-yl)-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide	
5 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-((1-methyl-1H-imidazol-5-yl)methyl)piperidin-4-yloxy)benzamide	
6 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(pyridin-4-ylmethyl)piperidin-4-yloxy)benzamide	

TABLE 2-continued


Cpd Name

Structure


7 N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(3-phenylpropyl)piperidin-4-yloxy)benzamide



8 N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide



9 N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(methylsulfonyl)piperidin-4-yloxy)benzamide



10 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(thiophen-2-yl)piperidin-4-yloxy)benzamide





TABLE 2-continued

Cpd Name	Structure
11 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide	
12 N-(1-benzylpiperidin-4-yl)-3-methoxy-4-(1-(furan-2-yl)piperidin-4-yloxy)benzamide	
13 N-(1-benzylpiperidin-4-yl)-3-(1-benzylpiperidin-4-yloxy)benzamide	

Another aspect of the invention relates to pharmaceutical composition comprising a compound having structural formula (XVII):



or a pharmaceutically acceptable salt, solvate, hydrate, or N-oxide thereof, and at least one pharmaceutically acceptable carrier, diluent or excipient, wherein

R<sup>1</sup> and R<sup>22</sup>, together with the nitrogen to which they are attached, form an optionally substituted monocyclic heterocycloalkyl; or

R<sup>1</sup> is H and R<sup>22</sup> is selected from —(C<sub>2</sub>-C<sub>4</sub> alkyl)-(morpholin-4-yl) and —(C<sub>2</sub>-C<sub>4</sub> alkyl)-NH—C(O)O—(C<sub>1</sub>-C<sub>6</sub> alkyl), and

all other variables are as described above with reference to structural formulae (I)-(XVI).

In one embodiment according to this aspect of the invention, R<sup>1</sup> and R<sup>22</sup>, together with the nitrogen to which they are attached, form an optionally substituted monocyclic heterocycloalkyl. The heterocycloalkyl can be, for example, piperi-

dine or piperazine. In certain embodiments according to this aspect of the invention, the heterocycloalkyl is piperazine substituted at its 4-position with —C(O)O—(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>4</sub>)-Het or —(C<sub>0</sub>-C<sub>4</sub>)-Ar. For example, the piperazine may be substituted at its 4-position with —C(O)O-tBu, -optionally-substituted pyridinylmethyl, optionally-substituted phenyl or optionally-substituted pyridinyl.

In another embodiment according to this aspect of the invention, R<sup>1</sup> is H and R<sup>22</sup> is selected from —(C<sub>2</sub>-C<sub>4</sub> alkyl)-(morpholin-4-yl) and —(C<sub>2</sub>-C<sub>4</sub> alkyl)-NH—C(O)O—(C<sub>1</sub>-C<sub>6</sub> alkyl). However, when w and x are zero, y is 1 and R<sup>5</sup> is methoxy substituted para to the benzyl methylene, R<sup>22</sup> is not —(C<sub>2</sub>-C<sub>4</sub> alkyl)-(morpholin-4-yl); and when w is 1, x and y are zero, and R<sup>3</sup> is methoxy substituted ortho to the ether oxygen, R<sup>22</sup> is not —(C<sub>2</sub>-C<sub>4</sub> alkyl)-(morpholin-4-yl). In certain embodiments of the invention, R<sup>22</sup> is —(C<sub>2</sub>-C<sub>4</sub> alkyl)-NH—C(O)O—(C<sub>1</sub>-C<sub>6</sub> alkyl). The C<sub>1</sub>-C<sub>6</sub> alkyl can be, for example, a tert-butyl group.

The pharmaceutical compositions according to this aspect of the invention can be formulated and administered as described above with respect to the pharmaceutical compositions of structural formulae (I)-(XVI), above.

Compounds suitable for use in the pharmaceutical compositions according to this aspect of the invention include compounds of Table 3, below. These compounds can be made according to the general scheme described above, for example using a procedure similar to that described below in Example 1.

TABLE 3

Cpd	Name	Structure
94	tert-butyl 3-(3-methoxy-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamido)propylcarbamate	
95	tert-butyl 3-(4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamido)propylcarbamate	
96	tert-butyl 4-(4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoyl)piperazine-1-carboxylate	
97	tert-butyl 3-(3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzamido)propylcarbamate	
98	4-(1-(4-(4-chlorobenzyl)piperidin-4-yloxy)-4-methoxy-N-(3-morpholinopropyl)benzamide	
99	tert-butyl 4-(4-(1-(4-benzyl)piperidin-4-yloxy)benzoyl)piperazine-1-carboxylate	
100	4-(1-(4-(4-chlorobenzyl)piperidin-4-yloxy)-N-(3-morpholinopropyl)benzamide	
101	3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)-N-(3-morpholinopropyl)benzamide	

TABLE 3-continued

Cpd	Name	Structure
102	tert-butyl 3-(4-(1-benzylpiperidin-4-yloxy)-3-chlorobenzamido)propylcarbamate	
103	(4-(1-benzylpiperidin-4-yloxy)phenyl)(4-phenylpiperazin-1-yl)methanone	
104	4-(1-(4-methylbenzyl)piperidin-4-yloxy)-N-(3-morpholinopropyl)benzamide	
105	(4-(1-benzylpiperidin-4-yloxy)phenyl)(4-(pyridin-2-ylmethyl)piperazin-1-yl)methanone	

While not intending to be bound by theory, the inventors surmise that compounds of structural formulae (I)-(XVII) are mimics of adiponectin which act as adiponectin receptor agonists, thereby activating the AMPK pathway. Activation of the AMPK pathway has the effect of increasing glucose uptake, decreasing glycogen synthesis and increasing fatty acid oxidation, thereby reducing glycogen, intracellular triglyceride and fatty acid concentration and causing an increase in insulin sensitivity. Because they activate the AMPK pathway, compounds of structural formulae (I)-(XVII) should also inhibit the inflammatory processes which occur during the early phases of atherosclerosis. Accordingly, compounds of structural formulae (I)-(XVII) can be useful in the treatment of type II diabetes and in the treatment and prevention of atherosclerosis, cardiovascular disease, obesity and non-alcoholic fatty liver disease.

Accordingly, another aspect of the invention is a method of activating the AMPK pathway. According to this aspect of the invention, a method for activating the AMPK pathway in a cell includes contacting the cell with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above. Data demonstrating activation of the AMPK pathway are provided below in Example 3.

Another aspect of the invention is a method of increasing fatty acid oxidation in a cell. According to this aspect of the invention, a method of increasing fatty acid oxidation in a cell includes contacting the cell with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above. Data demonstrating an increase in phosphorylated acetyl Co-A carboxylase (pACC) caused by certain compounds of the

present invention are provided below in Example 3. Acetyl Co-A carboxylase (ACC) catalyzes the formation of malonyl Co-A, a potent inhibitor of fatty acid oxidation; phosphorylation of ACC greatly reduces its catalytic activity, thereby reducing the concentration of malonyl Co-A and increasing the rate of fatty acid oxidation. Because compounds of the invention can increase the rate of phosphorylation of ACC, they can reduce the inhibition of fatty acid oxidation and therefore increase its overall rate.

Another aspect of the invention is a method of decreasing glycogen concentration in a cell. According to this aspect of the invention, a method of decreasing glycogen concentration in a cell includes contacting the cell with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above. Data demonstrating a decrease of glycogen concentration caused by certain compounds of the present invention are provided below in Example 4.

Another aspect of the invention is a method of increasing glucose uptake in a cell. According to this aspect of the invention, a method of increasing glucose uptake in a cell includes contacting the cell with an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above. Data demonstrating increase of glucose uptake caused by certain compounds of the present invention are provided below in Example 5.

Another aspect of the invention is a method of reducing triglyceride levels in a subject. According to this aspect of the invention, a method of reducing triglyceride levels in a subject includes administering to the subject an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above.

Another aspect of the invention is a method of increasing the insulin sensitivity of a subject. According to this aspect of the invention, a method of increasing insulin sensitivity of a subject includes administering to the subject an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above.

Another aspect of the invention is a method of treating type II diabetes. According to this aspect of the invention, a method of treating type II diabetes in a subject in need of such treatment includes administering to the subject an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above.

Another aspect of the invention is a method of treating or preventing atherosclerosis or cardiovascular disease. According to this aspect of the invention, a method of treating or preventing atherosclerosis or cardiovascular disease in a subject includes administering to the subject an effective amount of a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above.

In another aspect, the compounds of the invention, as activators of the AMPK pathway, the invention comprises modulating the AMPK pathway (either in vitro or in vivo) by contacting a cell with a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above, or administering a compound, pharmaceutically acceptable salt, prodrug, solvate, hydrate, N-oxide or composition described above to a mammal (e.g., a human) in an amount sufficient to modulate the AMPK activity and study the effects thereby induced. Such methods are useful for studying the AMPK pathway and its role in biological mechanisms and disease states both in vitro and in vivo.

The following examples are intended to further illustrate certain preferred embodiments of the invention and are not intended to limit the scope of the invention.

## EXAMPLES

### Example 1

#### Synthesis of N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)-piperidin-4-yloxy)benzamide

##### a. 1-(4-Methoxybenzyl)piperidin-4-ol

To a stirred solution of 4-hydroxypiperidine (0.97 g, 9.60 mmol) in anhydrous dimethylformamide (20 mL) at 0° C. was added 1-(bromomethyl)-4-methoxybenzene (1.93 g, 9.60 mmol) and triethylamine (2.16 g, 21.4 mmol). The reaction mixture was then warmed to room temperature and stirred overnight. After this time the mixture was concentrated under reduced pressure and the resulting residue was dissolved in ethyl acetate (40 mL), washed with water (20 mL), then brine (20 mL) and dried over sodium sulfate. The drying agent was filtered off and the filtrate concentrated under reduced pressure. The residue obtained was purified by flash chromatography (silica gel, 0-5% methanol/methylene chloride) to afford 1-(4-methoxybenzyl)piperidin-4-ol as a brown oil (1.70 g, 80%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.27 (d, J=7.8 Hz, 2H), 6.86 (d, J=7.8 Hz, 2H), 3.79 (s, 3H), 3.76 (m, 1H), 3.55 (s, 2H), 2.81 (m, 2H), 2.29 (m, 2H), 1.96 (m, 2H), 1.64 (m, 3H); MS (ESI): 222.1 (M+H).

##### b. Methyl 3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoate

To a stirred solution of 1-(4-methoxybenzyl)piperidin-4-ol (0.33 g, 1.49 mmol) in toluene (10 mL) at room temperature

was added triphenylphosphine (0.44 g, 1.67 mmol) and methyl 3-chloro-4-hydroxybenzoate (0.42 g, 2.25 mmol). The reaction was stirred at room temperature for 5 min. After this time, diisopropyl azodicarboxylate (0.33 g, 1.67 mmol) was added dropwise and the reaction was stirred at room temperature overnight. The mixture was concentrated under reduced pressure and the resulting residue was purified by flash chromatography (silica gel, methylene chloride/methanol/30% ammonium hydroxide=25/1/0.05) to afford methyl 3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoate as a white solid (0.05 g, 10%). MS (ESI): 390.1 (M+1).

##### c. 3-Chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoic acid

To a stirred solution of methyl 3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoate (0.04 g, 0.10 mmol) in methanol (0.5 mL) at room temperature was added 2 N sodium hydroxide (0.25 mL, 0.50 mmol) and the resulting mixture was stirred at room temperature overnight. After this time 2 N hydrochloric acid (0.3 mL, 0.60 mmol) was added and the mixture was concentrated under reduced pressure to afford 3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoic acid as a crude solid which was used without further purification. MS (ESI): 376.1 (M+1).

##### d. N-(1-Benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)-piperidin-4-yloxy)-benzamide

To a stirred mixture of 3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzoic acid in anhydrous dimethylformamide (0.5 mL) was added triethylamine (0.03 g, 0.30 mmol), 1-hydroxybenzotriazole hydrate (0.02 g, 0.15 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.03 g, 0.15 mmol) and 1-benzylpiperidin-4-amine (0.03 mg, 0.15 mmol). The mixture was stirred at room temperature overnight and then concentrated under reduced pressure. The residue obtained was purified by flash chromatography (silica gel, methylene chloride/methanol/30% ammonium hydroxide=15/1/0.05) to afford N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)-piperidin-4-yloxy)-benzamide as a white solid (0.01 g, 18%). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.76 (m, 1H), 7.60 (m, 1H), 7.30 (m, 7H), 6.94 (m, 1H), 6.87 (m, 2H), 5.83 (d, J=7.2 Hz, 1H), 4.52 (m, 1H), 3.97 (m, 1H), 3.80 (s, 3H), 3.54 (s, 4H), 2.86 (m, 4H), 2.11 (m, 8H), 1.58 (m, 4H); LCMS: >98%; MS (ESI): 548.2 (M+1), 546.2 (M-1).

### Example 2

#### <sup>1</sup>H-NMR and Mass Spectral Data

The following compounds were prepared using methods analogous to those described in Example 1 and in Scheme 1.

Compound 14: N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-chlorobenzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.74 (s, 1H), 7.59 (d, J=7.8 Hz, 1H), 7.32 (m, 10H), 6.94 (d, J=7.8 Hz, 1H), 5.83 (d, J=7.1 Hz, 1H), 4.47 (m, 1H), 3.96 (m, 1H), 3.58 (s, 4H), 2.85 (m, 2H), 2.71 (m, 2H), 2.38 (m, 2H), 2.19 (m, 2H), 1.96 (m, 6H), 1.57 (m, 2H) ppm; MS (ES) 518.2 (M+H).

Compound 15: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-((2,3-dihydrobenzo[b][1,4]dioxin-6-yl)methyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.74 (s, 1H), 7.60 (d, J=7.8 Hz, 1H), 7.28 (m, 4H), 6.93 (d, J=7.8 Hz, 1H), 6.82 (m, 4H), 5.82 (d, J=6.9 Hz, 1H), 4.47 (m, 1H), 4.24 (s, 4H), 3.96 (m, 1H), 3.52 (s, 2H), 3.44 (s, 2H), 2.84 (m, 2H),

## 65

2.70 (m, 2H), 2.37 (m, 2H), 2.17 (m, 2H), 1.95 (m, 6H), 1.61 (m, 2H) ppm; MS (ES) 576.1 (M+H).

Compound 16: N-(1-benzylpiperidin-4-yl)-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-chlorobenzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.74 (s, 1H), 7.60 (d, J=7.8 Hz, 1H), 7.30 (m, 9H), 6.93 (d, J=7.8 Hz, 1H), 5.84 (d, J=7.2 Hz, 1H), 4.48 (m, 1H), 3.96 (m, 1H), 3.52 (s, 4H), 2.86 (m, 2H), 2.72 (m, 2H), 2.40 (m, 2H), 2.18 (m, 2H), 1.94 (m, 6H), 1.57 (m, 2H), 1.32 (s, 9H) ppm; MS (ES) 574.3 (M+H).

Compound 17: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.76 (m, 1H), 7.67 (m, 1H), 7.48 (m, 2H), 7.33 (m, 5H), 6.97 (m, 3H), 5.84 (d, J=8.1 Hz, 1H), 4.68 (m, 1H), 3.97 (m, 1H), 3.56 (m, 4H), 3.35 (m, 2H), 2.87 (m, 2H), 2.19 (m, 2H), 2.03 (m, 6H), 1.59 (m, 2H) ppm; MS (ES) 572.2 (M+H).

Compound 18: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)benzoyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.77 (m, 1H), 7.65 (m, 3H), 7.53 (m, 2H), 7.30 (m, 5H), 6.95 (d, J=9.0 Hz, 1H), 5.92 (d, J=7.2 Hz, 1H), 4.76 (m, 1H), 4.04 (m, 2H), 3.70 (m, 2H), 3.51 (s, 2H), 3.42 (m, 1H), 2.85 (m, 2H), 2.16 (m, 2H), 2.07-1.78 (m, 6H), 1.54 (m, 2H) ppm; MS (ES) 600.8 (M+H).

Compound 19: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-fluorobenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.74 (m, 1H), 7.60 (m, 1H), 7.31 (m, 7H), 6.98 (m, 3H), 5.85 (d, J=8.1 Hz, 1H), 4.48 (m, 1H), 3.97 (m, 1H), 3.50 (m, 4H), 2.85 (m, 2H), 2.68 (m, 2H), 2.36 (m, 2H), 2.16 (m, 2H), 1.97 (m, 6H), 1.54 (m, 2H) ppm; MS (ES) 536.2 (M+H).

Compound 20: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.46 (m, 2H), 7.31 (m, 9H), 6.97 (m, 1H), 5.84 (d, J=8.1 Hz, 1H), 4.39 (m, 1H), 3.98 (m, 1H), 3.51 (m, 4H), 2.85 (m, 2H), 2.74 (m, 2H), 2.31 (m, 2H), 2.17 (m, 2H), 2.07-1.69 (m, 6H), 1.54 (m, 2H), 1.31 (s, 9H) ppm; MS (ES) 558.6 (M+H).

Compound 21: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(1-phenylethyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.44 (m, 2H), 7.31 (m, 10H), 6.94 (m, 3H), 5.82 (d, J=7.8 Hz, 1H), 4.32 (m, 1H), 3.97 (m, 1H), 3.51 (s, 2H), 3.46 (m, 1H), 2.85 (m, 2H), 2.72 (m, 2H), 2.21 (m, 4H), 2.04-1.63 (m, 6H), 1.54 (m, 2H), 1.38 (d, J=6.6 Hz, 3H) ppm; MS (ES) 516.5 (M+H).

Compound 22: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-fluorobenzoyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.77 (m, 1H), 7.63 (m, 1H), 7.38 (m, 2H), 7.31 (m, 5H), 7.10 (m, 2H), 6.95 (m, 1H), 5.93 (d, J=7.8 Hz, 1H), 4.74 (m, 1H), 3.97 (m, 1H), 3.72 (m, 2H), 3.52 (m, 4H), 2.86 (m, 2H), 2.17 (m, 2H), 1.97 (m, 8H), 1.55 (m, 2H) ppm; MS (ES) 550.8 (M+H).

Compound 23: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.49 (m, 4H), 7.31 (m, 5H), 7.02 (m, 1H), 6.94 (d, J=8.4 Hz, 2H), 5.84 (d, J=8.1 Hz, 1H), 4.60 (m, 1H), 3.97 (m, 1H), 3.60 (m, 2H), 3.52 (s, 2H), 3.26 (m, 2H), 2.85 (m, 2H), 2.18 (m, 2H), 2.06 (m, 6H), 1.54 (m, 2H) ppm; MS (ES) 556.6 (M+H).

Compound 24: N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(2-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.99 (m, 1H), 7.75 (m, 1H), 7.61 (m, 2H), 7.52 (m, 1H), 7.30 (m, 4H), 7.27 (m, 2H), 6.95 (d, J=9.0 Hz, 1H), 5.83 (d, J=7.5 Hz, 1H), 4.48 (m, 1H), 3.97 (m, 1H), 3.69 (s, 2H), 3.53 (s, 2H), 2.86 (m, 2H), 2.76 (m, 2H), 2.39 (m, 2H), 2.19 (m, 2H), 1.90 (m, 6H), 1.59 (m, 2H) ppm; MS (ES) 586.1 (M+H).

## 66

Compound 25: 3-fluoro-N-(1-(phenylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.52 (m, 4H), 7.25 (m, 2H), 6.97 (m, 6H), 5.88 (d, J=6.3 Hz, 1H), 4.60 (m, 1H), 4.13 (m, 1H), 3.65 (m, 4H), 3.26 (m, 2H), 2.93 (m, 2H), 2.02 (m, 6H), 1.64 (m, 2H) ppm; MS (ES) 542.1 (M+H).

Compound 26: tert-butyl 4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide)piperidine-1-carboxylate. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.54 (m, 4H), 7.03 (m, 1H), 6.93 (m, 2H), 6.11 (d, J=7.5 Hz, 1H), 4.60 (m, 1H), 4.42 (m, 2H), 4.20 (m, 1H), 3.60 (m, 2H), 3.26 (m, 2H), 2.96 (m, 2H), 2.08 (m, 2H), 1.98 (m, 4H), 1.40 (m, 2H), 1.46 (s, 9H) ppm; MS (ES) 566.1 (M+H).

Compound 27: 3-fluoro-N-(1-(4-(fluorobenzyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.49 (m, 4H), 7.28 (m, 2H), 6.99 (m, 5H), 5.86 (d, J=6.9 Hz, 1H), 4.59 (m, 1H), 3.98 (m, 1H), 3.60 (m, 2H), 3.48 (s, 2H), 3.27 (m, 2H), 2.84 (m, 2H), 2.17 (m, 2H), 2.04 (m, 6H), 1.54 (m, 2H) ppm; MS (ES) 574.6 (M+H).

Compound 28: 3-fluoro-N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.55 (m, 2H), 7.50 (m, 4H), 7.28 (m, 2H), 7.03 (m, 1H), 6.95 (d, J=8.7 Hz, 1H), 5.86 (d, J=7.5 Hz, 1H), 4.60 (m, 1H), 3.99 (m, 1H), 3.60 (m, 2H), 3.52 (s, 2H), 3.26 (m, 2H), 2.83 (m, 2H), 2.21 (m, 2H), 2.03 (m, 6H), 1.54 (m, 2H) ppm; MS (ES) 557.5 (M+H).

Compound 29: 3-fluoro-N-(1-(pyridin-3-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.54 (m, 2H), 7.65 (m, 1H), 7.49 (m, 4H), 7.24 (m, 1H), 7.04 (m, 1H), 6.94 (m, 1H), 5.85 (d, J=7.5 Hz, 1H), 4.60 (m, 1H), 3.98 (m, 1H), 3.60 (m, 2H), 3.53 (s, 2H), 3.26 (m, 2H), 2.84 (m, 2H), 2.20 (m, 2H), 2.02 (m, 6H), 1.57 (m, 2H) ppm; MS (ES) 557.5 (M+H).

Compound 30: 3-fluoro-N-(1-(pivaloylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.51 (m, 4H), 7.02 (m, 1H), 6.94 (d, J=8.7 Hz, 2H), 6.16 (d, J=7.5 Hz, 1H), 4.60 (m, 1H), 4.42 (m, 2H), 4.20 (m, 1H), 3.60 (m, 2H), 3.26 (m, 2H), 2.96 (m, 2H), 2.08 (m, 2H), 1.98 (m, 4H), 1.40 (m, 2H), 1.28 (s, 9H) ppm; MS (ES) 550.5 (M+H).

Compound 31: 3-fluoro-N-(1-(4-(fluorobenzoyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.56-7.38 (m, 6H), 7.06 (m, 3H), 6.95 (d, J=8.9 Hz, 1H), 5.98 (d, J=8.1 Hz, 1H), 4.61 (m, 1H), 4.23 (m, 1H), 3.60 (m, 2H), 3.28 (m, 2H), 3.10 (m, 2H), 2.18-1.82 (m, 6H), 1.48 (m, 4H) ppm; MS (ES) 588.5 (M+H).

Compound 32: 3-fluoro-N-(1-(pyridin-2-ylmethyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.57 (m, 1H), 7.65 (m, 1H), 7.49 (m, 4H), 7.38 (d, J=7.8 Hz, 1H), 7.17 (m, 1H), 7.05 (m, 1H), 6.95 (d, J=8.7 Hz, 1H), 5.83 (d, J=7.8 Hz, 1H), 4.60 (m, 1H), 3.99 (m, 1H), 3.67 (s, 2H), 3.59 (m, 2H), 3.26 (m, 2H), 2.88 (m, 2H), 2.28 (m, 2H), 2.04 (m, 6H), 1.62 (m, 2H) ppm; MS (ES) 557.7 (M+H).

Compound 33: methyl 4-((4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide)piperidin-1-yl)methyl)benzoate. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.99 (m, 2H), 7.44 (m, 6H), 7.03 (m, 1H), 6.94 (d, J=8.7 Hz, 1H), 5.84 (d, J=7.2 Hz, 1H), 4.60 (m, 1H), 3.98 (m, 1H), 3.90 (s, 2H), 3.61 (m, 2H), 3.56 (s, 3H), 3.26 (m, 2H), 2.84 (m, 2H), 2.20 (m, 2H), 2.03 (m, 6H), 1.58 (m, 2H) ppm; MS (ES) 614.6 (M+H).

67

Compound 34: 3-fluoro-N-(1-(isopropylsulfonyl)piperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.50 (m, 4H), 7.02 (m, 1H), 6.94 (d, J=8.1 Hz, 1H), 6.14 (m, 1H), 4.61 (m, 1H), 4.13 (m, 1H), 3.90 (m, 2H), 3.60 (m, 2H), 3.24 (m, 3H), 3.03 (m, 2H), 2.13-1.91 (m, 6H), 1.62 (m, 2H), 1.35 (m, 6H) ppm; MS (ES) 572.5 (M+H).

Compound 35: 4-((4-(3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamido)piperidin-1-yl)methyl)benzoic acid. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz) δ 8.02 (m, 2H), 7.62 (m, 2H), 7.48 (m, 4H), 7.24 (m, 1H), 7.06 (m, 2H), 4.72 (m, 1H), 4.01 (m, 3H), 3.65 (m, 2H), 3.28 (m, 2H), 2.74 (m, 2H), 2.12-1.73 (m, 10H) ppm; MS (ES) 600.6 (M+H).

Compound 36: 4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-fluoro-N-(1-phenylpiperidin-4-yl)benzamide. <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300 MHz) δ 7.63 (m, 2H), 7.45 (m, 2H), 7.35 (m, 2H), 7.22 (m, 3H), 6.99 (m, 2H), 6.83 (m, 1H), 4.66 (m, 1H), 3.98 (m, 3H), 3.70 (m, 2H), 3.10 (m, 2H), 2.82 (m, 4H), 2.03 (m, 6H), 1.79 (m, 2H), 1.31 (s, 9H) ppm; MS (ES) 544.6 (M+H).

Compound 37: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-4-ylmethyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.54 (m, 2H), 7.46 (m, 4H), 7.28 (m, 7H), 6.98 (m, 1H), 5.86 (d, J=8.1 Hz, 1H), 4.43 (m, 1H), 3.97 (m, 1H), 3.53 (s, 4H), 2.86 (m, 2H), 2.71 (m, 2H), 2.34 (m, 2H), 2.19 (m, 2H), 2.06-1.68 (m, 6H), 1.55 (m, 2H) ppm; MS (ES) 503.1 (M+H).

Compound 38: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-3-ylmethyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.52 (m, 2H), 7.67 (m, 1H), 7.46 (m, 2H), 7.31 (m, 6H), 6.97 (m, 1H), 5.84 (d, J=7.2 Hz, 1H), 4.42 (m, 1H), 3.97 (m, 1H), 3.53 (m, 4H), 2.86 (m, 2H), 2.71 (m, 2H), 2.34 (m, 2H), 2.18 (m, 2H), 2.08-1.68 (m, 6H), 1.55 (m, 2H) ppm; MS (ES) 503.5 (M+H).

Compound 39: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(pyridin-2-ylmethyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.57 (m, 2H), 7.58 (m, 2H), 7.31 (m, 6H), 7.19 (m, 1H), 6.98 (m, 1H), 5.83 (d, J=7.2 Hz, 1H), 4.41 (m, 1H), 3.96 (m, 1H), 3.53 (m, 4H), 2.86 (m, 2H), 2.70 (m, 2H), 2.33 (m, 2H), 2.18 (m, 2H), 2.07-1.69 (m, 6H), 1.55 (m, 2H) ppm; MS (ES) 503.5 (M+H).

Compound 40: 3-fluoro-N-(1-isonicotinoylpiperidin-4-yl)-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.70 (m, 2H), 7.49 (m, 4H), 7.26 (m, 2H), 7.03 (m, 1H), 6.94 (d, J=8.7 Hz, 1H), 5.99 (d, J=7.8 Hz, 1H), 4.72 (m, 1H), 4.61 (m, 1H), 4.22 (m, 1H), 3.60 (m, 3H), 3.26 (m, 3H), 2.99 (m, 1H), 2.21-1.91 (m, 6H), 1.56 (m, 2H) ppm; MS (ES) 571.4 (M+H).

Compound 41: N-(1-benzylpiperidin-4-yl)-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-3-fluorobenzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.63 (m, 3H), 7.47 (m, 5H), 7.31 (m, 3H), 6.98 (m, 1H), 5.82 (d, J=7.8 Hz, 1H), 4.43 (m, 1H), 3.99 (m, 1H), 3.54 (m, 4H), 2.85 (m, 2H), 2.70 (m, 2H), 2.34 (m, 2H), 2.18 (m, 2H), 2.08-1.72 (m, 6H), 1.56 (m, 2H) ppm; MS (ES) 527.5 (M+H).

Compound 42: N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.46 (m, 2H), 7.29 (m, 7H), 7.13 (m, 2H), 6.97 (m, 1H), 5.846 (d, J=7.5 Hz, 1H), 4.41 (m, 1H), 3.97 (m, 1H), 3.53 (s, 4H), 2.87 (m, 2H), 2.75 (m, 2H), 2.34 (m, 5H), 2.19 (m, 2H), 2.08-1.82 (m, 6H), 1.57 (m, 2H) ppm; MS (ES) 516.6 (M+H).

Compound 43: N-(1-(4-cyanobenzyl)piperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.61 (m, 2H), 7.49 (m, 6H), 6.98 (m, 3H), 5.84 (m, 1H), 4.61 (m, 1H), 3.99 (m,

68

1H), 3.570 (m, 4H), 3.26 (m, 2H), 2.82 (m, 2H), 2.22 (m, 2H), 2.02 (m, 6H), 1.58 (m, 2H) ppm; MS (ES) 581.5 (M+H).

Compound 44: 4-(1-(4-cyanophenyl)piperidin-4-yloxy)-3-fluoro-N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 8.54 (m, 2H), 7.50 (m, 3H), 7.7 (m, 3H), 7.03 (m, 1H), 6.88 (m, 2H), 5.84 (d, J=7.5 Hz, 1H), 4.63 (m, 1H), 3.98 (m, 1H), 3.63 (m, 2H), 3.52 (s, 2H), 3.357 (m, 2H), 2.83 (m, 2H), 2.21 (m, 2H), 2.01 (m, 6H), 1.58 (m, 2H) ppm; MS (ES) 514.6 (M+H).

Compound 45: N-(1-(4-cyanobenzyl)piperidin-4-yl)-4-(1-(4-cyanophenyl)piperidin-4-yloxy)-3-fluorobenzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.64 (m, 4H), 7.47 (m, 4H), 7.01 (m, 1H), 6.87 (m, 2H), 5.95 (d, J=7.8 Hz, 1H), 4.63 (m, 1H), 3.99 (m, 1H), 3.64 (m, 2H), 3.56 (s, 2H), 3.34 (m, 2H), 2.82 (m, 2H), 2.20 (m, 2H), 2.03 (m, 6H), 1.55 (m, 2H) ppm; MS (ES) 538.7 (M+H).

Compound 46: N-(1-benzylpiperidin-4-yl)-3-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.47 (d, 1H), 7.37-7.23 (m, 4H), 7.05 (d, 2H), 6.94 (d, 2H), 6.04 (d, 1H), 4.95 (m, 1H), 4.00 (m, 1H), 3.60 (m, 2H), 3.56 (s, 2H), 3.24 (m, 2H), 2.91 (m, 2H), 2.21 (m, 2H), 1.90-2.12 (m, 6H), 1.65 (m, 2H); LCMS (m/z): 538 (MH<sup>+</sup>).

Compound 47: N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.60 (d, 1H), 7.67 (d, 2H), 7.25-7.48 (m, 6H), 6.94 (d, 2H), 6.86 (d, 1H), 6.25 (d, 1H), 4.55 (m, 1H), 4.05 (m, 1H), 3.52-3.60 (m, 2H), 3.58 (s, 2H), 3.28 (m, 2H), 2.86 (m, 2H), 1.90-2.27 (m, 8H), 1.64 (m, 2H); LCMS (m/z): 573 (MH<sup>+</sup>).

Compound 48: N-(1-benzylpiperidin-4-yl)-3-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.48-7.25 (m, 10H), 7.02 (m, 1H), 6.86 (d, 2H), 6.32 (d, 1H), 4.61 (m, 1H), 4.05 (m, 1H), 3.73 (s, 2H), 3.31 (m, 2H), 3.05 (m, 2H), 2.40 (m, 2H), 2.04 (m, 4H), 1.87 (m, 4H); LCMS (m/z): 495 (MH<sup>+</sup>).

Compound 49: N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.63 (d, 2H), 7.47 (d, 2H), 6.80-7.56 (m, 7H), 6.94 (d, 1H), 6.50 (d, 1H), 4.59 (m, 1H), 4.18 (m, 1H), 4.11 (d, 2H), 3.51 (m, 2H), 3.40 (m, 2H), 3.12 (m, 2H), 2.80 (m, 2H), 2.42 (m, 2H), 2.20 (m, 4H), 2.01 (m, 2H); LCMS (m/z): 530 (MH<sup>+</sup>).

Compound 50: N-(1-benzylpiperidin-4-yl)-3-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.58 (m, 2H), 7.42 (m, 2H), 7.23-7.33 (m, 8H), 7.0 (m, 1H), 6.16 (d, 1H), 4.40 (m, 1H), 4.05 (m, 1H), 3.58 (s, 2H), 3.55 (s, 2H), 2.91 (m, 2H), 2.11-2.39 (m, 4H), 1.52-2.0 (m, 8H); LCMS (m/z): 509 (MH<sup>+</sup>).

Compound 51: N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.63 (d, 1H), 7.58 (m, 2H), 7.32-7.25 (m, 5H), 6.87 (d, 1H), 6.80 (m, 1H), 6.31 (d, 1H), 4.35 (m, 1H), 4.01 (m, 1H), 3.57 (s, 2H), 3.55 (s, 2H), 2.89 (m, 2H), 2.58 (m, 2H), 2.30 (m, 4H), 2.00 (m, 4H), 1.81 (m, 2H), 1.67 (m, 2H); LCMS (m/z): 544 (MH<sup>+</sup>).

Compound 52: N-(1-benzylpiperidin-4-yl)-3-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.76 (d, 2H), δ 7.45 (d, 2H), δ 7.38-7.26 (m, 8H), δ 7.02 (d, 1H), δ 6.21 (d, 1H), δ 4.40 (m, 1H), δ 4.05 (m, 1H), δ 3.66 (s, 2H), δ 3.58 (s, 2H), δ 3.0 (m, 2H), δ 2.70 (m, 2H), δ 2.30 (m, 4H), δ 1.85 (m, 4H), δ 1.61 (m, 2H); LCMS (m/z): 552 (MH<sup>+</sup>).

Compound 53: N-(1-benzylpiperidin-4-yl)-3-(1-(pyridin-4-yl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.21 (d, 2H), δ 7.24-7.39 (m, 8H), δ 7.02 (d, 1H), δ 6.67 (d, 2H), δ 6.28 (d, 1H), δ 4.63 (m, 1H), δ 4.05 (m, 1H),

69

δ 3.66 (m, 2H), δ 3.54 (s, 2H), δ 3.35 (m, 2H), δ 2.91 (m, 2H), δ 2.22 (m, 2H), δ 1.95 (m, 4H), δ 1.61 (m, 2H), δ 1.22 (m, 2H); LCMS (m/z): 471 (MH<sup>+</sup>).

Compound 54: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(4-trifluoromethoxybenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.16 (d, J=7.8 Hz, 1H), 7.89 (d, J=2.1 Hz, 1H), 7.74 (d, J=8.4 Hz, 1H), 7.41 (d, J=8.4 Hz, 2H), 7.28 (m, 6H), 4.62 (br s, 2H), 3.73 (br s, 2H), 3.51 (s, 2H), 3.45 (s, 2H), 2.79 (d, J=11.7 Hz, 2H), 2.61 (br s, 2H), 2.31 (t, J=7.8 Hz, 2H), 1.95 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 602 (MH<sup>+</sup>).

Compound 55: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(4-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.8 Hz, 1H), 7.65 (m, 3H), 7.52 (d, J=8.7 Hz, 2H), 7.28 (m, 5H), 4.53 (br s, 2H), 3.73 (br s, 1H), 3.58 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=10.5 Hz, 2H), 2.65 (br s, 2H), 2.28 (t, J=8.4 Hz, 2H), 1.99 (t, J=9.9 Hz, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 570 (MH<sup>+</sup>).

Compound 56: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(4-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.17 (d, J=7.2 Hz, 1H), 7.89 (d, J=1.2 Hz, 1H), 7.75 (d, J=8.4 Hz, 1H), 7.66 (d, J=7.8 Hz, 2H), 7.52 (d, J=8.1 Hz, 2H), 7.28 (m, 5H), 4.63 (br s, 2H), 3.71 (br s, 2H), 3.58 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=11.4 Hz, 2H), 2.62 (br s, 2H), 2.32 (t, J=8.1 Hz, 2H), 1.98 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 586 (MH<sup>+</sup>).

Compound 57: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(4-trifluoromethoxybenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.11 (d, J=7.5 Hz, 1H), 7.64 (m, 2H), 7.42 (d, J=8.4 Hz, 1H), 7.28 (m, 7H), 4.52 (br s, 1H), 3.71 (br s, 1H), 3.51 (s, 2H), 3.45 (s, 2H), 2.79 (d, J=10.5 Hz, 2H), 2.64 (br s, 2H), 2.26 (t, J=9.3 Hz, 2H), 1.97 (m, 4H), 1.72 (m, 4H), 1.54 (m, 2H); LCMS (m/z): 586 (MH<sup>+</sup>).

Compound 58: N-(1-Methylpiperidin-4-yl)-3-chloro-4-[1-(4-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.16 (d, J=7.8 Hz, 1H), 7.91 (s, 1H), 7.75 (d, J=9.0 Hz, 1H), 7.66 (d, J=7.8 Hz, 2H), 7.53 (d, J=7.8 Hz, 2H), 7.24 (d, J=8.4 Hz, 1H), 4.63 (br s, 1H), 3.68 (br s, 1H), 3.58 (s, 2H), 2.76 (d, J=11.7 Hz, 2H), 2.62 (br s, 2H), 2.33 (t, J=8.7 Hz, 2H), 1.92 (m, 4H), 1.73 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 510 (MH<sup>+</sup>).

Compound 59: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(3-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.5 Hz, 1H), 7.59 (m, 7H), 7.28 (m, 4H), 4.53 (d, J=4.2 Hz, 2H), 3.71 (br s, 1H), 3.58 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=10.5 Hz, 2H), 2.63 (br s, 2H), 2.28 (t, J=9.3 Hz, 2H), 2.01 (m, 4H), 1.72 (m, 4H), 1.54 (m, 2H); LCMS (m/z): 570 (MH<sup>+</sup>).

Compound 60: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(3-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.17 (d, J=7.8 Hz, 1H), 7.90 (d, J=1.8 Hz, 1H), 7.75 (dd, J=1.8 and 8.5 Hz, 1H), 7.58 (m, 5H), 7.28 (m, 5H), 4.62 (br s, 1H), 3.71 (br s, 1H), 3.58 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=11.4 Hz, 2H), 2.62 (br s, 2H), 2.33 (t, J=8.7 Hz, 2H), 1.98 (m, 4H), 1.71 (m, 4H), 1.56 (m, 2H); LCMS (m/z): 586 (MH<sup>+</sup>).

Compound 61: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(4-fluorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.11 (d, J=7.8 Hz, 1H), 7.64 (m, 3H), 7.21 (m, 7H), 7.11 (t, J=8.7 Hz, 2H), 4.51 (br s, 1H), 3.72 (br s, 1H), 3.46 (s, 2H), 3.45 (s, 2H), 2.80 (d, J=11.4 Hz, 2H), 2.62 (br s, 2H), 2.24 (t, J=9.3 Hz, 2H), 1.97 (m, 4H), 1.71 (m, 4H), 1.56 (m, 2H); LCMS (m/z): 520 (MH<sup>+</sup>).

Compound 62: N-(1-Benzylpiperidin-4-yl)-3,5-dichloro-4-[1-(3-trifluoromethylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.36 (d, J=7.5 Hz,

70

1H), 7.91 (s, 2H), 7.58 (m, 4H), 7.28 (m, 4H), 4.31 (br s, 1H), 3.70 (br s, 1H), 3.57 (s, 2H), 3.45 (s, 2H), 2.77 (t, J=12.6 Hz, 4H), 2.15 (t, J=9.9 Hz, 2H), 1.89 (m, 8H), 1.55 (m, 3H); LCMS (m/z): 620 (MH<sup>+</sup>).

Compound 63: N-(1-Benzylpiperidin-4-yl)-4-[1-(4-trifluoromethoxybenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.02 (d, J=7.5 Hz, 1H), 7.75 (d, J=8.1 Hz, 2H), 7.41 (d, J=8.1 Hz, 2H), 7.28 (m, 6H), 6.95 (d, J=8.7 Hz, 2H), 4.46 (br s, 1H), 3.71 (br s, 1H), 3.50 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=11.4 Hz, 2H), 2.64 (m, 2H), 2.25 (t, J=9.3 Hz, 2H), 1.61 (m, 6H); LCMS (m/z): 568 (MH<sup>+</sup>).

Compound 64: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(4-chlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.5 Hz, 1H), 7.64 (m, 2H), 7.37-7.21 (m, 9H), 4.51 (br s, 1H), 3.71 (br s, 1H), 3.47 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=10.8 Hz, 2H), 2.62 (m, 2H), 2.25 (t, J=10.5 Hz, 2H), 1.95 (m, 4H), 1.71 (m, 4H), 1.55 (t, J=11.5 Hz, 2H); LCMS (m/z): 536 (MH<sup>+</sup>).

Compound 65: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(4-chlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.17 (d, J=7.5 Hz, 1H), 7.90 (d, J=2.4 Hz, 1H), 7.74 (dd, J=2.1 and 8.5 Hz, 1H), 7.28 (m, 9H), 4.61 (br s, 1H), 3.71 (br s, 2H), 3.47 (s, 2H), 3.45 (s, 2H), 2.80 (d, J=11.1 Hz, 2H), 2.60 (m, 2H), 2.29 (t, J=9.0 Hz, 2H), 1.95 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 552 (MH<sup>+</sup>).

Compound 66: N-[1-(4-Fluorobenzyl)piperidin-4-yl]-3-chloro-4-[1-(4-chlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.17 (d, J=7.5 Hz, 1H), 8.11 (s, 1H), 7.89 (d, J=2.1 Hz, 1H), 7.74 (dd, J=2.1 and 8.5 Hz, 1H), 7.32 (m, 5H), 7.23 (d, J=8.7 Hz, 1H), 7.12 (t, J=8.7 Hz, 2H), 4.61 (br s, 1H), 3.70 (br s, 1H), 3.47 (s, 2H), 3.44 (s, 2H), 2.79 (d, J=11.7 Hz, 2H), 2.60 (m, 2H), 2.29 (t, J=8.1 Hz, 2H), 1.97 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 570 (MH<sup>+</sup>).

Compound 67: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.18 (d, J=7.8 Hz, 1H), 7.90 (d, J=2.4 Hz, 1H), 7.76 (m, 3H), 7.51 (d, J=8.1 Hz, 2H), 7.28 (m, 5H), 4.63 (br s, 1H), 3.70 (br s, 1H), 3.58 (s, 2H), 3.45 (s, 2H), 2.80 (d, J=11.4 Hz, 2H), 2.61 (m, 2H), 2.32 (t, J=8.1 Hz, 2H), 1.95 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 543 (MH<sup>+</sup>).

Compound 68: N-[1-(4-Fluorobenzyl)piperidin-4-yl]-3-chloro-4-[1-(4-methylbenzyl)piperidin-4-yloxy]benzamide (as the formate salt). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.41 (d, J=7.5 Hz, 1H), 7.92 (d, J=1.8 Hz, 1H), 7.77 (d, J=9.0 Hz, 1H), 7.53 (m, 2H), 7.41-7.28 (m, 7H), 4.29 (m, 4H), 3.34 (m, 5H), 3.06 (m, 4H), 2.33 (s, 3H), 2.06 (s, 4H), 1.99 (m, 1H), 1.74 (m, 3H); LCMS (m/z): 550 (MH<sup>+</sup>).

Compound 69: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(3,4-difluorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.5 Hz, 1H), 7.64 (m, 2H), 7.28 (m, 7H), 7.13 (m, 1H), 4.52 (br s, 1H), 3.71 (br s, 1H), 3.47 (s, 2H), 3.44 (s, 2H), 2.80 (d, J=11.4 Hz, 2H), 2.62 (m, 2H), 2.26 (t, J=8.7 Hz, 2H), 1.95 (m, 4H), 1.68 (m, 6H); LCMS (m/z): 538 (MH<sup>+</sup>).

Compound 70: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.11 (d, J=7.8 Hz, 1H), 7.77 (d, J=8.4 Hz, 2H), 7.64 (m, 2H), 7.50 (d, J=8.1 Hz, 2H), 7.31 (m, 5H), 4.53 (br s, 1H), 3.70 (br s, 1H), 3.57 (s, 2H), 3.44 (s, 2H), 2.78 (d, J=11.7 Hz, 2H), 2.63 (m, 2H), 2.28 (t, J=8.7 Hz, 2H), 1.96 (m, 4H), 1.72 (m, 4H), 1.56 (m, 2H); LCMS (m/z): 561 (MH<sup>+</sup>).

Compound 71: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(3,4-difluorobenzyl)piperidin-4-yloxy]benza-

71

amide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.11 (d, J=7.8 Hz, 1H), 7.64 (m, 2H), 7.34 (m, 7H), 7.14 (m, 1H), 4.52 (br s, 1H), 3.70 (br s, 1H), 3.47 (s, 2H), 3.44 (s, 2H), 2.78 (d, J=12.0 Hz, 2H), 2.63 (m, 2H), 2.26 (t, J=10.2 Hz, 2H), 1.97 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 572 (MH<sup>+</sup>).

Compound 72: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(4-chlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.13 (d, J=7.5 Hz, 1H), 7.64 (m, 2H), 7.34 (m, 9H), 4.52 (br s, 1H), 3.71 (br s, 1H), 3.47 (s, 2H), 3.44 (s, 2H), 2.77 (d, J=11.4 Hz, 2H), 2.63 (m, 2H), 2.24 (t, J=9.0 Hz, 2H), 1.96 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 570 (MH<sup>+</sup>).

Compound 73: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(4-methylbenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.5 Hz, 1H), 7.63 (t, J=11.9 Hz, 2H), 7.32 (m, 5H), 7.12 (m, 4H), 4.50 (br s, 1H), 3.73 (br s, 1H), 3.44 (s, 2H), 3.42 (s, 2H), 2.77 (d, J=11.4 Hz, 2H), 2.61 (m, 2H), 2.26 (s, 3H), 2.22 (t, J=12.6 Hz, 2H), 1.96 (m, 4H), 1.65 (m, 6H); LCMS (m/z): 550 (MH<sup>+</sup>).

Compound 74: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(3,4-difluorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.17 (d, J=7.5 Hz, 1H), 7.90 (d, J=2.1 Hz, 1H), 7.75 (d, J=8.4 Hz, 1H), 7.28 (m, 8H), 7.14 (m, 1H), 4.62 (br s, 1H), 3.70 (br s, 1H), 3.48 (s, 2H), 3.45 (s, 2H), 2.80 (d, J=11.4 Hz, 2H), 2.60 (m, 2H), 2.30 (t, J=8.7 Hz, 2H), 1.95 (m, 4H), 1.71 (m, 4H), 1.56 (m, 2H); LCMS (m/z): 554 (MH<sup>+</sup>).

Compound 75: N-(1-Benzylpiperidin-4-yl)-3,5-dichloro-4-[1-(4-chlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.36 (d, J=7.8 Hz, 1H), 7.91 (s, 2H), 7.28 (m, 8H), 7.14 (m, 1H), 4.30 (br s, 1H), 3.70 (br s, 1H), 3.45 (s, 4H), 2.77 (m, 4H), 2.10 (t, J=10.5 Hz, 2H), 2.00 (t, J=10.5 Hz, 2H), 1.88 (m, 2H), 1.77 (m, 4H); LCMS (m/z): 586 (MH<sup>+</sup>).

Compound 76: N-(1-Benzylpiperidin-4-yl)-3,5-dichloro-4-[1-(3,4-difluorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.36 (d, J=7.8 Hz, 1H), 7.91 (s, 2H), 7.28 (m, 6H), 7.13 (m, 1H), 4.30 (m, 1H), 3.70 (m, 1H), 3.45 (s, 4H), 2.77 (m, 4H), 2.12 (t, J=9.6 Hz, 2H), 2.00 (t, J=11.1 Hz, 2H), 1.81 (m, 8H), 1.55 (m, 3H); LCMS (m/z): 588 (MH<sup>+</sup>).

Compound 77: N-(1-Benzylpiperidin-4-yl)-3,5-dichloro-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide (as the formate salt). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.36 (d, J=7.2 Hz, 1H), 7.76 (d, J=8.1 Hz, 2H), 7.49 (d, J=8.1 Hz, 2H), 7.28 (m, 5H), 4.31 (m, 1H), 3.70 (m, 1H), 3.56 (s, 2H), 3.45 (s, 2H), 2.77 (t, J=11.1 Hz, 4H), 2.15 (t, J=11.4 Hz, 2H), 1.99 (t, J=9.9 Hz, 2H), 1.81 (m, 5H), 1.55 (m, 3H); LCMS (m/z): 577 (MH<sup>+</sup>).

Compound 78: N-(1-t-Butoxycarbonylpiperidin-4-yl)-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.67 (d, J=8.7 Hz, 2H), 7.59 (d, J=8.4 Hz, 2H), 7.45 (d, J=7.8 Hz, 2H), 7.25 (m, 1H), 6.89 (d, J=8.7 Hz, 2H), 5.53 (d, J=7.8 Hz, 1H), 4.40 (br s, 1H), 4.09 (m, 3H), 3.56 (s, 2H), 2.90 (t, J=10.5 Hz, 2H), 2.69 (m, 2H), 2.34 (m, 2H), 2.01 (m, 4H), 1.85 (m, 3H), 1.46 (s, 9H); LCMS (m/z): 519 (MH<sup>+</sup>).

Compound 79: N-(Piperidin-4-yl)-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide hydrochloride. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.83 (s, 3H), 8.33 (d, J=6.9 Hz, 1H), 7.88 (m, 5H), 7.03 (m, 2H), 4.84 (s, 1H), 4.43 (m, 2H), 3.38 (m, 1H), 3.28 (d, J=11.7 Hz, 3H), 3.18 (s, 2H), 2.99 (m, 2H), 2.21 (d, J=11.1 Hz, 2H), 2.00 (m, 4H), 1.77 (m, 2H); LCMS (m/z): 419 (MH<sup>+</sup>).

Compound 80: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(3,4-dichlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.11 (d, J=7.5 Hz,

72

1H), 7.64 (m, 2H), 7.54 (s, 1H), 7.31 (m, 5H), 4.51 (m, 1H), 3.72 (br s, 2H), 3.49 (s, 2H), 3.44 (s, 2H), 2.77 (d, J=12 Hz, 4H), 2.63 (m, 3H), 2.27 (t, J=8.7 Hz, 2H), 2.00 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 604 (MH<sup>+</sup>).

Compound 81: N-[1-(4-Chlorobenzyl)piperidin-4-yl]-3-fluoro-4-[1-(4-cyanophenyl)piperidin-4-yloxy]benzamide (as the formate salt). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.15 (d, J=7.5 Hz, 1H), 7.66 (t, J=9.3 Hz, 2H), 7.55 (d, J=8.7 Hz, 2H), 7.34 (m, 5H), 7.03 (d, J=9.0 Hz, 2H), 4.77 (m, 2H), 3.69 (m, 4H), 3.44 (s, 2H), 2.78 (d, J=11.1 Hz, 2H), 2.00 (m, 4H), 2.00 (m, 4H), 1.72 (m, 4H), 1.55 (m, 2H); LCMS (m/z): 547 (MH<sup>+</sup>).

Compound 82: N-(1-Benzylpiperidin-4-yl)-3-fluoro-4-[1-(3,4-dichlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.12 (d, J=7.5 Hz, 1H), 7.64 (m, 2H), 7.53 (m, 5H), 7.27 (m, 6H), 4.53 (s, 1H), 3.72 (s, 1H), 3.44 (s, 2H), 2.79 (d, J=11.4 Hz, 2H), 2.61 (m, 2H), 2.16 (t, J=8.7 Hz, 1H), 2.02 (m, 4H), 1.69 (m, 3H), 1.38 (m, 3H); LCMS (m/z): 570 (MH<sup>+</sup>).

Compound 83: N-[1-(4-Pyridinylmethyl)piperidin-4-yl]-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide (as the formate salt). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.50 (d, J=4.8 Hz, 2H), 8.11 (s, 1H), 8.06 (d, J=7.2 Hz, 1H), 7.77 (d, J=8.4 Hz, 3H), 7.51 (d, J=7.8 Hz, 2H), 6.96 (d, J=8.7 Hz, 2H), 6.50 (s, 1H), 4.49 (s, 1H), 3.75 (s, 1H), 3.64 (s, 2H), 3.57 (s, 2H), 2.84 (d, J=11.1 Hz, 2H), 2.67 (m, 2H), 2.34 (t, J=8.7 Hz, 2H), 2.15 (m, 2H), 1.93 (br s, 2H), 1.62 (m, 4H); LCMS (m/z): 510 (MH<sup>+</sup>).

Compound 84: N-(1-Benzylpiperidin-4-yl)-3-chloro-4-[1-(3,4-dichlorobenzyl)piperidin-4-yloxy]benzamide. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.16 (d, J=7.8 Hz, 1H), 7.90 (s, 1H), 7.75 (d, J=8.4 Hz, 1H), 7.54 (m, 3H), 7.28 (m, 5H), 4.62 (s, 1H), 3.71 (s, 1H), 3.49 (s, 2H), 3.44 (s, 2H), 2.81 (d, J=10.8 Hz, 2H), 2.61 (m, 2H), 2.31 (t, J=8.7 Hz, 2H), 1.99 (m, 4H), 1.71 (m, 4H), 1.38 (m, 3H); LCMS (m/z): 586 (MH<sup>+</sup>).

Compound 85: N-[1-(2-Pyridinylmethyl)piperidin-4-yl]-4-[1-(4-cyanobenzyl)piperidin-4-yloxy]benzamide (as the trifluoroacetate salt). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz): δ 8.67 (d, J=5.7 Hz, 1H), 8.29 (d, J=6.0 Hz, 1H), 7.95 (m, 2H), 7.82 (s, 2H), 7.71 (d, J=6.3 Hz, 2H), 7.51 (m, 2H), 7.02 (s, 2H), 4.49 (s, 1H), 4.02 (m, 5H), 3.45 (d, J=12.3 Hz, 2H), 3.20 (m, 4H), 2.25 (s, 2H), 2.00 (m, 4H), 1.88 (m, 2H); LCMS (m/z): 510 (MH<sup>+</sup>).

Compound 86: N-(1-benzylpiperidin-4-yl)-3-(1-(pyridin-2-yl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.74 (s, 1H), 8.18 (d, 1H), 8.04 (s, 1H), 7.81 (d, 1H), 7.24-7.58 (m, 6H), 7.00 (m, 2H), 6.81 (m, 2H), 4.80 (m, 1H), 4.22 (m, 1H), 4.18 (s, 2H), 3.94 (m, 4H), 3.52 (m, 2H), 2.90 (m, 2H), 2.44 (m, 2H), 2.01-2.21 (m, 6H); LCMS (m/z): 471 (MH<sup>+</sup>).

Compound 87: N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-3-(1-(4-(trifluoromethyl)phenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.70 (d, 2H), δ 7.65 (d, 2H), δ 7.45 (d, 2H), δ 7.25-7.39 (m, 2H), δ 7.01 (d, 1H), δ 6.92 (d, 2H), δ 6.80 (d, 1H), δ 4.60 (m, 1H), δ 4.25 (m, 1H), δ 4.09 (s, 2H), δ 3.48 (m, 2H), δ 3.44 (m, 2H), δ 3.22 (m, 2H), δ 2.86 (m, 2H), δ 2.40 (m, 2H), δ 2.20 (m, 2H), δ 2.10 (m, 2H), δ 1.95 (m, 2H); LCMS (m/z): 539 (MH<sup>+</sup>).

Compound 88: N-(1-(pyridin-4-ylmethyl)piperidin-4-yl)-3-(1-(4-cyanophenyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 8.64 (d, 2H), 7.25-7.52 (m, 7H), 7.03 (d, 1H), 6.85 (d, 2H), 6.87 (d, 1H), 4.60 (m, 1H), 4.15 (m, 1H), 3.89 (s, 2H), 3.61 (m, 2H), 3.30 (m, 4H), 2.62 (m, 2H), 2.02-2.13 (m, 8H); LCMS (m/z): 496 (MH<sup>+</sup>).

Compound 89: N-(1-benzylpiperidin-4-yl)-3-(1-(3-cyanobenzyl)piperidin-4-yloxy)benzamide. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz): δ 7.65 (s, 1H), 7.24-7.55 (m, 11H), 7.02 (d, 1H),

73

7.02 (d, 1H), 6.18 (d, 1H), 4.40 (m, 1H), 4.05 (m, 1H), 3.64 (s, 2H), 3.54 (s, 2H), 2.97 (m, 2H), 2.70 (m, 2H), 2.31 (m, 4H), 1.77-2.05 (m, 8H); LCMS (m/z): 510 (MH<sup>+</sup>).

## Example 3

5

## Screening of Compounds Using a Competitive Binding Assay

Candidate compounds are assayed for binding to a membrane-bound adiponectin receptor by performing a competitive binding assay with adiponectin. HEK 293 cellular membrane is coated onto a COSTAR 384 plate, which is then blocked with 1% casein. Polyhistidine-tagged globular adiponectin and a candidate compound is incubated with the membrane in HEPES buffer. Unbound ligands were washed away and the degree of binding of the adiponectin was determined using horseradish peroxidase-conjugated anti-polyhistidine. Compounds that compete with adiponectin binding to the membrane (i.e., give a reduced signal compared to a control performed without a candidate compound) are chosen as hits and are further screened using the below-described functional assays to identify adiponectin receptor agonists.

## Example 4

25

## Increase in AMPK Activity and Rate of Fatty Acid Oxidation

An in-cell western assay was performed to demonstrate the activation of AMPK in human liver cells by globular adiponectin using glutathione S-transferase (GST). AMPK activity was measured by the relative concentration of phosphorylated acetyl Co-A carboxylase, which is one of the products of AMPK. As described above, an increase in pACC correlates with an increase in the rate of fatty acid oxidation. FIG. 1 is a plot of AMPK activity vs. concentration for glutathione S-transferase (GST) and its fusion protein with globular adiponectin (gAd). The presence of gAd clearly increases AMPK. FIG. 2 is a plot of AMPK activity vs. concentration for both gAd and its polyhistidine-tagged analog. The presence of the polyhistidine tag caused little difference in AMPK activity.

Compounds of Tables 1, 2 and 3 were assayed for their ability to activate AMPK using an enzyme-linked immunosorbent assay. The EC<sub>50</sub> values for AMPK activation for compounds 1-16 are presented in Table 4 below, in which "A" is less than 1 μM; "B" is 1-10 μM; "C" is 10-20 μM; "D" is 20-50 μM; "E" is 50-100 μM, and "F" is >100 μM:

TABLE 4

Cpd No.	AMPK EC <sub>50</sub> (μM)
1	A
2	A
3	B
4	F
5	E
6	A
7	A
8	B
9	D
10	B
11	B
12	B
13	D
14	A
15	A
16	A

74

TABLE 4-continued

Cpd No.	AMPK EC <sub>50</sub> (μM)
17	A
18	A
19	A
20	A
21	B
22	B
23	A
24	C
25	F
26	A
27	A
28	A
29	A
30	A
31	A
32	A
33	A
34	A
35	B
36	A
37	A
38	B
39	B
40	A
41	A
42	A
43	A
44	A
45	A
46	A
47	A
48	A
49	A
50	B
51	A
52	B
53	C
54	A
55	A
56	A
57	A
58	B
59	A
60	A
61	A
62	A
63	A
64	A
65	A
66	A
67	A
68	A
69	A
70	A
71	A
72	A
73	A
74	A
75	A
76	A
77	A
78	A
79	F
80	A
81	A
82	A
83	A
84	A
85	A
86	A
87	A
88	B
89	B
90	B
91	A
92	B
93	F
94	A

75

TABLE 4-continued

Cpd No.	AMPK EC <sub>50</sub> (μM)
95	A
96	B
97	B
98	B
99	B
100	B
101	C
102	C
103	D
104	C
105	C

Western blot assays for AMPK activity were performed on compounds 1, 2, 94 and 95. FIG. 3 presents gel electrophoresis data for compounds 1, 2, 94 and 95 relative to GST-gAd. Each of these compounds demonstrated strong activity in the western blot assay.

## Example 5

## Decrease in Glycogen Concentration

The ability of compounds of the present invention to reduce the glycogen content in liver cells was determined using a functional assay. Human liver cells (HepG2) were seeded in a 96 well plate and on the next day were treated with compounds 1, 2, 94 and 95 for 16 hours. The glycogen content of the cells was determined using Amplex Red. Data for compounds 1, 2, 94 and 95 are presented in FIG. 4. Each of these compounds demonstrated a sub-micromolar IC<sub>50</sub> value.


## Example 6

## Increase in Glucose Uptake

The ability of compounds of the present invention to increase glucose uptake in skeletal muscle cells was determined by a functional assay. Rat differentiated L6 myotube cells were incubated for two hours with GST-gAd and compounds 1, 2, 94 and 95, then with 2-deoxy-D-[6-<sup>3</sup>H]glucose for 10 minutes. Uptake of 2-deoxy-D-[6-<sup>3</sup>H]glucose was determined by liquid scintillation counting. FIG. 5 presents data for compounds 1, 2, 94 and 95, shown relative to adiponectin.

What is claimed is:

1. A compound having the structural formula



or a pharmaceutically acceptable salt or N-oxide thereof, wherein

each R<sup>3</sup> is independently selected from halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), acyl, carboxylate, carboxamide and nitro; w is 0, 1, 2 or 3;

76

each R<sup>4</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), -halogen, —NO<sub>2</sub> and —CN, and two R<sup>4</sup> on the same carbon optionally combine to form oxo;

x is 0, 1, 2, 3 or 4;

each R<sup>5</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-R<sup>7</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>8</sup>R<sup>9</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-OR<sup>10</sup>, —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>R<sup>10</sup>, -halogen, —NO<sub>2</sub> and —CN;

y is 0, 1, 2, 3 or 4;

G is a single bond, —CH<sub>2</sub>— or —C(O)—;

v is 0, 1, 2, 3 or 4;

each R<sup>15</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), -halogen, —NO<sub>2</sub> and —CN, and two R<sup>15</sup> on the same carbon optionally combine to form oxo; and

R<sup>17</sup> is optionally-substituted aryl;

in which

each L is independently selected

from —NR<sup>9</sup>C(O)O—, —NR<sup>9</sup>C(O)—NR<sup>9</sup>—, —NR<sup>9</sup>C(O)S—, —NR<sup>9</sup>C(O)—, —NR<sup>9</sup>C(S)O—, —NR<sup>9</sup>C(S)—NR<sup>9</sup>—, —NR<sup>9</sup>C(S)S—, —NR<sup>9</sup>C(S)O—, —OC(O)NR<sup>9</sup>—, —SC(O)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —OC(S)NR<sup>9</sup>—, —SC(S)NR<sup>9</sup>—, —C(S)NR<sup>9</sup>—, —S(O)<sub>0-2</sub>—, —C(O)O—, —OC(O)—, —C(S)O—, —OC(S)—, —C(O)S—, —SC(O)—, —C(S)S—, —SC(S)—, —OC(O)O—, —SC(O)O—, —OC(O)S—, —SC(S)O—, —OC(S)S—, —NR<sup>9</sup>C(NR<sup>9</sup>)NR<sup>9</sup>—, —NR<sup>9</sup>SO<sub>2</sub>—, —SO<sub>2</sub>NR<sup>9</sup>— and —NR<sup>9</sup>SO<sub>2</sub>NR<sup>9</sup>—,

each R<sup>7</sup>, R<sup>8</sup> and R<sup>10</sup> is independently selected from H, —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-L-(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-NR<sup>9</sup>(C<sub>0</sub>-C<sub>6</sub> alkyl), —(C<sub>0</sub>-C<sub>6</sub> alkyl)-O—(C<sub>0</sub>-C<sub>6</sub> alkyl) and —(C<sub>0</sub>-C<sub>6</sub> alkyl)-S(O)<sub>0-2</sub>—(C<sub>0</sub>-C<sub>6</sub> alkyl),

each R<sup>9</sup> is independently selected from —H, —(C<sub>1</sub>-C<sub>4</sub> alkyl) and —C(O)—(C<sub>1</sub>-C<sub>4</sub> alkyl), and

each R<sup>16</sup> is independently selected from —(C<sub>1</sub>-C<sub>6</sub> alkyl), —(C<sub>1</sub>-C<sub>6</sub> fluoroalkyl), and two R<sup>16</sup> on the same carbon optionally combine to form oxo;

provided that the compound is not

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-methoxybenzamide; or

N-(1-benzylpiperidin-4-yl)-3-(1-benzylpiperidin-4-yloxy)benzamide.

2. A compound according to claim 1, wherein y is 0.


3. A compound according to claim 1, wherein y is 1.

4. A compound according to claim 1, wherein y is 1, 2, 3 or 4 and at least one R<sup>5</sup> is halo, cyano, trifluoromethyl or trifluoromethoxy.


5. A compound according to claim 1, wherein y is 1 and R<sup>5</sup> is selected from the group consisting of halo, cyano, —(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), —O—(C<sub>1</sub>-C<sub>4</sub> fluoroalkyl), carboxylate, carboxamide and nitro.

6. A compound according to claim 1, wherein the compound has the structural formula


77




78



7. A compound according to claim 1, wherein the compound has the structural formula




10



15

17. A compound according to claim 1, having the structural formula



8. A compound according to claim 1, wherein G is  $-\text{CH}_2-$ .

9. A compound according to claim 1, wherein G is  $-\text{C}(\text{O})-$ .

10. A compound according to claim 1, wherein  $\text{R}^{17}$  is phenyl substituted with an electron withdrawing group.


in which

$\text{R}^3$  is H or halo;

$\text{R}^{11}$  is H, halo, cyano, or a carboxylate; and

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

18. A compound according to claim 1, the having structural formula



11. A compound according to claim 1, wherein  $\text{R}^{17}$  is unsubstituted phenyl.

12. A compound according to claim 1, wherein w is 0.

13. A compound according to claim 1, wherein w is 1.

14. A compound according to claim 1, wherein x is 0 and v is 0.

15. A compound according to claim 1, wherein

w is 0 or 1;

x is 0;

v is 0;

y is 0, 1, 2 or 3; and

G is  $-\text{CH}_2-$  or  $-\text{CO}-$ .

16. A compound according to claim 15, having the structural formula

in which

$\text{R}^{11}$  is H, halo, cyano, or a carboxylate; and

$\text{R}^{12}$  and  $\text{R}^{13}$  are independently H, trifluoromethyl, trifluoromethoxy, halo or cyano.

19. A compound according to claim 1, wherein the compound is selected from the group consisting of:

N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-chlorobenzamide;

N-(1-benzylpiperidin-4-yl)-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-chlorobenzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-fluorobenzyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)benzamide;

N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(2-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;

79

4-(1-(4-tert-butylbenzyl)piperidin-4-yloxy)-3-fluoro-N-(1-phenylpiperidin-4-yl)benzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-2-chloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-fluoro-4-(1-(4-fluorobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(3-(trifluoromethyl)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-(4-(trifluoromethoxy)benzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)benzamide;  
 3-chloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-N-(1-(4-fluorobenzyl)piperidin-4-yl)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide;  
 3-chloro-N-(1-(4-fluorobenzyl)piperidin-4-yl)-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;

80

N-(1-(4-chlorobenzyl)piperidin-4-yl)-3-fluoro-4-(1-(4-methylbenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(4-chlorobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(3,4-difluorobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3,5-dichloro-4-(1-(4-cyanobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-(4-chlorobenzyl)piperidin-4-yl)-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)-3-fluorobenzamide;  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(3,4-dichlorobenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-3-(1-(3-cyanobenzyl)piperidin-4-yloxy)benzamide;  
 or a pharmaceutically acceptable salt or N-oxide thereof.  
**20.** A pharmaceutical composition comprising a compound according to claim **1** or a compound selected from the group consisting of  
 N-(1-benzylpiperidin-4-yl)-3-chloro-4-(1-(4-methoxybenzyl)piperidin-4-yloxy)benzamide;  
 N-(1-benzylpiperidin-4-yl)-4-(1-benzylpiperidin-4-yloxy)-3-methoxybenzamide; and  
 N-(1-benzylpiperidin-4-yl)-3-(1-benzylpiperidin-4-yloxy)benzamide,  
 or a pharmaceutically acceptable salt or N-oxide thereof.  
**21.** A method for activating the AMPK pathway in a cell, the method comprising contacting the cell with an effective amount of a compound according to claim **1** or a pharmaceutically acceptable salt or N-oxide thereof.  
**22.** A method for reducing triglyceride levels in a subject, the method comprising administering to the subject an effective amount of a pharmaceutical composition described in claim **1** or a pharmaceutically acceptable salt or N-oxide thereof.  
**23.** A method for treating type II diabetes in a subject, the method comprising administering to the subject an effective amount of a pharmaceutical composition described in claim **1** or a pharmaceutically acceptable salt or N-oxide thereof.  
**24.** A method for treating atherosclerosis in a subject, the method comprising administering to the subject an effective amount of a pharmaceutical composition described in claim **1** or a pharmaceutically acceptable salt or N-oxide thereof.

\* \* \* \* \*