



US009255299B2

(12) **United States Patent**
Ji et al.

(10) **Patent No.:** **US 9,255,299 B2**
(45) **Date of Patent:** **Feb. 9, 2016**

(54) **COMPOSITIONS AND METHODS FOR DETECTION OF MYCOBACTERIUM AVIUM PARATUBERCULOSIS**

(71) Applicant: **LIFE TECHNOLOGIES CORPORATION**, Carlsbad, CA (US)

(72) Inventors: **Yongmei Ji**, Chicago, IL (US); **Pius Brzoska**, Woodside, CA (US); **Angela Burrell**, Austin, TX (US); **Craig Cummings**, Pacifica, CA (US); **Catherine O'Connell**, Austin, TX (US)

(73) Assignee: **LIFE TECHNOLOGIES CORPORATION**, Carlsbad, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 172 days.

(21) Appl. No.: **13/785,409**

(22) Filed: **Mar. 5, 2013**

(65) **Prior Publication Data**
US 2013/0260374 A1 Oct. 3, 2013

Related U.S. Application Data
(60) Provisional application No. 61/619,196, filed on Apr. 2, 2012.

(51) **Int. Cl.**
C12P 19/34 (2006.01)
C12Q 1/68 (2006.01)

(52) **U.S. Cl.**
CPC **C12Q 1/689** (2013.01); **C12Q 2600/112** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS

7,108,968 B2	9/2006	Gingeras et al.	
2007/0042383 A1*	2/2007	Kapur et al.	435/6
2007/0105167 A1	5/2007	Ausubel et al.	
2013/0260375 A1	10/2013	Cummings et al.	

FOREIGN PATENT DOCUMENTS

EP	1371985	12/2003
WO	WO2008/119332	10/2008
WO	WO2013/151647	10/2013
WO	WO2013/151648	10/2013

OTHER PUBLICATIONS
Ravva et al. (2005) J of microbial meth vol. 63 pp. 305-317.*
Stratagene 1988 catalog.*
Alvarez et al., "Genetic Diversity of *Mycobacterium avium* Isolates Recovered from Clinical Samples and from the Environment:

Molecular Characterization for Diagnostics Purposes", *Journal of Clinical Microbiology*, vol. 46, No. 4, Feb. 13, 2008, 1246-1251.
Bartos et al., "Identification of members of *Mycobacterium avium* species by Accu-Probes, serotyping, and single IS900, IS901, IS1245 and IS901-flanking region PCR with internal standards", *Journal Microbiological Methods*, vol. 64, No. 3, 2006, 333-345.
Cayrou et al., "Genotyping of *Mycobacterium avium* complex organisms using multispacer sequence typing", *Microbiology*, vol. 156, No. 3, Mar. 1, 2010, 687-694.
Covert et al., "Occurrence of nontuberculous mycobacteria in environmental samples", *Applied and Environmental Microbiology*, vol. 65, No. 5, Jun. 1999, 2492-2496.
Falkinham et al., "Factors influencing numbers of *Mycobacterium avium*, *Mycobacterium intracellulare*, and other mycobacteria in drinking water distribution systems", *Applied and Environmental Microbiology*, vol. 67, No. 3, Jun. 2001, 1225-1231.
Higgins et al., "Identification of *Mycobacterium* spp. of veterinary importance using *rpOB* gene sequencing", *BMC Veterinary Research*, vol. 7, No. 1, Jan. 1, 2011, 1-14.
Hilborn et al., "Molecular comparison of *Mycobacterium avium* isolates from clinical and environmental sources", *Applied and Environmental Microbiology*, vol. 74, No. 15, Aug. 2008, 4966-4968.
Intl Application No. PCT/US2013/029072, International Search Report and Written Opinion mailed May 29, 2013, 1-14.
Intl Application PCT/US2013/029093, International Search Report and Written Opinion mailed May 27, 2013, 1-13.
Johansen et al., "Distribution of IS1311 and IS1245 in *Mycobacterium avium* Subspecies Revisited", *Journal of Clinical Microbiological*, vol. 45, No. 5, May 1, 2005, 2500-2502.
Le Dantec et al., "Occurrence of *Mycobacteria* in Water Treatment Lines and in Water Distribution Systems", *Applied and Environmental Microbiology*, vol. 68, No. 11, Nov. 2002, 5318-5325.
Pate et al., "IS1245 RFLP-based genotyping study of *Mycobacterium avium* subsp. *hominissuis* isolates from pigs and humans", *Comparative Immunology, Microbiology & Infectious Diseases*, vol. 31, No. 6, Nov. 1, 2008, 537-550.
Santos et al., "Detection and identification of mycobacteria in the Lisbon water distribution system", *Water and Science Technology*, vol. 52, No. 8, 2005, 177-180.
Shin et al., "Efficient Differentiation of *Mycobacterium avium* Complex Species and Subspecies by Use of Five-Target Multiplex PCR", *Journal of Clinical Microbiology*, vol. 48, No. 11, Nov. 2010, 4057-4062.

* cited by examiner

Primary Examiner — Suchira Pande

(57) **ABSTRACT**

Disclosed are compositions, assays, methods, diagnostic methods, kits and diagnostic kits for the specific and differential detection of *Mycobacterium avium* subsp. *paratuberculosis* from samples including veterinary samples, clinical samples, food samples, forensic sample, an environmental sample (e.g., soil, dirt, garbage, sewage, air, or water), including food processing and manufacturing surfaces, or a biological sample.

9 Claims, No Drawings

1

COMPOSITIONS AND METHODS FOR DETECTION OF *MYCOBACTERIUM AVIUM* PARATUBERCULOSIS

CROSS-REFERENCE(S) TO RELATED APPLICATION

This application claims priority under 35 U.S.C. §119(e) of U.S. Provisional Application No. 61/619,196, filed Apr. 2, 2012, the entire contents of which are incorporated herein by reference.

FIELD

The present teachings relate to compositions, methods and kits for specific detection, diagnosis and differentiation of *Mycobacterium avium* subspecies *paratuberculosis* (MAP) for differentially detecting MAP from other non-MAP microorganisms including other *Mycobacterium avium* subspecies (such as *M. avium* subsp. *hominissuis* (MAH), *Mycobacterium avium* subsp. *avium* (MAA), and *Mycobacterium avium* subsp. *silvaticum* (MAS)) in a variety of mammalian host species.

EFS INCORPORATION PARAGRAPH FOR SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. The ASCII copy, created on Feb. 28, 2013, is named LT00645_SL.txt and is 27,112 bytes in size.

BACKGROUND

Mycobacterium avium subsp. *paratuberculosis* (MAP) is a bacterium that causes Johne's Disease (chronic granulomatous enteritis of the small intestine) in livestock. Johne's disease results in decreased milk production, fetal loss, diarrhea and early death resulting in substantial economic loss to the livestock and dairy industry.

There is a need for assays for the rapid, sensitive and specific detection of infectious pathogens for differential identification of MAP from other bacteria and *Mycobacterium avium* subspecies. Such assays are sought for diagnostic identification of animals infected by MAP.

SUMMARY

The present disclosure, in some embodiments, describes compositions, methods and kits for specific and/or differential detection and diagnosis of MAP from other *Mycobacterium avium* subspecies and/or other non-MAP organisms. Differentially detecting MAP microorganisms from non-MAP organisms in a variety of mammalian host species provides elimination of false positives and superior diagnostic tests which may include one-step diagnosis.

In some embodiments, the present disclosure describes nucleic acid target sequences that are unique to MAP and not present in other *Mycobacterium avium* subspecies. These unique nucleic acid sequences comprise regions of MAP genomes including non-coding regions, coding regions, genes, alleles and variants thereof and/or portions and/or fragments and/or complements thereof. In some embodiments, a nucleic acid target sequence unique to MAP comprises the sequence of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5, fragments

2

thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences. In some embodiments, fragments comprise fragments having at least 10, at least 20, at least 25, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100, contiguous nucleotides, including all values in-between at least 10 and at least 100 contiguous nucleotides (such as 11, 12, . . . 21, 27 . . . 29 etc.), of any of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5. Sequences having 80%-99% sequence identity may include nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and include all values in-between 80% and 99% not explicitly disclosed.

In some embodiments, compositions comprising isolated nucleic acid sequences of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences are described. In some embodiments, isolated nucleic acid fragments comprise fragments having at least 10, at least 20, at least 25, or at least 30 contiguous nucleotides of any of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5. Isolated nucleic acids having 80%-99% sequence identity include nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and include all values in-between 80% and 99% not explicitly disclosed.

Some embodiments describe oligonucleotide primers for use in a nucleic acid amplification method (such as but not limited to PCR) for the detection of target nucleic acid sequences that are unique to MAP. Oligonucleotide primers of the present disclosure comprise at least two primers, having at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting primer sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 49, complements thereof and sequences having about 90% identity to the foregoing sequences.

In some embodiments, primer sets are described having at least 1 or at least 2 primer sets, wherein each primer set has at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. In some embodiments, primer sets may be nested primers. In some embodiments, primer sets or primers may be degenerate primers.

Some embodiments describe oligonucleotide probe sequences for use in detection of target nucleic acid sequences and/or amplified target sequences that are unique to MAP. Oligonucleotide probes of the present disclosure are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting probe sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

In some embodiments, isolated nucleic acid sequence compositions of the disclosure, including primers, probes and polynucleotides/oligonucleotides, can further comprise one or more label, such as, but not limited to, a dye, a radioactive isotope, a chemiluminescent label, a fluorescent moiety, a bioluminescent label an enzyme, and combinations thereof.

The present disclosure, in some embodiments, describes methods (assays) utilizing molecular methods such as nucleic acid sequence specific amplification and detection that offer significant improvements in speed, sensitivity and specificity over traditional microbiological methods. Embodiments relate to design and development of molecular detection assays comprising identification of one or more target nucleic acid sequence that is present in a MAP organism to be detected and absent or divergent in organisms not to be detected. Some embodiments further relate to designing primers, including designing degenerate primers, that can bind to and amplify one or more target nucleic acid sequences encoding for a MAP-specific target nucleic acid and/or a complement and/or a fragment thereof, and using the designed primers to amplify and detect such target nucleic acid sequences.

In some embodiments, methods of detecting in a sample the presence of a MAP microorganism are disclosed. In some embodiments, methods of detecting the presence of a MAP strain are described.

The specification also discloses methods for detection of a MAP organism from a sample and methods to exclude the presence of MAP organism in a sample, wherein the detection of at least one nucleic acid sequence that is specific for a MAP organism is indicative of the presence of a MAP organism and the absence of detection of any nucleic acid sequence unique to a MAP organism is indicative of the absence of a MAP organism in the sample.

In some embodiments, methods of detection of MAP comprise detection of one or more target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto.

Accordingly, a method of the disclosure, in some embodiments, can comprise detecting, in a sample, at least one (or more) nucleic acid sequence(s) having at least 10 to at least 25 contiguous nucleic acids of one (or more) MAP specific nucleic acid targets and/or complementary sequences thereof, wherein detection of at least one nucleic acid sequence indicates the presence of an MAP organism in the sample. Non-limiting examples of MAP specific nucleic acids that can be detected include SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5,

fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences (see also Table 6). Methods of detection can also comprise identification steps and can further comprise steps of sample preparation. Such embodiments are described in detail in sections below.

One embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of an MAP-specific target nucleic acid and/or a fragment or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof by contacting nucleic acids present in the sample with at least one primer set, having one forward primer and one reverse primer that can hybridize to and amplify the MAP-specific nucleic acid and/or a fragment and/or a complement thereof, under conditions suitable for amplification, and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid amplified by the primers confirms the presence of a MAP organism in a sample. In some embodiments, more than one primers can be used to amplify one or more amplification products.

Non-limiting exemplary primer pairs comprise a primer pair such as SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; complements thereof and sequences having about 90% identity to the foregoing sequences, wherein one of the two primers of each primer set is a forward primer and the other is a reverse primer.

Accordingly, an example embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of a MAP-specific nucleic acid and/or a fragment and/or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof contacting nucleic acids present in the sample with at least one primer set, each primer set having one forward primer and one reverse primer, comprising the at least one primer set selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49, wherein the contacting is performed under conditions suitable for a nucleic acid amplification reaction; and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in a sample.

In one embodiment a method for detection of a MAP organism from a sample comprises: detecting the presence of one or more MAP specific nucleic acids and/or a fragment or a complement thereof comprising: contacting nucleic acids present in a sample with a multiplex of primer sets each primer set having one forward primer and one reverse primer, comprising a first primer set, a second primer set, and optionally a third (a fourth etc.) primer sets, under conditions opti-

mal for an amplification reaction to obtain one or more amplified nucleic acids; and detecting the one or more amplified nucleic acids, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in the sample.

Some embodiments describe a method for detection of a MAP organism from a sample comprising: detecting the presence of one or more MAP-specific nucleic acids including detecting a first MAP specific nucleic acid and/or a fragment or a complement thereof comprising: a) amplifying from a sample a first MAP specific nucleic acid and/or a fragment or a complement thereof by contacting nucleic acids present in the sample with at least a first primer set, having one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid and/or a fragment or a complement thereof; and b) amplifying simultaneously from the same sample a second MAP specific nucleic acid and/or a fragment or a complement thereof by simultaneously contacting nucleic acids present in the sample with at least a second primer set, having one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid and/or a fragment or a complement thereof, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction; and detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) and/or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample. In some embodiments a first and a second amplification product can be detected to indicate the presence of a MAP organism in the sample. Furthermore, the process can be optionally repeated to detect a third, a fourth and additional MAP specific nucleic acid molecules.

In other embodiments, not detecting any amplified product using one or more methods described above can be used to exclude the presence of a MAP organism in a sample.

Amplification reactions can comprise a PCR amplification, an end-point determination, a quantitative amplification, a real-time PCR such as a SYBR® Green Assay or a TaqMan® Assay.

Detection can be performed by a variety of methods, such as but not limited to, a nucleic acid amplification reaction such as described in the paragraph above. Detection in some embodiments can be performed by hybridization using probes specific to amplified nucleic acid sequences encoding a MAP specific target nucleic acid sequence. Combinations of amplification and hybridization can be used for detection according to some embodiments. Example probe sequences of the disclosure that can be used for detecting in a method of the present disclosure are SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

In some embodiments, probes can be used to detect and/or to identify a nucleic acid sequence amplified described in the methods above. For example, a probe comprising SEQ ID NO:8 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 6 and SEQ ID NO 7; a probe having SEQ ID NO:11 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 8 and SEQ ID NO 9; a probe having SEQ ID NO 14 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 12 and SEQ ID NO: 13; a probe comprising SEQ ID NO:17 can be used to detect

an amplified nucleic acid amplified by using primer set having SEQ ID NO: 15 and SEQ ID NO 16; a probe having SEQ ID NO:20 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 18 and SEQ ID NO 19; a probe having SEQ ID NO 23 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 21 and SEQ ID NO:22; a probe comprising SEQ ID NO:26 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 24 and SEQ ID NO 25; a probe having SEQ ID NO:29 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 27 and SEQ ID NO 28; a probe having SEQ ID NO 32 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 30 and SEQ ID NO:31; a probe comprising SEQ ID NO:35 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 33 and SEQ ID NO 34; a probe having SEQ ID NO:38 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 36 and SEQ ID NO 37; a probe having SEQ ID NO 41 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 39 and SEQ ID NO:40; a probe comprising SEQ ID NO:44 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 42 and SEQ ID NO 43; a probe having SEQ ID NO:47 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 45 and SEQ ID NO 46; and a probe having SEQ ID NO 50 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 48 and SEQ ID NO:49.

In one embodiment, disclosed is an assay for the detection of a MAP organism in a sample comprising a) hybridizing a first pair (or set) of PCR primers selected from a row in the Table 3 described as forward primers and reverse primers (selected from primer sets described in the paragraph above), and complements thereof to at least a first MAP target polynucleotide sequence and/or fragment thereof; b) amplifying the at least first target MAP polynucleotide sequences; and c) detecting the at least first and the at least second amplified target polynucleotide sequence products; wherein the detection of the at least first amplified target polynucleotide sequence product and the detection of the at least second amplified target polynucleotide sequence product is indicative of the presence of a MAP organism in the sample. The method can also comprise hybridizing a second pair of PCR primers selected from another row in Table 3 described as forward primers and reverse primers, and complements thereof to at least a second target MAP polynucleotide sequence and/or fragment thereof.

In further embodiments, the detection can comprise using hybridization with one or more probes. Probes that can be used are described in Table 3 where different probes specific to different amplified MAP target sequences. Primer-probe combinations are outlined in the Table 3 and also set forth in the sections above (for example, amplification product amplified using primers SEQ ID NO:6 and SEQ ID NO:7 can be detected using probe SEQ ID NO:8).

In some embodiments, hybridization can comprise at least a first probe and a second probe, the first probe further comprising a first label and the second probe further comprising a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye.

In some embodiments, any detection method described above can further comprise preparing a sample for PCR amplification (prior to hybridizing), for example, but not limited to (1) bacterial enrichment, and/or (2) separation of bacterial cells from the sample, and/or (3) cell lysis, and/or (4) nucleic acid extraction. In some embodiments, a detection method can comprise DNA isolation from a sample.

Samples may include without limitation, veterinary samples, animal-derived samples, clinical samples, food/beverage samples, water samples, and environmental sample. Veterinary samples may be derived from animals such as but not limited to all ruminants, cattle, sheep, bison, deer, rabbits, foxes and birds including poultry birds, other domesticated and wild birds.

Methods can include multiplex assays such as polymerase chain reactions, wherein hybridizing and amplifying of the first pair of polynucleotide primers occurs in a first vessel and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a second vessel, or hybridizing and amplifying of the first pair of polynucleotide primers and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a single vessel, the detection is a real-time assay, the real-time assay is a SYBR® Green dye assay or a TaqMan® assay. Methods can also comprise using additional primers such as a third primer pair and a fourth primer pair and so on.

A method of the disclosure can further comprise providing a first probe and a second probe (and additional probes such as a third probe and a fourth probe and so on), wherein the first and second probes are different from each other, the first probe operable to identify the first amplified target polynucleotide sequence and the second probe operable to identify the second amplified target nucleotide sequence, the first probe further comprises a first label and the second probe further comprises a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye; and hybridizing the first and second probes to the PCR amplified fragments to detect the presence of the first amplified target polynucleotide sequence and the second amplified target polynucleotide sequence from the sample.

Some embodiments describe kits suitable for identifying the presence of a MAP organism. Such a kit can comprise at least one set of oligonucleotide primers for use in a PCR process for the detection of a MAP target nucleic acid sequence.

A first probe can further comprise a first label and a second probe further comprise a second label, both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, and the first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye. Kits can without limitation contain other buffers, molecular bio reagents, and one or more container means for kit components.

In the following description, certain aspects and embodiments will become evident. It should be understood that a given embodiment need not have all aspects and features

described herein. It should be understood that these aspects and embodiments are merely exemplary and explanatory and are not restrictive of the invention. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

DETAILED DESCRIPTION

The present disclosure, in some embodiments, describes compositions, kits and methods for detection of MAP. Some embodiments describe specific detection of MAP from a sample even when non-MAP *Mycobacterium avium* organisms are present in the sample.

MAP is a bacterium that infects livestock and several herd animals and causes Johne's Disease which is characterized by chronic granulomatous enteritis of the small intestine. In livestock especially dairy animals such as cows, sheep and bison, Johne's disease results in decreased milk production, fetal loss, diarrhea and early death resulting in substantial economic loss to the livestock industry. Existing assays that are used to detect the presence of MAP in samples obtained from animals and to diagnose Johne's disease are based on the detection of insertion sequences present in MAP. MAP is also believed to be linked to human Crohn's disease.

Mycobacterium avium complex (MAC) consists of multiple *Mycobacterium avium* subspecies that can be found as environmental contaminants in soil and water (Covert et al., 1999; Falkinham et al., 2001; Le Dantec et al., 2002; Santos et al., 2005; Hilborn et al., 2008), as well as infectious agents for animals such as pigs, cattle, sheep and birds (C. Cayrou et al 2010). Organisms classified in the MAC complex include *Mycobacterium* subsp. *avium*, *Mycobacterium avium* subsp. *paratuberculosis*, *Mycobacterium avium* subsp. *silvaticum*, and *Mycobacterium* subsp. *hominissuis*. Symptoms of infection vary by host species and MAC organism. Common methods used for identifying MAC subspecies include DNA sequencing, particularly of the 16S and rpoB genes, and RFLP (restriction fragment length polymorphism) analysis (C. Cayrou et al 2010).

In some cases, certain nucleic acid insertion sequences have been associated with some *Mycobacterium* subspecies and this has been used to detect these organisms. For example, insertion sequences known as ISMAP02 and IS900 are detected in existing PCR-based MAP detection assays which rely on detecting or determining the presence of the ISMAP02 or IS900 target nucleic acid sequences in a sample.

Next-generation sequencing, using an Ion Torrent Personal Genome Machine (PGM) was performed herein to get complete genome sequences for 16 *Mycobacterium avium* subspecies. Sequences were obtained for one *M. avium* subsp. *silvaticum* sample; four *M. avium* subsp. *avium* samples; two *M. avium* subsp. *hominissuis* samples; three *M. avium* samples with an unknown subtype; and six *M. avium* subsp. *paratuberculosis* samples. The sequenced genomes were analyzed to look for the presence of all previously known insertion sequences as well as for new highly specific single-gene targets. A partial ISMAP02 sequence was found in all organisms indicating that ISMAP02 is not an optimal sequence to use for a MAP-specific assay. Results were also confirmed with qPCR testing using the TaqMan® MAP reagents. Results of this analysis are presented in Table 1.

TABLE 1

Sample	Pathogen	Host	1:100X Sample Dilution		
			MAP CT 1	MAP CT 2	ISMAP02 Sequencing Results
ATCC 49884	<i>M. silvaticum</i>	Wood pigeon	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240)
10-5581	<i>M. avium</i> subsp. <i>avium</i>	Asian Elephant	16.94	16.96	ISMAP02 positive, full sequence
10-9275	<i>M. avium</i> subsp. <i>avium</i>	Red Tailed Hawk	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240)
11-4751	<i>M. avium</i> subsp. <i>avium</i>	Ruddy Duck	Undetermined	Undetermined	parital ISMAP02 sequence (position 1-240 and 360-519)
05-4293	<i>M. avium</i>	Avian broadbill from Dallas Zoo	Undetermined	37.45	parital ISMAP02 sequence (position 1-240)
10-4249	<i>M. avium</i> subsp. <i>hominissuis</i>	White lipped deer	14.78	14.76	ISMAP02 positive, full sequence
10-5606	<i>M. avium</i> subsp. <i>hominissuis</i>	Pig	30.24	30.95	parital ISMAP02 sequence (position 1-240 and 360-519)
10-5560	Unknown MAC subtype	Pig	15.49	15.48	ISMAP02 positive, full sequence
09-5983	Unknown MAC subtype	Dog	16.43	16.43	ISMAP02 positive, full sequence
11-0986	Unknown MAC subtype	deer	16.22	16.23	ISMAP02 positive, full sequence
10-4404	<i>M. avium</i> subsp. <i>paratuberculosis</i>	bison	14.38	14.40	ISMAP02 positive, full sequence
10-5975	<i>M. avium</i> subsp. <i>paratuberculosis</i>	bison	14.74	14.79	ISMAP02 positive, full sequence
10-5864	<i>M. avium</i> subsp. <i>paratuberculosis</i>	cattle	14.44	14.45	ISMAP02 positive, full sequence
10-8425	<i>M. avium</i> subsp. <i>paratuberculosis</i>	cattle	14.74	14.70	ISMAP02 positive, full sequence
08-8281	<i>M. avium</i> subsp. <i>paratuberculosis</i>	sheep	14.82	14.82	ISMAP02 positive, full sequence
11-1786	<i>M. avium</i> subsp. <i>paratuberculosis</i>	sheep	14.76	14.75	ISMAP02 positive, full sequence

Other insertion sequences, including IS901 were also found to be non-specific for MAP. The IS901 sequence was present in the *M. avium* subsp. *silvaticum* sequence, and two of the *M. avium* subsp. *avium* sequences (isolates 11-4751 and 10-9275).

In one embodiment of the current teachings, bioinformatic and direct DNA sequencing comparisons of several *Mycobacterium avium* species were conducted to identify MAP-specific target nucleic acids sequences. Alignment of these sequences using custom algorithms identified several MAP-specific target regions to which primer pairs of the disclosure were designed for each identified MAP-specific target region to specifically amplify only the unique MAP-specific target sequences against both inclusion (organism to be detected, i.e., MAP organisms) and exclusion genomes (organisms not to be detected, non-MAP organisms).

Exemplary embodiments of nucleic acid target sequence unique to MAP identified herein include nucleic acids having SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences.

Genome coordinates for SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, are described in Table 2 below in relation to the MAP genome described in GenBank accession number NC_002944. For example, nucleic acid target sequence unique to MAP comprising the sequence of SEQ ID. NO: 1 corresponds to a 1,863 bp nucleic acid sequence starting from position 238,936 and continuing through position 240,798 of the MAP genome described as NC_002944. SEQ ID. NOs: 1-5 are described here and in the Sequence Listing below.

GenBank accession number NC_002944 can be found at http://www.ncbi.nlm.nih.gov/nuccore/nc_002944 referenced as "*Mycobacterium avium* subsp. *paratuberculosis* K-10, complete genome" 16 Sep. 2011, (1. Li, L., Bannan-

tine, J. P., Zhang, Q., Amonsin, A., May, B. J., Alt, D., Banerji, N., Kanjilal, S. and Kapur, V. "The complete genome sequence of *Mycobacterium avium* subspecies *paratuberculosis*," Proc. Natl. Acad. Sci. U.S.A., 102 (35), 12344-12349 (2005); 2. CONSRM NCBI Genome Project Direct Submission, Submitted (11 Sep. 2004) National Center for Biotechnology Information, NIH, Bethesda, Md. 20894, USA; 3. Li, L., Bannantine, J., Zhang, Q., Amonsin, A., Alt, D. and Kapur, V., Direct Submission, Submitted (5 Sep. 2003) Biomedical Genomics Center, University of Minnesota, 1971 Commonwealth Ave., St. Paul, Minn. 55108, USA).

TABLE 2

MAP genome coordinates (NC_002944)				
Assay ID Number	Signature left	Signature right	Signature length	SEQ ID. NO:
61801, 61802, 61803, 61805	238936	240798	1863	SEQ ID. NO: 1
61808, 61813, 61815, 61816, 61817	3088062	3089612	1551	SEQ ID. NO: 2
61819, 61822, 61823, 61825, 61826, 61829	4168103	4169949	1847	SEQ ID. NO: 3
	4169965	4174680	4716	SEQ ID. NO: 4
	4194685	4197076	2392	SEQ ID. NO: 5

In some embodiments, compositions of the disclosure comprise isolated nucleic acids having SEQ ID. NO: 1, SEQ ID. NO: 2, SEQ ID. NO: 3, SEQ ID. NO: 4 and/or SEQ ID. NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences. Sequences hav-

ing 80%-99% sequence identity include isolated nucleic acid sequences that have 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% and 99% sequence identity to any one of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and/or fragments thereof and/or complementary sequences thereof and include all values in-between 80% and 99% not explicitly disclosed. Fragments include oligonucleotides or polynucleotides having at least 10 contiguous nucleotide sequences, or at least 20 contiguous nucleotide sequences, or at least 25 contiguous nucleotide sequences, or at least 30 contiguous nucleotide sequences in any part of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5 and/or fragments thereof and/or complementary sequences thereof and sequences having at least 90% identity to the foregoing sequences.

Compositions of the disclosure comprise probe and/or primer sequences that are specific to hybridize to (and amplify in the case of primers or detect in the case of probes) one or more target nucleic acid sequences that are unique to a MAP organism. A target nucleic acid sequence unique to a MAP organism may include a gene, a non-coding region, an allele or a complement thereof that is present in a MAP organism but absent from a non-MAP organism. In some embodiments, a target nucleic acid sequence unique to a MAP organism is absent from other *Mycobacterium avium* subspecies which are otherwise very similar or closely related. Target nucleic acid sequences as described in this disclosure can comprise portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto and/or from about 70% to about 90% identity to a target nucleic acid sequence unique to a MAP organism and can include a gene, a non-coding region, an allele or a complement thereof that is present in a MAP organism but absent from a non-MAP organism. Some example probe and primer sequences are described in Table 3 in sections ahead.

Several programs for designing primers such as Primer3 (Steve Rozen and Helen J. Skaletsky (2000) "Primer3" on the World Wide Web for general users and for biologist programmers as published in: Krawetz S, Misener S (eds) *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa, N.J., pp 365-386), Primer Express® software (Applied Biosystems), and OLIGO 7 (Wojciech Rychlik (2007). "OLIGO 7 Primer Analysis Software". *Methods Mol. Biol.* 402: 35-60) and variations thereof can be used for primer designing. Custom algorithms were used in the present disclosure in conjunction with the new uniquely identified targets to arrive at primer and probe compositions described herein. The presently designed PCR primers and probes are useful in assays such as real-time PCR assays to specifically and unambiguously detect MAP organisms with great sensitivity.

In some embodiments, exemplary non-limiting compositions comprising primer sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence of SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 19, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 40, SEQ ID NO: 42, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 49, complements thereof and sequences having about 90% identity to the foregoing sequences.

In some embodiments, primer sets are described having at least 1 or at least 2 primer sets, wherein each primer set has at least a forward primer and at least a reverse primer, that are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. In some embodiments, primer sets can be nested primers. In some embodiments, primer sets or primers can be degenerate primers. In some embodiments the present disclosure describes designing multiplex primers that are suitable for multiplex PCR assays.

Some embodiments describe oligonucleotide probe sequences for use in detection of target nucleic acid sequences and/or amplified target sequences that are unique to MAP. Oligonucleotide probes of the present disclosure are operable to hybridize to target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity, and/or sequences having at least about 80% identity thereto. Some exemplary non-limiting probe sequences of the present disclosure comprise isolated nucleic acid sequences having the nucleotide sequence, of SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

Isolated nucleic acid sequence compositions of the disclosure, including primers and probes according to the disclosure, can further comprise one or more label, such as, but not limited to, a dye, a radioactive isotope, a chemiluminescent label, a fluorescent moiety, a bioluminescent label an enzyme, and combinations thereof.

The specification also discloses methods for detection and/or diagnosis of a MAP organism from a sample, wherein the detection of at least one unique or target nucleic acid sequence that is expressed in a MAP organism is indicative of the presence of a MAP organism and the absence of detection of any nucleic acid sequence unique to MAP organism is indicative of the absence of a MAP organism in the sample. Accordingly, some methods of the disclosure exclude the presence of a MAP organism in a sample absent of detection of any nucleic acid sequence unique to MAP organism in the sample. In some embodiments, methods of detecting and/or diagnostically determining in a sample the presence of a MAP microorganisms in the presence of other *Mycobacterium avium* organisms of different subspecies and subtypes are disclosed.

Diagnosis and diagnostic methods of the disclosure can include detecting the presence of a MAP organism to diagnose a MAP organism induced disease or condition. Any animal or human disease or condition caused by a MAP organism is contemplated to be diagnosable by the detection methods disclosed herein that can detect the presence of a MAP organism as defined herein. Non limiting examples include diagnosis of John's disease and Crohn's disease.

A method of detection and/or diagnosis of MAP comprises detection of one or more target nucleic acid sequences that are unique to MAP including portions and/or fragments and/or complements thereof and/or sequences having at least about 90% identity to, and/or sequences having at least about 80% identity thereto. Accordingly, a method of the disclosure, in some embodiments, can comprise detecting, in a sample, at least one (or more) nucleic acid sequence(s) having at least 10 to at least 25 nucleic acids of one (or more) MAP specific

nucleic acid targets and/or complementary sequences thereof, wherein detection of at least one nucleic acid sequence indicates the presence of a MAP organism in the sample. Non-limiting examples of MAP specific nucleic acids that can be detected include SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4 and/or SEQ ID NO: 5, fragments thereof, complementary sequences thereof as well as sequences having from about 80% to about 99% sequence identity to one of these sequences.

Methods of detection and/or diagnosis of MAP in a sample can also comprise identification steps and can further comprise steps of sample preparation. Preparing a sample for PCR amplification (prior to hybridizing with primers), can, in some embodiments, comprise one or more steps such as, but not limited to (1) bacterial enrichment, and/or (2) separation of bacterial cells from the sample (isolation of DNA from sample), and/or (3) cell lysis, and/or (4) nucleic acid (NA) extraction.

Detection of MAP organisms by the use of methods described herein, in some embodiments, is by an amplification reaction such as a polymerase chain reaction for rapid detection. In general methods of the disclosure include comparing for presence of an MAP organism using suitable controls, for example, typically, an internal positive control can be used in a PCR reaction which will have a detectable signal/positive result and a suitable negative control that will have no detectable signal. Amplification reactions can comprise one or more of the following: a PCR amplification, an end-point determination, a quantitative amplification, a real-time PCR such as a SYBR® Green Assay and/or a TaqMan® Assay.

Methods can also comprise detecting at least one amplified nucleic acid by hybridization, mass spectrometry, nanostring, microfluidics, chemiluminescence, enzyme technologies and combinations thereof.

In one embodiment, a method for detection of a MAP organism from a sample comprises: detecting the presence of an MAP-specific target nucleic acid and/or a fragment or a complement thereof comprising: amplifying an MAP-specific nucleic acid and/or a fragment and/or a complement thereof by contacting nucleic acids present in the sample with at least one primer set, having one forward primer and one reverse primer that can hybridize to and amplify the MAP-specific nucleic acid and/or a fragment and/or a complement thereof, under conditions suitable for amplification, and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid amplified by the primers confirms the presence of a MAP organism in a sample.

In some embodiments, more than one primers can be used to amplify one or more amplification products. Non-limiting exemplary primer pairs comprise a primer pair such as SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; complements thereof and sequences having about 90% identity to the foregoing sequences, wherein one of the two primers of each primer set is a forward primer and the other is a reverse primer.

One example embodiment method for detection (and/or diagnosis) of a MAP organism from a sample comprises: detecting the presence of a MAP-specific nucleic acid and/or

a fragment and/or a complement thereof comprising: amplifying a MAP-specific nucleic acid and/or a fragment and/or a complement thereof contacting nucleic acids present in the sample with at least one primer set, each primer set having one forward primer and one reverse primer, comprising the at least one primer set selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49, wherein the contacting is performed under conditions suitable for an nucleic acid amplification reaction; and detecting an amplified nucleic acid, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in a sample.

In one embodiment method for detection of a MAP organism from a sample comprises: detecting the presence of one or more MAP specific nucleic acids and/or a fragment or a complement thereof comprising: contacting nucleic acids present in a sample with a multiplex of primer sets each primer set having one forward primer and one reverse primer, comprising a first primer set, a second primer set, and optionally a third (a fourth etc.) primer sets, under conditions optimal for an amplification reaction to obtain one or more amplified nucleic acids; and detecting the one or more amplified nucleic acids, wherein detecting an amplified nucleic acid using the primers confirms the presence of a MAP organism in the sample.

Some embodiments describe a method for detection of a MAP organism from a sample comprising: detecting the presence of one or more MAP-specific nucleic acids including detecting a first MAP specific nucleic acid and/or a fragment or a complement thereof comprising: a) amplifying from a sample a first MAP specific nucleic acid and/or a fragment or a complement thereof by contacting nucleic acids present in the sample with at least a first primer set, having one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid and/or a fragment or a complement thereof; and b) amplifying simultaneously from the same sample a second MAP specific nucleic acid and/or a fragment or a complement thereof by simultaneously contacting nucleic acids present in the sample with at least a second primer set, having one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid and/or a fragment or a complement thereof, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction; and c) detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) and/or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample. In some embodiments a first and a second amplification product are detected to indicate the presence of a MAP organism in the sample. Furthermore, the method can be optionally repeated to detect a third, a fourth and additional MAP specific nucleic acid molecules.

In other embodiments, not detecting any amplified product using one or more methods described above can be used to exclude the presence of a MAP organism in a sample.

Methods of the disclosure can also use other detection methods in addition to nucleic acid amplification reactions described in the paragraphs above. Detection in some

embodiments can be performed by hybridization using probes specific to MAP specific target nucleic acid sequence. Accordingly, probes specific to detect one or more target MAP nucleic acids can be used in a sample, under conditions to promote sequence specific hybridization, to detect presence of a MAP nucleic acid in the sample. In some embodiments, hybridization comprises using at least a first probe and a second probe, the first probe further comprising a first label and the second probe further comprising a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye.

In some embodiments, combinations of amplification and hybridization can be used for detection and amplified nucleic acid sequences encoding a MAP specific target nucleic acid sequence can be detected as MAP-specific amplification products using MAP specific probes. Some non-limiting example probe sequences of the disclosure that can be used for detecting in a method of the present disclosure are described in SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences.

In one embodiment, a method for the detection of a MAP organism in a sample comprises: a) hybridizing a first pair (or set) of PCR primers selected from a row in the Table 3 described as forward primers and reverse primers (selected from primer sets described in the paragraph above), and complements thereof to at least a first MAP target polynucleotide sequence and/or fragment thereof; b) amplifying the at least first target MAP polynucleotide sequences; and c) detecting the at least first and the at least second amplified target polynucleotide sequence products; wherein the detection of the at least first amplified target polynucleotide sequence product and the detection of the at least second amplified target polynucleotide sequence product is indicative of the presence of a MAP organism in the sample. The method can also comprise hybridizing a second pair of PCR primers selected from another row in Table 3 described as forward primers and reverse primers, and complements thereof to at least a second target MAP polynucleotide sequence and/or fragment thereof.

In further embodiments, the detection can comprise using hybridization with one or more probes. Some example probes are described in Table 3 where different probes specific to different amplified MAP target sequences are described. Primer-probe combinations are also outlined in the Table 3. For example, in some non-limiting embodiments, a probe comprising SEQ ID NO:8 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 6 and SEQ ID NO 7; a probe having SEQ ID NO:11 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 8 and SEQ ID NO 9; a probe having SEQ ID NO 14 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 12 and SEQ ID NO: 13; a probe comprising SEQ ID NO:17 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 15 and SEQ ID NO 16; a probe having SEQ ID NO:20 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 18 and SEQ ID NO 19; a probe having SEQ ID NO 23 can be used to detect an amplified nucleic acid

amplified by using primer set having SEQ ID NO: 21 and SEQ ID NO:22; a probe comprising SEQ ID NO:26 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 24 and SEQ ID NO 25; a probe having SEQ ID NO:29 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 27 and SEQ ID NO 28; a probe having SEQ ID NO 32 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 30 and SEQ ID NO:31; a probe comprising SEQ ID NO:35 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 33 and SEQ ID NO 34; a probe having SEQ ID NO:38 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 36 and SEQ ID NO 37; a probe having SEQ ID NO 41 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 39 and SEQ ID NO:40; a probe comprising SEQ ID NO:44 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 42 and SEQ ID NO 43; a probe having SEQ ID NO:47 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 45 and SEQ ID NO 46; and a probe having SEQ ID NO 50 can be used to detect an amplified nucleic acid amplified by using primer set having SEQ ID NO: 48 and SEQ ID NO:49.

In some embodiments of the present methods, one assay alone may not be definitive for detecting a MAP organism due to genomic similarity between the genomic regions of other non-MAP organisms. Yet, when two (or more) assays such as but not limited to the assays shown in Table 3 are used either in parallel or as a multiplex assay, e.g., in a real-time TaqMan® assay, for example, where each probe in each of the two (or more) assays has a different label for distinguishing results on a real-time PCR instrument, e.g., a 7500 Fast Real-Time PCR System (Applied Biosystems), a positive result from such an assay is indicative of the presence of a MAP organism. Such dual or multiplex (more than 2 assay sets) assay approach can be used to detect and distinguish MAP organism. Some embodiments describe detecting at least two (or more) of MAP-specific target nucleic acid target regions as positive identification of a MAP organism.

Methods of the disclosure include multiplex assays such as polymerase chain reactions, wherein hybridizing and amplifying of the first pair of polynucleotide primers occurs in a first vessel and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a second vessel, or hybridizing and amplifying of the first pair of polynucleotide primers and the hybridizing and amplifying of the second pair of polynucleotide primers occurs in a single vessel, the detection is a real-time assay, the real-time assay is a SYBR® Green dye assay or a TaqMan® assay. Methods also comprise using additional primers such as a third primer pair and a fourth primer pair and so on.

A method of the disclosure can further comprise providing a first probe and a second probe (and additional probes such as a third probe and a fourth probe and so on), wherein the first and second probes are different from each other, the first probe operable to identify the first amplified target polynucleotide sequence and the second probe operable to identify the second amplified target nucleotide sequence, the first probe further comprises a first label and the second probe further comprises a second label, wherein both labels are selected from a dye, a radioactive isotope, a chemiluminescent label, and an enzyme, the dye comprises a fluorescein dye, a rhodamine dye, or a cyanine dye, the dye is a fluorescein dye and first probe is labeled with FAM™ dye and the second probe is labeled with VIC® dye; and hybridizing the first and

second probes to the PCR amplified fragments to detect the presence of the first amplified target polynucleotide sequence and the second amplified target polynucleotide sequence from the sample.

Compositions and methods of the present disclosure are ideally suited for the preparation of a kit suitable for identifying the presence of a MAP organism, such as for example a diagnostic kit. Such a kit can comprise at least one set of oligonucleotide primers for use in a PCR process for the amplification and detection of a MAP-specific target nucleic acid sequence. Some kits of the disclosure comprise at least two sets of oligonucleotide primers for simultaneous use in a multiplex PCR process for the amplification and detection of MAP-specific target nucleic acid sequences. Kits can additionally comprise one or more reagents such as but are not limited to, buffers, nucleotide triphosphates, DNA polymerases, intercalating dye, primers, probes, salt, and instructions for the use of the kit.

In some embodiments, kit primers can be labeled. A kit comprising multiple pairs of primers can have primer pairs each labeled with different labels that may be detectable separately. Probes comprised in kits of the disclosure can be labeled. If a kit comprises multiple probes each probe can be labeled with a different label to allow detection of different target nucleic acids and/or amplification products that are the targets of each different probe.

An example kit for the detection and/or diagnosis of a MAP organism comprises: at least one pair of forward and reverse PCR primers having primer pairs selected from primer pairs described in Table 3 (i.e., at least one primer set selected from: a first primer set having SEQ ID NO: 6 and SEQ ID NO 7; and/or a second primer set having SEQ ID NO: 9 and SEQ ID NO: 10; and/or a third primer set having SEQ ID NO: 12 and SEQ ID NO:13; and/or a fourth primer set having SEQ ID NO: 15 and SEQ ID NO 16; and/or a fifth primer set having SEQ ID NO: 18 and SEQ ID NO: 19; and/or a sixth primer set having SEQ ID NO: 21 and SEQ ID NO: 22; and/or a seventh primer set having SEQ ID NO: 24 and SEQ ID NO: 25; and/or an eighth primer set having SEQ ID NO: 27 and SEQ ID NO: 28; and/or a ninth primer set having SEQ ID NO: 30 and SEQ ID NO: 31; and/or a tenth primer set having SEQ ID NO: 33 and SEQ ID NO: 34; and/or a eleventh primer set having SEQ ID NO: 36 and SEQ ID NO: 37; and/or a twelfth primer set having SEQ ID NO: 39 and SEQ ID NO: 40; and/or a thirteenth primer set having SEQ ID NO: 42 and SEQ ID NO: 43; and/or a fourteenth primer set having SEQ ID NO: 45 and SEQ ID NO: 46; and/or a fifteenth primer set having SEQ ID NO: 48 and SEQ ID NO: 49, or sequences comprising at least 90% nucleic acid sequence identity thereof, or labeled derivatives thereof). In some embodiments, a kit can also have at least one probe selected from probes described Table 3 (i.e., probe sequences of the disclosure having SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, complements thereof and sequences having about 90% identity to the foregoing sequences, each probe corresponding to the first primer set, second primer set, the third primer set, etc. as described above, respectively, selected based on which primer set is selected).

Another example and kit for the detection and/or diagnosis of a MAP organism comprise: at least two pairs of forward and reverse PCR primers (two primer pairs) selected from primer pairs described in Table 3; and optionally at least two probes selected from probes described Table 3.

A kit of the disclosure can further comprise one or more components such as but not limited to: at least one enzyme, dNTPs, at least one buffer, at least one salt, at least one control nucleic acid sample, loading solution for preparation of the amplified material for electrophoresis, genomic DNA as a template control, a size marker to insure that materials migrate as anticipated in a separation medium, and an instruction protocol and manual to educate a user and limit error in use. Components of kits can be individually and/or in various combinations comprised in one or a plurality of suitable container means.

It is within the scope of these teachings to provide test kits for use in manual applications or test kits for use with automated sample preparation, reaction set-up, detectors or analyzers. In some embodiments, a kit amplification product may be further analyzed by methods such as but not limited to electrophoresis, hybridization, mass spectrometry, nanostring, microfluidics, chemiluminescence and/or enzyme technologies.

For the purposes of interpreting of this specification, the following definitions may apply and whenever appropriate, terms used in the singular will also include the plural and vice versa. In the event that any definition set forth below conflicts with the usage of that word in any other document, including any document incorporated herein by reference, the definition set forth below shall always control for purposes of interpreting this specification and its associated claims unless a contrary meaning is clearly intended (for example in the document where the term is originally used). It is noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the," include plural referents unless expressly and unequivocally limited to one referent. The use of "or" means "and/or" unless stated otherwise. The use of "comprise," "comprises," "comprising," "include," "includes," and "including" are interchangeable and not intended to be limiting. Furthermore, where the description of one or more embodiments uses the term "comprising," those skilled in the art would understand that, in some specific instances, the embodiment or embodiments can be alternatively described using the language "consisting essentially of" and/or "consisting of."

As used herein, the phrase "nucleic acid," "oligonucleotide", and polynucleotide(s)" are interchangeable and not intended to be limiting.

As used herein, the phrase "stringent hybridization conditions" refers to hybridization conditions which can take place under a number of pH, salt and temperature conditions. The pH can vary from 6 to 9, preferably 6.8 to 8.5. The salt concentration can vary from 0.15 M sodium to 0.9 M sodium, and other cations can be used as long as the ionic strength is equivalent to that specified for sodium. The temperature of the hybridization reaction can vary from 30° C. to 80° C., preferably from 45° C. to 70° C. Additionally, other compounds can be added to a hybridization reaction to promote specific hybridization at lower temperatures, such as at or approaching room temperature. Among the compounds contemplated for lowering the temperature requirements is formamide. Thus, a polynucleotide is typically "substantially complementary" to a second polynucleotide if hybridization occurs between the polynucleotide and the second polynucleotide. As used herein, "specific hybridization" refers to hybridization between two polynucleotides under stringent hybridization conditions.

As used herein, the term "polynucleotide" refers to a polymeric form of nucleotides of any length, either ribonucleotides, deoxynucleotides, or peptide nucleic acids (PNA), and includes both double- and single-stranded RNA, DNA, and

PNA. A polynucleotide may include nucleotide sequences having different functions, including, for instance, coding regions, and non-coding regions such as regulatory regions. A polynucleotide can be obtained directly from a natural source, or can be prepared with the aid of recombinant, enzymatic, or chemical techniques. A polynucleotide can be linear or circular in topology. A polynucleotide can be, for example, a portion of a vector, such as an expression or cloning vector, or a fragment. An "oligonucleotide" refers to a polynucleotide of the present invention, typically a primer and/or a probe.

As used herein a "target-specific polynucleotide" refers to a polynucleotide having a target-binding segment that is perfectly or substantially complementary to a target sequence, such that the polynucleotide binds specifically to an intended target without significant binding to non-target sequences under sufficiently stringent hybridization conditions. The target-specific polynucleotide can be e.g., a primer or probe and the subject of hybridization with its complementary target sequence.

The term "target sequence", "target signature sequence" "target nucleic acid", "target" or "target polynucleotide sequence" refers to a nucleic acid present in a MAP organism that is not present in other non-MAP organisms and is unique to MAP. The target sequence can be a polynucleotide sequence that is the subject of hybridization with a complementary polynucleotide, e.g. a primer or probe. The target sequence can be composed of DNA, RNA, an analog thereof, and including combinations thereof. The target sequence may be known or not known, in terms of its actual sequence and its amplification can be desired. The target sequence may or may not be of biological significance. Typically, though not always, it is the significance of the target sequence which is being studied in a particular experiment. As non-limiting examples, target sequences may include regions of genomic DNA, regions of genomic DNA which are believed to contain one or more polymorphic sites, DNA encoding or believed to encode genes or portions of genes of known or unknown function, DNA encoding or believed to encode proteins or portions of proteins of known or unknown function, DNA encoding or believed to encode regulatory regions such as promoter sequences, splicing signals, polyadenylation signals, etc.

As used herein an "amplified target polynucleotide sequence product" refers to the resulting amplicon from an amplification reaction such as a polymerase chain reaction. The resulting amplicon product arises from hybridization of complementary primers to a target polynucleotide sequence under suitable hybridization conditions and the repeating in a cyclic manner the polymerase chain reaction as catalyzed by DNA polymerase for DNA amplification or RNA polymerase for RNA amplification.

As used herein, the "polymerase chain reaction" or PCR is an amplification of nucleic acid consisting of an initial denaturation step which separates the strands of a double stranded nucleic acid sample, followed by repetition of (i) an annealing step, which allows amplification primers to anneal specifically to positions flanking a target sequence; (ii) an extension step which extends the primers in a 5' to 3' direction thereby forming an amplicon polynucleotide complementary to the target sequence, and (iii) a denaturation step which causes the separation of the amplicon from the target sequence (Mullis et al., eds, *The Polymerase Chain Reaction*, BirkHauser, Boston, Mass. (1994). Each of the above steps may be conducted at a different temperature, preferably using an automated thermocycler (Applied Biosystems LLC, a division of Life Technologies Corporation, Foster City,

Calif.). If desired, RNA samples can be converted to DNA/RNA heteroduplexes or to duplex cDNA by methods known to one of skill in the art.

As used herein, "amplifying" and "amplification" refers to a broad range of techniques for increasing polynucleotide sequences, either linearly or exponentially. Exemplary amplification techniques include, but are not limited to, PCR or any other method employing a primer extension step. Other non-limiting examples of amplification include, but are not limited to, ligase detection reaction (LDR) and ligase chain reaction (LCR). Amplification methods may comprise thermal-cycling or may be performed isothermally. In various embodiments, the term "amplification product" includes products from any number of cycles of amplification reactions.

In certain embodiments, amplification methods comprise at least one cycle of amplification, for example, but not limited to, the sequential procedures of: hybridizing primers to primer-specific portions of target sequence or amplification products from any number of cycles of an amplification reaction; synthesizing a strand of nucleotides in a template-dependent manner using a polymerase; and denaturing the newly-formed nucleic acid duplex to separate the strands. The cycle may or may not be repeated.

Descriptions of certain amplification techniques can be found, among other places, in H. Ehrlich et al., *Science*, 252:1643-50 (1991), M. Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press, New York, N.Y. (1990), R. Favis et al., *Nature Biotechnology* 18:561-64 (2000), and H. F. Rabenau et al., *Infection* 28:97-102 (2000); Sambrook and Russell, *Molecular Cloning*, Third Edition, Cold Spring Harbor Press (2000) (hereinafter "Sambrook and Russell"), Ausubel et al., *Current Protocols in Molecular Biology* (1993) including supplements through September 2005, John Wiley & Sons (hereinafter "Ausubel et al.>").

The term "label" refers to any moiety which can be attached to a molecule and: (i) provides a detectable signal; (ii) interacts with a second label to modify the detectable signal provided by the second label, e.g. FRET; (iii) stabilizes hybridization, i.e. duplex formation; or (iv) provides a capture moiety, i.e. affinity, antibody/antigen, ionic complexation. Labeling can be accomplished using any one of a large number of known techniques employing known labels, linkages, linking groups, reagents, reaction conditions, and analysis and purification methods. Labels include light-emitting compounds which generate a detectable signal by fluorescence, chemiluminescence, or bioluminescence (Kricka, L. in *Nonisotopic DNA Probe Techniques* (1992), Academic Press, San Diego, pp. 3-28). Another class of labels are hybridization-stabilizing moieties which serve to enhance, stabilize, or influence hybridization of duplexes, e.g. intercalators, minor-groove binders, and cross-linking functional groups (Blackburn, G. and Gait, M. Eds. "DNA and RNA structure" in *Nucleic Acids in Chemistry and Biology*, 2.sup.nd Edition, (1996) Oxford University Press, pp. 15-81). Yet another class of labels effect the separation or immobilization of a molecule by specific or non-specific capture, for example biotin, digoxigenin, and other haptens (Andrus, A. "Chemical methods for 5' non-isotopic labeling of PCR probes and primers" (1995) in *PCR 2: A Practical Approach*, Oxford University Press, Oxford, pp. 39-54).

The terms "annealing" and "hybridization" are used interchangeably and mean the base-pairing interaction of one nucleic acid with another nucleic acid that results in formation of a duplex or other higher-ordered structure. The primary interaction is base specific, i.e. A/T and G/C, by Watson/Crick and Hoogsteen-type hydrogen bonding.

The term “end-point analysis” refers to a method where data collection occurs only when a reaction is substantially complete.

The term “real-time analysis” refers to periodic monitoring during PCR. Certain systems such as the ABI 7700 Sequence Detection System (Applied Biosystems, Foster City, Calif.) conduct monitoring during each thermal cycle at a pre-determined or user-defined point. Real-time analysis of PCR with FRET probes measures fluorescent dye signal changes from cycle-to-cycle, preferably minus any internal control signals.

The term “quenching” refers to a decrease in fluorescence of a first moiety (reporter dye) caused by a second moiety (quencher) regardless of the mechanism.

A “primer,” as used herein, is an oligonucleotide that is complementary to a portion of target polynucleotide and, after hybridization to the target polynucleotide, may serve as a starting-point for an amplification reaction and the synthesis of an amplification product. Primers include, but are not limited to, spanning primers. A “primer pair” refers to two primers that can be used together for an amplification reaction. A “PCR primer” refers to a primer in a set of at least two primers that are capable of exponentially amplifying a target nucleic acid sequence in the polymerase chain reaction.

The term “probe” comprises a polynucleotide that comprises a specific portion designed to hybridize in a sequence-specific manner with a complementary region of a specific nucleic acid sequence, e.g., a target nucleic acid sequence. In certain embodiments, the specific portion of the probe may be specific for a particular sequence, or alternatively, may be degenerate, e.g., specific for a set of sequences. In certain embodiments, the probe is labeled. The probe can be an oligonucleotide that is complementary to at least a portion of an amplification product formed using two primers.

The terms “complement” and “complementary” as used herein, refer to the ability of two single stranded polynucleotides (for instance, a primer and a target polynucleotide) to base pair with each other, where an adenine on one strand of a polynucleotide will base pair to a thymine or uracil on a strand of a second polynucleotide and a cytosine on one strand of a polynucleotide will base pair to a guanine on a strand of a second polynucleotide. Two polynucleotides are complementary to each other when a nucleotide sequence in one polynucleotide can base pair with a nucleotide sequence in a second polynucleotide. For instance, 5'-ATGC and 5'-GCAT are complementary.

A “label” refers to a moiety attached (covalently or non-covalently), or capable of being attached, to an oligonucleotide, which provides or is capable of providing information about the oligonucleotide (e.g., descriptive or identifying information about the oligonucleotide) or another polynucleotide with which the labeled oligonucleotide interacts (e.g., hybridizes). Labels can be used to provide a detectable (and optionally quantifiable) signal. Labels can also be used to attach an oligonucleotide to a surface.

A “fluorophore” is a moiety that can emit light of a particular wavelength following absorbance of light of shorter wavelength. The wavelength of the light emitted by a particular fluorophore is characteristic of that fluorophore. Thus, a particular fluorophore can be detected by detecting light of an appropriate wavelength following excitation of the fluorophore with light of shorter wavelength.

The term “quencher” as used herein refers to a moiety that absorbs energy emitted from a fluorophore, or otherwise interferes with the ability of the fluorescent dye to emit light. A quencher can re-emit the energy absorbed from a fluorophore in a signal characteristic for that quencher, and thus a quencher can also act as a fluorophore (a fluorescent

quencher). This phenomenon is generally known as fluorescent resonance energy transfer (FRET). Alternatively, a quencher can dissipate the energy absorbed from a fluorophore as heat (a non-fluorescent quencher).

As used herein the term “sample” refers to a starting material suspected of harboring a particular microorganism or group of microorganisms. A “contaminated sample” refers to a sample harboring a pathogenic microbe (such as MAP) thereby comprising nucleic acid material from the pathogenic microbe. Examples of samples include, but are not limited to, veterinary samples (samples obtained from animals suspected of harboring or being infected by a microorganism—these may include any body fluid or tissue sample), food samples (including but not limited to samples from food intended for human or animal consumption such as processed foods, raw food material, produce (e.g., fruit and vegetables), legumes, meats (from livestock animals and/or game animals), fish, sea food, nuts, beverages, drinks, fermentation broths, and/or a selectively enriched food matrix comprising any of the above listed foods), water samples, environmental samples (e.g., soil samples, dirt samples, garbage samples, sewage samples, industrial effluent samples, air samples, or water samples from a variety of water bodies such as lakes, rivers, ponds etc., samples obtained from animal pens/barns/farms), air samples (from the environment or from a room or a building), forensic samples, agricultural samples, pharmaceutical samples, biopharmaceutical samples, samples from food processing and manufacturing surfaces, and/or biological samples. A “biological sample” refers to a sample obtained from eukaryotic or prokaryotic sources. Examples of eukaryotic sources include mammals, such as a human, a ruminant a cow, a bison, a sheep, a horse, a pig, a chicken, a turkey, a livestock animal, a fish, a crab, a crustacean, a rabbit, a game animal, a murine animal such as rat or mouse, etc. A biological sample may include tissue samples, cell samples, blood, serum, plasma, pus, cerebrospinal fluid, bone marrow, urine, feces, saliva, milk, mucus, or other materials from a human or a livestock animal. A biological sample can be, for instance, in the form of a single cell, plurality of cells, in the form of a tissue, or in the form of a fluid.

A sample may be tested directly, or may be prepared or processed in some manner prior to testing. For example, a sample may be processed to enrich any contaminating microbe and may be further processed to separate and/or lyse microbial cells contained therein. Lysed microbial cells from a sample may be additionally processed or prepares to separate, isolate and/or extract genetic material from the microbe for analysis to detect and/or identify the contaminating microbe. Analysis of a sample may include one or more molecular methods. For example, according to some exemplary embodiments of the present disclosure, a sample may be subject to nucleic acid amplification (for example by PCR) using appropriate oligonucleotide primers that are specific to one or more microbe nucleic acid sequences that the sample is suspected of being contaminated with. Amplification products may then be further subject to testing with specific probes (or reporter probes) to allow detection of microbial nucleic acid sequences that have been amplified from the sample. In some embodiments, if a microbial nucleic acid sequence is amplified from a sample, further analysis may be performed on the amplification product to further identify, quantify and analyze the detected microbe (determine parameters such as but not limited to the microbial strain, pathogenicity, quantity etc.).

As used herein “preparing” or “preparing a sample” or “processing” or processing a sample” refers to one or more of the following steps to achieve extraction and separation of a

nucleic acid from a sample: (1) bacterial enrichment, (2) separation of bacterial cells from the sample, (3) cell lysis, and (4) nucleic acid extraction and/or purification (e.g., DNA extraction, total DNA extraction, genomic DNA extraction, RNA extraction). Embodiments of the nucleic acid extracted include, but are not limited to, DNA, RNA, mRNA and miRNA.

As used herein, “presence” refers to the existence (and therefore to the detection) of a reaction, a product of a method or a process (including but not limited to, an amplification product resulting from an amplification reaction), or to the “presence” and “detection” of an organism such as a pathogenic organism or a particular strain or species of an organism.

As used herein, “detecting” or “detection” refers to the disclosure or revelation of the presence or absence in a sample of a target polynucleotide sequence or amplified target polynucleotide sequence product. The detecting can be by end point, real-time, enzymatic, and by resolving the amplification product on a gel and determining whether the expected amplification product is present, or other methods known to one of skill in the art.

The presence or absence of an amplified product can be determined or its amount measured. Detecting an amplified product can be conducted by standard methods well known in the art and used routinely. The detecting may occur, for instance, after multiple amplification cycles have been run (typically referred to an end-point analysis), or during each amplification cycle (typically referred to as real-time). Detecting an amplification product after multiple amplification cycles have been run is easily accomplished by, for instance, resolving the amplification product on a gel and determining whether the expected amplification product is present. In order to facilitate real-time detection or quantification of the amplification products, one or more of the primers and/or probes used in the amplification reaction can be labeled, and various formats are available for generating a detectable signal that indicates an amplification product is present. For example, a convenient label is typically a label that is fluorescent, which may be used in various formats including, but are not limited to, the use of donor fluorophore labels, acceptor fluorophore labels, fluorophores, quenchers, and combinations thereof. Assays using these various formats may include the use of one or more primers that are labeled (for instance, scorpions primers, amplifluor primers), one or more probes that are labeled (for instance, adjacent probes, TaqMan® probes, light-up probes, molecular beacons), or a combination thereof. The skilled person will understand that in addition to these known formats, new types of formats are routinely disclosed. The present invention is not limited by the type of method or the types of probes and/or primers used to detect an amplified product. Using appropriate labels (for example, different fluorophores) it is possible to combine (multiplex) the results of several different primer pairs (and, optionally, probes if they are present) in a single reaction. As an alternative to detection using a labeled primer and/or probe, an amplification product can be detected using a polynucleotide binding dye such as a fluorescent DNA binding dye. Examples include, for instance, SYBR® Green dye or SYBR® Gold dye (Molecular Probes). Upon interaction with the double-stranded amplification product, such polynucleotide binding dyes emit a fluorescence signal after excitation with light at a suitable wavelength. A polynucleotide binding dye such as a polynucleotide intercalating dye also can be used.

As used herein a “MAP organism” or “MAP microorganism” refers to an organism of the subspecies *Mycobacterium avium* subspecies *paratuberculosis* (MAP).

As used herein, a “non-MAP organism” is an organism of the *Mycobacterium avium* Complex (MAC) with the exception of *Mycobacterium avium* subsp. *paratuberculosis*. Non-MAP organisms include *Mycobacterium avium* subspecies *avium* (MAA), *Mycobacterium avium* subsp. *hominissuis* (MAH) and *Mycobacterium avium* subsp. *silvaticum* (MAS), as well as MAC organisms of indeterminate subspecies.

As used herein, a “MAP-specific polynucleotide” refers to a nucleic acid sequence that is able to specifically hybridize to a MAP specific target nucleic acid sequence and/or to a portion and/or complement thereof, under suitable hybridization conditions and which does not hybridize with other nucleic acid sequences that do not encode for a MAP-specific nucleic acid sequence, portions thereof or complements thereof. In some embodiments, a “MAP-specific polynucleotide” of the disclosure may be a probe or primer sequence specific to MAP. It is well within the ability of one skilled in the art, using the present teachings, to determine suitable hybridization conditions based on probe length, G+C content, and the degree of stringency required for a particular application.

It is expected that minor sequence variations in MAP-specific target nucleotide sequences associated with nucleotide additions, deletions and mutations, whether naturally occurring or introduced in vitro, would not interfere with the usefulness of primer and probe sequences disclosed herein in the detection of MAP organisms, as would be understood by one of skill in the art. Therefore, the scope of the present invention as claimed is intended to encompass minor variations in the sequences of the MAP specific target nucleic acid sequences described here and the MAP specific nucleotides, such as, exemplary primer sets and probe sequences set forth here, and all sequences disclosed also comprise sequences having at least a 90% sequence homology to these sequences.

A probe may be RNA or DNA. Depending on the detection means employed, the probe may be unlabeled, radiolabeled, chemiluminescent labeled, enzyme labeled, or labeled with a dye. The probe may be hybridized with a sample in solution or immobilized on a solid support such as nitrocellulose, a microarray or a nylon membrane, or the probe may be immobilized on a solid support, such as a silicon chip or a microarray.

Conditions that “allow” an event to occur or conditions that are “suitable” for an event to occur, such as hybridization, strand extension, and the like, or “suitable” conditions are conditions that do not prevent such events from occurring. Thus, these conditions permit, enhance, facilitate, and/or are conducive to the event. Such conditions, known in the art and described herein, may depend upon, for example, the nature of the nucleotide sequence, temperature, and buffer conditions. These conditions may also depend on what event is desired, such as hybridization, cleavage, or strand extension. An “isolated” polynucleotide refers to a polynucleotide that has been removed from its natural environment. A “purified” polynucleotide is one that is at least about 60% free, preferably at least about 75% free, and most preferably at least about 90% free from other components with which they are naturally associated.

The words “preferred” and “preferably” refer to embodiments of the invention that may afford certain benefits, under certain circumstances. However, other embodiments may also be preferred, under the same or other circumstances. Furthermore, the recitation of one or more preferred embodi-

ments does not imply that other embodiments are not useful, and is not intended to exclude other embodiments from the scope of the invention.

The terms “comprises” and variations thereof do not have a limiting meaning where these terms appear in the description and claims. Unless otherwise specified, “a,” “an,” “the,” and “at least one” are used interchangeably and mean one or more than one.

Also herein, the recitations of numerical ranges by end-points include all numbers subsumed within that range (e.g., 1 to 5 includes 1, 1.5, 2, 2.75, 3, 3.80, 4, 5, etc.). The term “and/or” means one or all of the listed elements or a combination of any two or more of the listed elements.

There are many known methods of amplifying nucleic acid sequences including e.g., PCR. See, e.g., PCR Technology: Principles and Applications for DNA Amplification (ed. H. A. Erlich, Freeman Press, NY, N.Y., 1992); PCR Protocols: A Guide to Methods and Applications (eds. Innis, et al., Academic Press, San Diego, Calif., 1990); Mattila et al., Nucleic Acids Res. 19, 4967 (1991); Eckert et al., PCR Methods and Applications 1, 17 (1991); PCR (eds. McPherson et al., IRL Press, Oxford); and U.S. Pat. Nos. 4,683,202, 4,683,195, 4,800,159 4,965,188 and 5,333,675 each of which is incorporated herein by reference in their entireties for all purposes.

Nucleic acid amplification techniques are traditionally classified according to the temperature requirements of the amplification process. Isothermal amplifications are conducted at a constant temperature, in contrast to amplifications that require cycling between high and low temperatures. Examples of isothermal amplification techniques are: Strand Displacement Amplification (SDA; Walker et al., 1992, Proc. Natl. Acad. Sci. USA 89:392 396; Walker et al., 1992, Nuc. Acids. Res. 20:1691 1696; and EP 0 497 272, all of which are incorporated herein by reference), self-sustained sequence replication (3SR; Guatelli et al., 1990, Proc. Natl. Acad. Sci. USA 87:1874 1878), the Q.beta. replicase system (Lizardi et al., 1988, BioTechnology 6:1197 1202), and the techniques disclosed in WO 90/10064 and WO 91/03573.

Examples of techniques that require temperature cycling are: polymerase chain reaction (PCR; Saiki et al., 1985, Science 230:1350 1354), ligase chain reaction (LCR; Wu et al., 1989, Genomics 4:560 569; Barringer et al., 1990, Gene 89:117 122; Barany, 1991, Proc. Natl. Acad. Sci. USA 88:189 193), transcription-based amplification (Kwoh et al., 1989, Proc. Natl. Acad. Sci. USA 86:1173 1177) and restriction amplification (U.S. Pat. No. 5,102,784).

Other exemplary techniques include Nucleic Acid Sequence-Based Amplification (“NASBA”; see U.S. Pat. No. 5,130,238), Q β replicase system (see Lizardi et al., BioTechnology 6:1197 (1988)), and Rolling Circle Amplification (see Lizardi et al., Nat Genet 19:225 232 (1998)). The amplification primers of the present invention may be used to carry out, for example, but not limited to, PCR, SDA or tSDA. Any of the amplification techniques and methods disclosed herein can be used to practice the claimed invention as would be understood by one of ordinary skill in the art.

PCR is an extremely powerful technique for amplifying specific polynucleotide sequences, including genomic DNA, single-stranded cDNA, and mRNA among others. Various methods of conducting PCR amplification and primer design and construction for PCR amplification will be known to those of skill in the art. Generally, in PCR a double-stranded DNA to be amplified is denatured by heating the sample. New DNA synthesis is then primed by hybridizing primers to the target sequence in the presence of DNA polymerase and excess dNTPs. In subsequent cycles, the primers hybridize to

the newly synthesized DNA to produce discreet products with the primer sequences at either end. The products accumulate exponentially with each successive round of amplification.

The DNA polymerase used in PCR is often a thermostable polymerase. This allows the enzyme to continue functioning after repeated cycles of heating necessary to denature the double-stranded DNA. Polymerases that are useful for PCR include, for example, Taq DNA polymerase, Tth DNA polymerase, Tfl DNA polymerase, Tma DNA polymerase, Tli DNA polymerase, and Pfu DNA polymerase. There are many commercially available modified forms of these enzymes including: AmpliTaq® and AmpliTaq Gold® both available from Applied Biosystems. Many are available with or without a 3- to 5' proofreading exonuclease activity. See, for example, Vent® and Vent®. (exo-) available from New England Biolabs.

Other suitable amplification methods include the ligase chain reaction (LCR) (e.g., Wu and Wallace, Genomics 4, 560 (1989) and Landegren et al., Science 241, 1077 (1988)), transcription amplification (Kwoh et al., Proc. Natl. Acad. Sci. USA 86, 1173 (1989)), and self-sustained sequence replication (Guatelli et al., Proc. Nat. Acad. Sci. USA, 87, 1874 (1990)) and nucleic acid based sequence amplification (NABSA). (See, U.S. Pat. Nos. 5,409,818, 5,554,517, and 6,063,603). The latter two amplification methods include isothermal reactions based on isothermal transcription, which produce both single-stranded RNA (ssRNA) and double-stranded DNA (dsDNA) as the amplification products in a ratio of about 30 or 100 to 1, respectively.

Those having ordinary skill in the art, in light of this specification, will understand that many modifications, alternatives, and equivalents of the embodiments described above are possible. All such modifications, alternatives, and equivalents are intended to be encompassed herein.

EXAMPLES

The following procedures are representative examples of embodiments according to the disclosure that may be employed for the detection of a MAP organism. These examples are not intended to be limiting to the scope of the claims and/or the disclosure in any way.

Example 1

Compositions & Methods to Detect MAP Organisms

The present example describes exemplary assays designed to detect MAP organism using probe and primer sequences designed as described herein. The described assays can be used in diagnostic methods and kits to detect a MAP organism in a sample, such as for example a sample obtained from an animal suspected of being infected with a MAP organism, with no cross-reactivity to a non-MAP or other MAC organism.

A set of real-time qPCR primers and probes that can specifically amplify and detect the presence of *Mycobacterium avium* subsp. *paratuberculosis* without cross-reactivity with other subspecies in the *Mycobacterium avium* complex including *M. avium hominissuis*, and *M. avium avium* are described in Table 3 below.

Table 3 lists a set of TaqMan® primer/probe sets, where primer sequences comprise sets/pairs of primers, each primer set including a forward and a reverse primer, and a corresponding probe sequence, that have been shown to be specific to the amplification of MAP with a CT cutoff < 38.

TABLE 3

Assay	Coordinate on NC_002944.2						
	ID No	Forward	Reverse	Probe	Left	Right	Locus
61801	CGGTGCGCGGTA GGA SEQ ID NO: 6	CGACCTGGTCTGCG TCATC SEQ ID NO: 7	CCGCGAGGACCTG CTGACGC SEQ ID NO: 8	239559	239635	MAP0225c	Hypothetical protein
61802	GCGATCCGGCGA AACG SEQ ID NO: 9	GACGAGTTAGACGC ACAGATTCT SEQ ID NO: 10	ACGCCCTGCAGTT AGTCCCCG SEQ ID NO: 11	239855	239922	MAP0225c	Hypothetical protein
61803	ATCGACCGGCTG GTGTAC SEQ ID NO: 12	GCGAGCAGCACGT TGAAC SEQ ID NO: 13	TCGACGCGATGAC CGCGCTCG SEQ ID NO: 14	240155	240255	MAP0226	Hypothetical protein
61805	CAGACACCTGCA GAGCTACTAC SEQ ID NO: 15	AGCATGTCGGCGA CGTT SEQ ID NO: 16	CAAGCCCCACTAC GCCGCCCT SEQ ID NO: 17	240712	240777	MAP0226	Hypothetical protein
61808	TCGCGGTGCCG ATCAG SEQ ID NO: 18	GTTTGGATGCCCG TGACATTC SEQ ID NO: 19	CCGCGCTCCCCGA AATGCTC SEQ ID NO: 20	3088984	3089043	MAP2756c	Large hypothetical protein
61813	CGGCGAACCCG TGACA SEQ ID NO: 21	CACAGGTGCTCCC ATTGAGT SEQ ID NO: 22	CCGCGCCAGCGT GTAGGC SEQ ID NO: 23	4168688	4168782	MAP3740	Large hypothetical protein
61815	GCGCAGCTTAC CATCGA SEQ ID NO: 24	GTGCGAAGGCTTC GAAGAC SEQ ID NO: 25	AACAGGCTGCCGC CACGCT SEQ ID NO: 26	4169324	4169424	MAP3740	Large hypothetical protein
61816	GCCACGACGGTG ACATCA SEQ ID NO: 27	CCAGGTTGCTGGT GAACA SEQ ID NO: 28	CCGCGTTGCGGCC AGCCCC SEQ ID NO: 29	4169626	4169706	MAP3740	Large hypothetical protein
61817	GGAATCGATCTC GGTGGCTTTT SEQ ID NO: 30	CCACCGCCGGAGC AA SEQ ID NO: 31	CCGCGTGGCGCCG CA SEQ ID NO: 32	4169849	4169946	MAP3740	Large hypothetical protein
61819	GCCGCGACGGTA GGT SEQ ID NO: 33	CCAGATCTCGGCAA GCGT SEQ ID NO: 34	ACGACGCCGGCAC AGCC SEQ ID NO: 35	4170784	4170872	MAP3740	Large hypothetical protein
61822	CGGACGGAGAGG TGGAC SEQ ID NO: 36	GGGCTCCGGCACCA SEQ ID NO: 37	TTGCCGACCCGCA GACCTG SEQ ID NO: 38	4174593	4174666	MAP3742	Large hypothetical protein
61823	GGTAACCAGCAAC CCGATCTG SEQ ID NO: 39	CCCAGGTGAACCA CTTTGA SEQ ID NO: 40	TCGGTTGATGATTT CACCCAGGACGC SEQ ID NO: 41	4194693	4194779	MAP3756c	Hypothetical protein
61825	GGCTCCTCCGAA CCA SEQ ID NO: 42	GCAACATTGGCGAT CGACATC SEQ ID NO: 43	CAGCGGAGCATCAC CGACCCCGG SEQ ID NO: 44	4195499	4195561	MAP3757c	Hypothetical protein
61826	GGCTCGGGCCTGA TAGC SEQ ID NO: 45	GCGGCCTGAACTCC TATGATC SEQ ID NO: 46	CCGCGTTCAGTGCC GTGGCC SEQ ID NO: 47	4195904	4195980	MAP3757c	Hypothetical protein
61829	GCAATACTCGTGA ACCGTAAACTG SEQ ID NO: 48	GTGGCGGGTGTT G SEQ ID NO: 49	CCCTGGCGGGCATT TCCTCTACG SEQ ID NO: 50	4196988	4197054	MAP3758c- MAP3759c	Intergenic

*Assay ID describes an assay comprising using Forward Primers and Reverse Primers described in one row and in some embodiments Probe described in the same row.

An exemplary method of detecting the presence of a MAP organism in a sample comprises: 1) isolating nucleic acid from a sample suspected to contain MAP and 2) detecting the presence of at least one MAP-specific target nucleic acid and/or fragment thereof and/or complement thereof as shown using an assay described in Table 3 above comprising using for example a set of primers, comprising a forward and a reverse primer described in a row to amplify the at least one MAP-specific target nucleic acid and/or fragment and/or complement thereof; and detecting the amplified product. In

some optional embodiments, a probe described in the same row can be used to detect an amplified product amplified by forward and reverse primers from the same row. Such a method can be a diagnostic method, where a sample is derived from a mammalian animal suspected of being infected with a MAP organism. A diagnostic method can be used for example to diagnose a MAP related/caused disease or condition, such as but not limited to Johne's disease and/or Crohn's disease, by detecting the presence of a MAP organism in a sample derived from a mammal suspected of having this disease.

As shown in Table 3, an Assay ID number (such as 61801, 61802 etc.) is assigned to describe an associated primer pair and associated probes that can be used for amplification and/or detection of a target MAP sequence (such as in non-limiting examples SEQ ID. NOs: 1-5, Table 1) and/or fragments and/or complements thereof. These specific combinations of primer pairs and probe sequences have been designed to selectively amplify MAP specific target nucleic acid sequences. In some embodiments these primer pairs are degenerate.

In some embodiments, multiplex assays may be performed by simultaneously contacting a sample with the one or more primer pairs. Multiplex assays of the disclosure can be performed in parallel or sequentially.

Each of the assays was tested against the 16 sequenced *Mycobacterium avium* strains to confirm diagnostic sensitivity and specificity. For every assay the forward primer, reverse primer and probe were diluted in a mix to 900 nM for each primer and 250 nM for the probe (final, 1x, concentration). 84, of sample DNA (18-39 ng) was input into a 25 µL qPCR using the Path-ID™ qPCR master mix and each candidate MAP assay. The thermal profile and qPCR set-up is shown below in Table 4.

TABLE 4

MAP AgPath-ID Master Mix		
25 ul	reaction volume	
1X	Component	100 rxns
5 ul	Sample	500 ul
8.50 ul	NF H2O	650 ul
12.50 ul	2X Path-ID PCR Master Mix	1250 ul
1.00 ul	25X MAP Primer Probe Mix	100 ul
Thermal Profile		
1. 95dC: 10 mins: 1Rep		
2. 95dC: 15 sec		
80dC: 80 sec		
40 Repts		

Table 5, below, has results of sensitivity and specificity testing. There was no cross-reactivity with any other non-MAP species (with a CT cutoff<38 considered positive).

The above experiment demonstrates the feasibility of using any of the assays listed below as a specific and highly sensitive detection method for *Mycobacterium avium* subsp. *paratuberculosis*. In some embodiments, the disclosure provides a set of highly-specific TaqMan® assays for the detection of *Mycobacterium avium* subsp. *paratuberculosis*.

One or more advantages of the methods of the present disclosure are described in the following section. Each individual assay can independently serve as a highly-accurate diagnostic assay for MAP detection. Each assay has the advantage of targeting single-copy gene in the MAP genome instead of genomic insertion sequences that are utilized in other MAP detection assays. Insertion sequences have the capacity to mobilize between genomes and insert into new species, making them suboptimal for a species-specific assay.

The currently described assays and methods are able to detect MAP in multiple host species including sheep, cattle, and bison quickly and accurately with no cross-reactivity between very closely related *M. avium* subspecies making it ideal for veterinary diagnostics.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be appreciated by one skilled in the art from reading this disclosure that various changes in form and detail can be made without departing from the spirit and scope of the invention. These methods are not limited to any particular type of sample or nucleic acid contained therein for e.g., total genome DNA, RNA, cDNA and the like may be analyzed using some or all of the methods disclosed in this disclosure. This disclosure provides powerful tools for analysis of complex nucleic acid samples. From experiment design to detection of MAP microbes, the above disclosure provides for fast, efficient and inexpensive methods for detection of pathogenic organisms.

All publications and patent applications cited above are incorporated by reference in their entirety for all purposes to the same extent as if each individual publication or patent application were specifically and individually indicated to be so incorporated by reference. Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

TABLE 5

Sample Info Assay ID No.	Wood pigeon M.	Asian Elephant MAA	Red Tailed Hawk MAA	Ruddy Duck MAA	Avian broadbill M. avium	White lipped deer MAH	Pig MAH	Pig Intermediate sp. MAC	Dog Intermediate sp. MAC	deer Intermediate sp. MAC	bison MAP	bison MAP	cattle MAP	cattle MAP	sheep MAP	sheep MAP
	silvaticum	10- ATCC	10- MAA	11- MAA	05- M. avium	10- MAH	10- MAH	10- MAC	09- MAC	11- MAC	10- MAP	10- MAP	10- MAP	10- MAP	08- MAP	11- MAP
61801	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	27.12	28.50	28.47	29.31	30.57	27.03
61802	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	20.10	21.17	20.30	21.15	22.04	20.68
61803	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	26.01	27.56	26.35	27.30	28.06	25.11
61805	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	19.73	20.12	19.60	20.69	21.49	20.09
61808	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	18.33	19.69	19.05	19.57	20.04	19.53
61813	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	29.65	30.04	29.86	30.88	31.10	29.91
61815	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	19.78	21.45	21.36	22.16	22.14	21.03
61816	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	21.03	22.30	21.21	23.19	22.74	21.11
61817	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	32.05	33.05	31.55	33.60	34.49	30.34
61819	40.00	40.00	40.00	40.00	40.00	40.00	39.76	40.00	40.00	40.00	26.62	25.74	28.12	27.34	28.55	27.02
61822	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	19.59	20.26	20.00	21.08	20.75	20.04
61823	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	19.31	20.06	19.83	20.33	20.27	20.28
61825	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	18.21	19.23	19.26	19.63	19.57	19.20
61826	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	20.73	21.02	22.59	23.12	22.45	21.58
61829	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	40.00	19.12	19.91	19.78	20.25	20.22	19.56

TABLE 6

SEQ ID NOs: 1-5

SEQ ID NO: 1:

CGGGTGC GGATTGATCAGCCGCCCATGGGCCACCAGCGATCCGTGTTGCTTGAGCCGTGGG
TGCGGACGTTGACCTCGTAGCCAGACACCCCGTCGAGCTGGCCGAGTTGCTCGCCAGGTAG
CGGTACAGGTCGTCGGAATCTCGGCAGATGGCCACGGCCATCAGGTTTTTGGATCCGCTTGT
GGCGCCGGCGAAAGCAATCTGCGGATGGGCCGCGATACGTTCCGCCGACCGTGTGCACGTCG
CGAGGCGCAACCGTGAAGCACAGCATGGCGCTGAGGCGATGGCCGAGCCGTTCCGGTAGTA
CGTGCAGCTCGTAGACCAGGGCGCCGAGGCTCCAGCGCCGCGATCCGCTGCTCACCCGC
GCCACCGACCAACCGGTGCGCGCGCCAGTTGGGACTGCGGAGTGGCCGCTCGTCGGCGA
GGCCGCGAGCAGCGGGGGTCTTCCGGCGTGGGCCGGGACCGTGAAGTTCCCGACTCCGG
CTGGGCCCGCGCGCCACGATCCGCTGCGCTTGC CGGTCCGAGAGCGCGTCCGCTAGCCCG
TCCAGGGCGCGTTCACCGGGTCCCGAACGAATGACAGCATCAGATCGATGCTGATGTCGGTG
ACCGCGGGCGTGC GCGTAGGAGCTGCGTCAAGCAGTCTCGCGGGTGTCCGCCAGCGCGC
CACGGATGACGCGAGCCAGGTCCGCTCCACCCCGCAGCACATTGGCGTGGGACACGTCGGG
GCGTGCACCCAGCGCGTCCGCCAGCCGCGATATGCGGTCCGGCGCCGACCGATCGGCAC
ACCCAGTCCGCGTCCGATCCGGCCACCGGTTCTCCAGCCGACGACGCGCACCCCGTCC
GCGCGCAGCCGTTGATACCGCCGGGCGACCGTCTGCTCGGTGGCGCCACCCAGCTGGCG
ATCCGGCGAAACGAGACCCGGGAGCTAACTGACGGGCGTGCAGAATCTGTGCTTAACT
CGTCCGTCATCAGGAAAAGTATCCGCGTGGACAGGTCCAAGTGAGGATTTCCAGGCCAATTTG
GTCGCGCGGAGCGCAATTTCAACGCCTCCGCTGTGAGCTGGAAGCTCGGCCTAAGGAGCGC
TCACATGCACATCGAAATCGGACTGCCAGCCATATCCGCCACGTGGGGGGCGGCTGAC
GTGAGTGGGCGCGCGGGCCGAAACCGCGGCTTTCGCGGCTGGCCGCGATCGACCGGC
TGGTGTACCCGAGCCTCGACGCGATGACCGCGCTCGCGGTGGCCGCGGGCGGACCCCGG
CATCGGGTGAAGTTCAACGTCGTCGCTCGCTCCGCTGTACCCCGCGGTGCTGCTGGCCAAAG
AGGTGATACCCCTCGCGGAGGCGTCCGGCGCGCGCTGCGCTGGGGTGGGCGTCCGGGAG
CCGCTGCGACGACTACCCGCGTCCGCGTGCAGTATCGCGCGCGCGGGCGCATTTCTCGACG
AGACGGCGGGCGTGTGCGTGCCTGCGTGTGAGGCGAGGTTGTTGACCCGCGACAGCCGCT
GTGCCCGCGCGCGTGC GCGATCCCGATCCTGTTCGGTGGACGGGCGGACGCGATCCGC
CGCGCCGTGACCGTCCGTCGAGCGCTGGACCCCGGGCGCTGCGCGATTACGCCAACCGAT
CGGTGTTCCGCGAGAGGGTCCGGCGCGCATGGGCGCGCGGGCGCTCCGAGACGGCCGTG
GCTGCAGGCCAGCGTCAACTTCCGCTTCCGGCGATGACGAGGCGCTCGCGCTGGCCGAGAC
ACCTGCAGAGCTACTACGGCTTCAAGCCGACTACGCGCCCTCAACGTGCGCCGACATGC
TCACCACGCCGAGGAGCGGC

SEQ ID NO: 2:

CCCGCACCGCCGCTCTGCCGCGGACAATGCCGCTTGGCTTCGCGGCTGCCGCGTGAC
GGCGCTTCGCGCGGAGTTCGGCGTCCGCGCGCGCCAGCACCTCGGTTCCACTGCCATGG
CCGCGGGGCCCATGTTGCGCATCGCCGCTGCTGGCTGTGGTGGCTGGGCTTCGACGCTGAAC
AGCTTGTCTTCAAACATGCCAGACCCGTCGCCGATCGGTGACAGGATCGTCGAGGCGATGTT
GTAGCCCTTGATCGCAACCATGCGCGTCCCATTTGCACCCACACGCGGTCCCGCATCCA
GCAACCGCTTGTGCTCGGTGAGCCAATGCGCCGACTCCCCAAGCACCCCAACGACTTCAGTC
ATCGCTGGCAGAAAGTCTCGGCGGATTCGATGCGGGTGCCTCAACGCCCTTTGAAGTC
CTTCACTTGGCGTGTAGGTTCTCTGGGATTCGGTGAATCCCTTGGTATTGGCCGACCCGTC
GGCTTGCCTGACTGACCTCTTGACCGCGGACGCGGCTTTGGGCCGTTCTCCCGACCA
ACTGCAATACCGTCCGAGATGTTCTTGGCCGCGACCATGAGGGCCAGCGCTTGGATCATC
GACTGGACATCACCCTTCCCGACTTGAGCTGATGTTGAAGCTGTCAATGCTTTTTCTTGG
TTCCACCCTGCTCGACAAGGTTGGCTTCTCAACGCTCAAGCCGCGCGGTTCTTGGCGAA
CTGCTTGTACGCTCAGCATTGTTCTGATCGCCGTCCGCGCCGCTGCGCTGCGCGGGCA
GCTTCGAGAACATCTCATTAGCGCTGTCCGCCACCTGCTGCTCTTGAACAGGCGTCCAGC
GTGACCATCCCGCAGCGGCCAAGCTTCGACTGGAATCGTGTATACAAACAGGTTCCGCGTGC
GATCAGGCGCGCTCCCGAAATGCTCTGAAATGTCACGGGCATCCAAACGAGCTGTCCCA
TCTCTGGGACATTTGCTGGGTTGGCTTGGCCAGCGAGTGTGCGTGCATATTTCTGG
GTGGCTGATCAGCACCATACCCGACTGAGTGAGCATCGCCAACGACGCATACAGGTCATT
GACGCTTTCGCAACTCCGCGCGCGACCGGCTCGACGTTGTGCAACGCCCGCGAAGTCTT
GCAGCGAAACCTTCGACAAACCCACGGCCACATTCATTTGAGGTGACGTTGGCGGCTGCT
TCGACCCGGATGTGGTAGTCGTGATCGTGGTCTGACCGCGCTGATCGCTCGCCAGGTC
CGCGCCCTTTCGAGCGCAAGCTGCGCTGAAGCCGCTCAACACCTTGTGCGATCACTGCCCC
GATAGCCGGCTTCTCGACACCCAGCGCAGCGTCCATCAGCTTTTGGGCGCATAAACGACC
ACGCTCGACAGTTTCAGCACGCCATCGCTGATCGCTTCCAGATTGGCCGCGACTCTCCGGC
CACCGTGTGACGCTTGCACAGACTGCTGGAAATCCCGCCGATCAGTCCGCGAGACA

SEQ ID NO: 3

ACATCGGGTGGAGTGGCGGAGATCGAGCACGCGCTGCGCGGCCACCCGCTGGTCCGCGCC
GCGACGGTGGTCCCATCCACAACCTGCACTGCGCTGGGCGCGGGATCGTCGTAACCGGCA
GCGGCGCAGAGCAATTTGACGACTCCACACCCGGCGCGCTGCGCGCCATCTCGCCGTCGGA
CTCCCGCAGTACATGATCCCAAGGTTTCGTTCTCATGCCCGAGCTACCGCTCACCGCCAA
CGGCAAGGTCGACCGGGGCAAAATCGCGCGCGCTCGAAGCGCGCGCGGGCACCCAG
CCGCTCGACACGTCATCCACCTCACTGTGGTTCGAGCGGCTGGTCCGCGAGGTCGTTCCGA
TGTGCTGGGCGCGCGATCACCGCGCGGAGGACAAATTTCTCGCCAGGGCGGCGATAGCC
TGCGCGCCACGCAAGCGTCCGCCGACTGACGCGCAGGGGAGTGGCCGAGCGGAGGTTGG
CCAGTGTCTAGCCACCAGCGCTTGGGCGTTCAGCGCGCGTGTGCTCGCCGACCCGG
CATCCGAGGATCCGATCGCGCGCGCGATGTCGGCGAAACCGTGACACCGGGCGAGGGTT
CCCGCTCACCGGTTGACGAGGCTTACACGCTGGGCGCGCGGACTCAATGGGAGCACCC
TGTGCACCAACGATTTTCGCGGTGCTGTCGGCCGCGCGCCGAGTCTGCCGTTATAGACCT
GGATCGGTTTCCCGGTGGTCCACAGATCGTTCGACGAATTCGCGATGCTCGGTGCGCGC
TGGACCGCGACACCCCAACGGTTCAGGTCGACGCGCGGGCGGTTGCCGCTCATGACCT
TGATATACAGAGCACCCGACTGTTACTGCGCGCATGGCGGCGCCCGCTTCGATCCGC
ATTCCGTTCCGGTGTGATCCAGTGTTCGACCGCTGAGGTCACCCGCTCACGTCGCTGCTG
ATCAGCTATCTGGGCTCGATGCCCGCAGCTGTCCACCGTGTCCACGATCATCGCCGA

TABLE 6-continued

SEQ ID NOs: 1-5

ATACAGTCGCAACCCCGGCCGCGGAGGTGACCCGACCGCGCGGTCTTCGCCCGGTTTG
 CCTCCGAAAGCGTTGGGGCGAAAAAGATGTCGCAACAGCGTTGCCGGCCCTCCGCTGCTG
 CCGCTGCACGACCAGCGAGTACCCCTTCGAGCGGGTACCTTCGCGCGGCGCAGCTTCAC
 CATCGAAGAACAGGCTGCCGCCACGCTGCGTGAGCACGCCGCACACCTCGGGCTCACCCCC
 ACCGCGCTGGTCTTCAAGCCCTTCGCACATGCGCTGGCGTGCATCGGCGCCGGTCAGCGATT
 CGCGGTGACAGTCCCAAGTCGTACCGTCCCGACTACGCCCCCGCGGACCGCGAGGTGTTGG
 GCAACTTCACCCGCTGGCGCTGTGCGAGGTGCACTACGGCGCGGTGAGACCGGGATGTCGC
 GAAGCGGTGCCGCGGCGCAGCGGAACTGTGGCGCGCGGTGAGCACGACGGTGACATCA
 CCGGCGGGCTGGCCCAACCGCGGACCGGGTGGCTACCCCGTGGTGTACCCAGCACCCCT
 GGGGCTCACCCATCAGGACGCCAGCGGGCTGACCAACGTGCGGACATTGACCCAGACCCCG
 GGCGTTCGGCTGGACTGCCAGACCGAGGACGAGGTCCGCGAATTCGTATGAGCTGGGACA
 TAGCTACCAATGTGGTTCCCGCGAATCGATCTCGTGGCTTTTTCCCGATTCGAGGAGGCG
 GTGCGGCCACCGCGGGCAAGCCGAGCCCGCGGACCGGCCGTTGCTCCGGCGGTGGGGC

SEQ ID NO: 4

AGTGGGCGAGCGCGGTGATCGCCGCGCGCTGCGCCACTGCCGACCCGAGCAGGTGCTGCC
 GCAATACACCATGCTGGTGGCGCGCTGGGAGGCACTGCGATACGTCCTCCCTCAGGATACGCC
 GCTTCCGACGTCGAACGGCGGCTCGCCGGTTGGCCGCATCGTACCGGAGCCGCTGTACCC
 CCAGACGCTGATCGGAGACCCGAGCTGACGCCCGAGGCGCTGTGCTGCGTGACGACCCG
 ATGCGGATGGCCCTCGACGACCTGGCCGGGGCCTTTCGGGACCGCGCGCACGCTGGGGC
 GGCGCTTCGCGTCTGGAAGTCCGCTCCCGGACCGGCTGATCACCGAAGGTTGACCGA
 GCTGGTGGGCTGGTGGTTCGAGGAATACTTGTGCTGGAGCCAAACCGACGCTCGCCGGA
 ATCGCCGCGGGGCGGCGCTTCCCGGCCCGACCCGCGCACGTCGACGACCCGACGCGCGCTC
 GGGCTCGACGTGGTGTGCTGTGGTTCGCTGATCAGCTGCCCGACCGGAGGCGGGTCC
 TCGAGGCGATCACCGTGTCCGACGACGGTGGCTGTGGATGGTGGAGAATTCGAGGGCCAC
 CCAAGCGACGCTCATCAGCGCGGCGCTTCCGACCCCGCGCTGCTCGCGTCCGATTCAAGA
 CGCTGCGTCCGCGCATCGGTGGTGGCGGCTCATCGCCGACCGGTTGGCGACCGACGCGAC
 ATGATCCAGGACCGACCGGCTCAGCTCATCGCGACCCCGCCGCAAGCCCGGATGC
 CGACACCGCCGGCCGAAACAGCGCCGCGACCGTAGGTGGTCCGCGACCGGCTGTGCCGGCTC
 GTCGCTGCCGACCGACCGCCAGGTTGGTGGCCAGCTTCCCGAGATCTGGCAGCGTCTATCTCG
 CCATTCACAACCCCGGCTCGACGACGACTTCTTCTGCTCGGCGGTGACAGCCCTCGTCCGG
 ACCCGGTTCTACGCGCACTTCGGGCGCCGGTTTCGGCCAACTCGCTTCGTGACACTGTTT
 AACCACTCGACGCTCGGTGAGCTCGCGGACACCGCCGCGCGCACCGGCCGCGAAGTGT
 CGGTGGCGGCTGAGTCGACCCGGGGCGGCACCCAGACCCGAAACCGATTCCCGCTCACCGT
 CGTGAGAACCGCTATCGGGCCGGCGAGAAGGCGCGTTGATCTCGCGCGCGTCCGCGCG
 CACTGCTACTTCGAGTTCGAGCTCGCGGACTTCGACCGCGGAGATTGATTCGCGCGCAGC
 CCAACTCGTAGCACGCCACCGCGGACTGCGCACCCGCGTGCACCGCGGGCACCCGACGCG
 GCCTCCTCGGTTGAGGTGCGCGTCTGTCACACCGCGCGGATCGAGCCCGCTCGTGCGAGACCA
 CGACGACGTGCGAGCCGCGATGCGCGACCGATCATCGACTTGACGGCCCGCCCGGGCATC
 GACTTCGGGGTGCAAAACCGCGGACCGGGCGCACCGTTCGTCGCGCATCAGCATGGACAACA
 CCATGCTCGACGGCCGACGATGATGATCGCCCTGTCCGAACTCGATCACCTCTATCGCGCG
 GAAACCGTTGACCAATTCGCGCCGCTGGAACGCTTTCGCGCACTACGTTGGAAACCCACCC
 GGAGCTGCTGCCGACCGCGAGGCGGTTGCTCCCGCGGCTGGCCCGCAGCCGAGACTAT
 TGGCGCGACCGCTCCATCTTTCGCGCCGCGCGAAATGGCCGACATGCTACTGCTGTT
 CGAGATCGAGGAGCGAGGTTCAACCGGCAACCGCGACCATTCGCGCGTTCGACTGGTTCG
 CAGGTAACCGGATCGTCCGCTGCGAGGGCGTACCCGTCGCTCATTTCTGCTCGCCAACTA
 TGCACGGGTGCTGTCTCGGTGGTTCGGGACCGGACCACTTCGATCAACGTCACGCTATTTCG
 ACCGCGACCCGATGCTCGTGGGATCGAAAACGTCGTCGGAGATTCACTTCCCTGGTGTG
 TTGGAGTCCGAGTCATGAGCCCGCTCGATCTGGGAGCGTTCGCGCTCTCGACGCGC
 AATTGATGACCGACTGCTCCGACCGCGCGCGGACCGCGGTGGTTCGCAACCGGAACTGCT
 GCGGTTTCACGGCAACCCGACGGCCGCGCTGTTCCCGTCTGTTCAACGCGGACTGGGCC
 TTGTGACGCTCGGCTCGGCGCGGTCGCGTTTCGCGAACCGGTTATTCGCGCCCTCGCAG
 ACACCGCAGACGCTTGGACTTCCAGGTGTTGGGAAAGCGCGGGGCGCTGAAGCTGTCTG
 GGGACTTCGTAGTCAGCGGTGTCGCGCGCCACCGCGCACTCAGCTCGAGTCTGTTGGT
 GACGGCATCACCGTGTGCGCACGCGAGCCCGCATCGAACACAAGTTGGGCGAGGGGG
 CATCCAATGACGAGCTCCTGCAACGTTTCGAGGATCTGCGCGTCCGCGCTGGGTGAGCGG
 AGGGTCAACCTCACGACAATTTCTTCAGCTCGGCGGATTCGGTTCAGCGGACCAAGGT
 GGTCAACAGATCGGCCGTGAGCTGTCAGCTCGGCCACCCCTCGACTGCTGTTCCGCAATC
 CGGTGATCGGCGACTTCGCGCCAAAAATCGCGCACTGACAAACCGCGACGAAACCGGACT
 GACCGTTGAGGAGGCAATGTTATGACCGCGCGGAGCTCGTGCACACCTCGGGGTATCG
 GCGTCCAATGTTGGGCCGCGGTGAGAATTTGCGCTACCGAGCACCGCAACAAGTCTCACC
 CGGACTGAAAGCTCAACTGCGCGCGCTCAAAAAGGACGTTGATTAACCTGCTGGCGGAA
 AGACGACCTGCTGCGCGCGCGCAGGACCGGTTGAGCCGTTTCGCTCACCGACGTCGAA
 GCCGATATCTGGTGGGCGCACGTCGCGCTTCAGTGGGGCGGGTGGCTGCCACGGCTA
 CGCCGAGTTCGCGGTCGACACACCGTGGCAACAACGAGCGCGGAGCAATATCGGGAGGGC
 TGGCGCAAGGTTGCGACCGCCACGACATGTTGCGCTGCGTCTGTTATCCCGAGGGGTATCA
 GGTGATATGCCCGACGTCGCCGACGACGGGCTGGTTCATCATCAGTGTACACCGTCAAG
 ACGTTGCCCGGTAAGTTCGCGGGGTCAAGCAATCTGCGTAACCGGATATATCCGCTGGG
 GAAGCGCGATGATGACCTGGTATCAGATGGGCCCTGACGACACCGTGGTCCATCTGTC
 CGTTGACCTGCTATCGCGACTTCGTGACATCTCCATCTGATGACCGACTTTCAGCAGTG
 CTTGCTGACCCGATCGACCTTTCGCGCGTTCGATTCAGCTTCGCGACTACCTGCTGAA
 TCTCGCTCGCGAGCGAAGCTCGGCCGCGGTTAGTCCCGCGGGAAACCGGATCTCGCTACT
 GCGGGATCGGCTCGATCAGCTGCCGTCACCACTGTCTGTTACCGTGTGCTGCCGACGACTAG
 ACGCGTACGCGTCCGCGACCGCACCGACTTTTCAAGCGGTTTCGATGCGCTGTGCGCGA
 GCGTTTCGAGGTGCTCAGCCGCGCGCGGACGAGCACGAGCGACCGTCAACGTCGCCATC
 GTGACGGCATTCAGCCGGCCATCGCACGCTATGGCGATCGCGACCATTTCTGCTGACCCCT
 GACGACATGACCCCGCCACCGGTTCACTCCCGCGTGGGCGAGTTGGTGGGGGATTTCAACG
 GCACGAGCGTCTGAAGTCATGTCGCGGACGCGCACCTTCGCCGACTGCTGACGCGC

TABLE 6-continued

SEQ ID NOs: 1-5

GTCGGTGATCGTCTCTTCGACGACATGGACCATTGACACCCGGCGGCGTCAACGTGCCCCG
 GCTCCTGGGGCAGCGCAGCAGCAGCAGCAGGAGTGAGCAGACGCCCCGTCGTGTTACCTCGACA
 CTCGGCGCCACCGACCGAATCGACAGCGCGCCAAATCTTTGTTGACCCGATTCAAGGTGCG
 CGGTCTGAGCCAGACTCCTCAGGTGCTGCTGGACTGTGAGGTCGCGCGAAATCGACGGCATGC
 TTGAGGTCAACTGGGACACCCGCGATCAGGCCGTGCCGCGAGGTCTCGACCGCGGTTTC
 GCCGACTTCCGCCACGCGCTGGATCTGCTCAGCAGCCGCGCTCGGCTGGCACCGGCGCT
 ATTGCCGGCCAGCCCCGAGACCACACCGGTGAGGGGCCAGCACACACCATGAACCG
 GCGCTGATCCATACCGGTTCTCGCGCAACGTGCTGGTGACACCGGATGCGGTCGCCATCCG
 CCACGGTGATCGGGCCACCGTACCGCGAATTGCTCGCTGCCGCGACCGGGTGGCCGAC
 ACGCTGGCCGCGACGGGGTGGCGCGGTGACTACGTGGTATCCGGTTGCCGCGAGGGCC
 CCGCCAGATCGCGGCTCTCTGGGCGATTGCTGGCCGGCCGCTATGTGCGGTTGGAC
 GTCGGCTGGCCACCTACCGCGTCGACAGATCGCGGCCAATGCTCGCTGGCAGCGCTGTG
 CGAGCCGACGGAGAGGTGGACCGGCTGCTTGGCCAGCCGACAGCTGGTCACCGCGCGCG
 CGGTTGGTCCGGAGCCCCACAGCGAGGTGCT

SEQ ID NO: 5

CCGACTCCGGTAACACAGCAACCCGATCTGGGCTCGGTTGATGATTTACCCCGAGGACGCAAG
 GGCGTCCAAACCTCAAAGTGGTTCACCTCGGGCTGATGTGGGGACAACCGGGCCGGCA
 TCGGTGGCCTTGTCAAAAACCGGTTCTGTAATAAGGTCGATCCAAAGATCAGGTCGGCCCC
 CATACTCGGCTTGATGACCGCGCACCGCCAGGTCCGGTAGTTCGGGGTCCCGCCGGCT
 GGATCTGAACACCAACGAGGAGATCATGCCCCACCCCGCTGTCGATCCCGGAGGGT
 TGCAGTCGGTGGGTGATCCAGTCGAGGATGATGACTGGGTATCTGCCGTTAGCCCATGTG
 CCGGTGGCAAGAACCGCGCGGATTTGGGGACCCGTGCTCGAGCAGCAGGTAGTAGTCG
 CTGATCGGGAATACGGTCTGCTGAGCCGTTGATGAGCAGCGTCTCAGCGCAGGGCCGGTC
 CAGGATACCTTGGCGTACAAGGACAGCTTCCGGCGGTAGTCCACCCATTGCTCGAAGGTG
 CGCGCCCAAATGCAATAGCGAGGGTCTCGGCCAGCTCGAAGGGTACTCGCTGATTGGGC
 CTGCTCGATCCACTCGGCTCAAAGCCATGGTCTGATGAGCAGCGCTTGGCTGACCACAGG
 CGAGGGAGTCCGGTGGGTGTGGCGATTTTGGCCGCCAGTAGCCACCGGTACTACACCC
 CAATAGCCCAACCCCGCGGTCGAGTTCGGGTGCGCCGCGATCCACTCCAACACGGGG
 TGAACTTCGCTCGGCGTCTCCGAACAGCCAGCGGAGCATACCGACCCCGGGATGTGCG
 ATCGCAATGTTGCGACGTTCCGGCGAGCACCTGATCGGTGTGCATGCTTCTTGAAGGT
 GTCGATCCCGCCGAGACGAGCAATACAGGACCGGTCGGTCCGCTTCGGGACACGCAAA
 TGGGCGATGATGCGGTCAACCTCCCGGGCCGCGCGGAACGGGATCTCGACAGCTCGAT
 GGGGATGTGCAAGTACCGGATGCGCGCAGCAACATTTCTGCGACTTGGGTAGGCGCTCT
 TCTTACCGCCGAATTCATCGTCCGGTAGCGGGCCATGCGGTAGTAGCCATAGGCGCGAAA
 TAGTTTTCGTGTGCGCACTGGTGTGCGCAATCTGTTCGGCTCTTCCGCTCGGCTGATAG
 CGTGGCCACCGGCACTGAACCGCGCCCGCCAGGCGTCCGATCATAGGAGTTCAGGCCG
 TCAACACGCCCTGACATCATCGCGGAGCGTGTACTGAAACGGATAAATTCGCCCGTGGCC
 GAGACCGCATAGCGGCTAGCTCGGTACCCGGTCCGCTGTGTCGCTTCAACACGGCCAAAT
 TATGACCGCTCGCGGTCAGATATTTGGAGAAAACGGACATCTCACCGGTGCCGTGGGCGA
 GCCGACCGGGAAGTTCTGTGGGAGCCGCTCCAGCCAGTCCAGGCAATTCGCTGATCAGGTG
 GGACTGATCGGCGTAACCCGCTCGATCGCTAGATCGGCCACCGGATTCGGTGTACCCG
 CATGGATCAAATCCAGCAGCCGCTGGAACCGGAAAATCCGGCCAGCCGCTTGGGGCCGAA
 GCCGACCGCGGCTCGAACCGACGGCGAAGCTGCCGCTCGCTCAACCCAAACAGAGGAGGCG
 ACATCCGAACAGAGCGGGCCGGTGGACACGAGCATCTCGATTGAGTGCCTACTGGCT
 GATCGACACCGGATCGATCTTGACCAATACCGCGCCAGCACCGCCCAATAATGCGAC
 CCGTTGCCGAAAAGCTTGGCTCCAGCACATCTCAGCCATCCGGACGCGGATGTGCCGA
 ACACCGAGTCAACCGGATCTGGGTATCTTTAACTCGTGCAGGATGGCCAGCACCGCG
 GCGGACGACCCGTCGTAACCGCAGACCGATCATCGCACCTTCGGTCCCGCACATTGGT
 GTAAAGGTTGTGGCCGACGAAACCATGACTCGCCCTCGGCGGTGACGAACAGATCG
 ACGCAACCGTCCGGATCACCCGACGCGCCACCCGCGGATTCGAGCGCACCCAGCCAC
 AGTCCACATACGGCGCAGCGCACCAAAGTTCGACATCCAAATACCCGACCATCACGCT
 CACATTCAGCCAGGTCTACGTCGCTCCGCAACAAATGGGCGGCAATCTTCGAGCCGCG
 CGGTGAGCGCAGACTCCAGCGGCAATCTGTAACCGTAACTTCGGCGTAGAGGAAA
 TGCCCGCCAGGCGCAACCCCGCCACTTCCGGCGAAGCGTGACAGG

SEQ ID NO: 6:

CGGTGCGCGTAGGA

SEQ ID NO: 7

CGACCTGGTCTGCGTCATC

SEQ ID NO: 8

CCGCGAGGACCTGCTGACCG

SEQ ID NO: 9

GCGATCCGGCGAAACG

SEQ ID NO: 10

GACGAGTTAGACGCACAGATTCT

SEQ ID NO: 11

ACGCCCTGCAGTTAGTCCCCG

SEQ ID NO: 12

ATCGACCGGCTGGTGTAC

TABLE 6-continued

SEQ ID NOs: 1-5

SEQ ID NO: 13
GCGAGCAGCAGTTGAAC

SEQ ID NO: 14
TCGACGCGATGACCGCGCTCG

SEQ ID NO: 15
CAGACACCTGCAGAGTACTAC

SEQ ID NO: 16
AGCATGTTCGGCGACGTT

SEQ ID NO: 17
CAAGCCCGACTACGCCGCCCT

SEQ ID NO: 18
TCGCGGTGCCGATCAG

SEQ ID NO: 19
GTTTGGATGCCCGTGACATTC

SEQ ID NO: 20
CCGCGCTCCCCGAAATGCTC

SEQ ID NO: 21
CGGCGAACCCGTGACA

SEQ ID NO: 22
CACAGGTGCTCCCATTGAGT

SEQ ID NO: 23
CCGCGCCAGCGTGTAGGC

SEQ ID NO: 24
GCGCAGCTTCACCATCGA

SEQ ID NO: 25
GTGCGAAGGCTTCGAAGAC

SEQ ID NO: 26
AACAGGCTGCCGCCACGCT

SEQ ID NO: 27
GCCACGACGGTGACATCA

SEQ ID NO: 28
CCAGGGTGCTGGTGAACA

SEQ ID NO: 29
CCGCGTTGCGGCCAGCCCG

SEQ ID NO: 30
GGAAATCGATCTCGGTGGCTTTT

SEQ ID NO: 31
CCACCGCCGGAGCAA

SEQ ID NO: 32
CCGCGTGGCGCCGCA

SEQ ID NO: 33
GCCGCGACGGTAGGT

SEQ ID NO: 34
CCAGATCTCGGCAAGCGT

SEQ ID NO: 35
ACGACGCCGGCACAGCC

SEQ ID NO: 36
CGGACGGAGAGGTGGAC

SEQ ID NO: 37
GGGCTCCGGCACCA

SEQ ID NO: 38
TTGCCGACCCGACACCTG

TABLE 6-continued

SEQ ID NOS: 1-5

SEQ ID NO: 39
GGTAACCAGCAACCCGATCTG

SEQ ID NO: 40
CCCGAGGTGAACCACTTGA

SEQ ID NO: 41
TCGGTTGATGATTTACCCAGGACGC

SEQ ID NO: 42
GGCGTCTCCGAACCA

SEQ ID NO: 43
GCAACATTGGCGATCGACATC

SEQ ID NO: 44
CAGCGGAGCATCACCGACCCCGG

SEQ ID NO: 45
GGCTCGGCCTGATAGC

SEQ ID NO: 46
GCGGCCTGAACTCCTATGATC

SEQ ID NO: 47
CCGCGTTCAGTGCCGTGGCC

SEQ ID NO: 48
GCAATACTCGTGAACCGTAAACTG

SEQ ID NO: 49
GTGGCGCGGGTGTG

SEQ ID NO: 50
CCCTGGCGGGCATTTCCTCTACG

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 50

<210> SEQ ID NO 1

<211> LENGTH: 1863

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 1

```

cgggtgcgga ttgatcagcc gcccatgggc caccagcgat ccgtgttget tgagccgctg      60
ggtgcggaac ttgacctcgt agccgagcac cccgtcgagc tggccgagtt gctcgcgcag      120
gtagcggtag aggtcgtcgg aatctcggca gatggccaag gccatcaggt ttttgatcc      180
gcttgtggcg ccggcgaaag caatctgcgg atgggcccgc atacgttcgc cgaccgtgtg      240
cacgtcgcga ggcgcaaccg tgagccacag catggcgctg aggcgatggc cgagccgttc      300
gggtagtacg tcgacgtcgt agaccagggc gccgcaggcc tccagcgccg cgatccgcct      360
gctcaccgcg gccaccgacc aaccggtgcg cgcggccagt tgggactgcg gactgcggcc      420
gtcgtcggcg agcgcgcgca gcagcggcg gtcttcggcg gtgggcccgg accggtgagt      480
tcccgaactc ggctggggccc gcgcggccac gatccgctgc gcttgccggt ccgagagcgc      540
gctgcccgtg cccgtccagg gcgcgttcac cgggtcggcc aacgaatgca gcatcagatc      600
gatgctgatg tcggtgaccg cggcgggtgc cggtaggagc tgcgtcagca ggtcctcgcg      660
ggtgtcgcgc agcggcgcac ggatgacgca gaccaggctg gtccaccccg ccagcacatt      720
ggcgtgggac acgtcggggc gtcgcaccag cgcgtcggcc agcccgata tgcggtccgg      780

```

-continued

cgccgcacgg atgcggcaca ccagtcgcg gtcgccatcg gccacccggg tctccagccc	840
gacgacgcgc accaccccggt cgcgggcgag ccggtgatac gcgcgggcga ccgtctgctc	900
ggtggcgccc accacgctgg cgatccggcg aaacgagacc cggggagcta actgcagggc	960
gtgcagaatc tgtgctgcta actcgtcggg catcaggaaa agtatccgcg tggacaggtc	1020
caagtgagga tttcaggcca atttggtcgc ggcgagcgca atttcaacgc ctccgctgtc	1080
agcctggaag ctccgctaa ggagcgctca catgcacatc gaaatcggac tgcacagcca	1140
tatcgcaccac gtgcgggggc ggctgaccgt cgagtgggccc gcgcgggccc aacaccgcg	1200
ctttgccggc ctggccgca tcgaccggct ggtgtaccgg agcctcgacg cgatgaccgc	1260
gctcgcggtg gcccggggcg cgaccaccgg catcgggttg acgttcaacg tgetgctcgc	1320
tccgctgtac ccccggtgct tgctggccaa gcaggtcacc accctcgcgg aggcgtccgg	1380
cggccggctg cgcctggggc tgggcgctcg gagccgctgc gacgactaca ccgccgctcg	1440
cgctgactat cggcgggcgcg ggcgcattct cgacgagacg gcggcgctgc tgcgtgccc	1500
gtgtgagggc gaggtggtga ccggcgacca gccgctgtgc ccggcgccgg tgcggatccc	1560
gatcctgttc ggtggacggg cggacgcgac gatccgccc gccgtgaccg tgcgtgacgg	1620
ctggaccgcc gggcgctgc gcgattacgc caaccagtcg gtgttcgccc agcgggtccg	1680
ggcgcatgg gcggcgggg gccgctccgg acggccgctgg ctgcaggcca gcgtcaactt	1740
cgccttcggc gatgacgagg ccgtcgcgcg tggccgcaga cacctgcaga gctactacgg	1800
cttcaagccc gactacgccc cctcaacgt cgccgacatg ctcaccacgc cgcaggaggc	1860
ggc	1863

<210> SEQ ID NO 2

<211> LENGTH: 1551

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 2

cccgcaccgc cggctctgcc gccggacaat gcccggttgg cttcgcgggc tgcgcgctg	60
acgcggcctt cggcggcgac ttcggcgctc gcggcgccca gcacctcggc ttcacctgcc	120
atggcccgcg ggcccatggt gcgcatcgcc gtgctggctg tgggtggcctg ggtctcgacg	180
ctgaacagct tgtccttcaa catgcccgaga cccgtcccga tcggtgacag gatcgtcgag	240
gcgatgttgt agcccttgat cgccaaccat gccgtgccc ttgcaccac accgacggtc	300
accgcatcca gcaacggcct gtgctcggtg agccaatgcg ccgactcccc aagcacccca	360
acgacttcag tcatcgctgg cagaaagtcc tggccgatct cgatgcgggc tgcgctcaac	420
gcccccttga agtccctcat cttggcgctg taggtctcct gggattcggg gaatcccttg	480
gtattgccec caccgtcggc ctgcgcctga ctgaacctct tgaccgcggc agcggctttg	540
ggcccgttct ccccgaccaa ctgcaatacc gtccgcagat tgtcttggcc gccgaccatg	600
agggccagcg cttggatcat cgactggaca tcacccttgc cggacttgag ctgattgttg	660
aagcctgtca atgcttttcc ttggttccac cactgctcga caaggttggc ttcctcaacg	720
ctcaagccgc cgcggctcct gcggaactgc ttgtacgct cagcattgtt ctggatcgcc	780
gtcgcgacgg cctgcgcctg cgcgggcgac ttcgagaaca tctcattagc gctgtcggcc	840
acctgcttgc tcttgaacca ggcgtccagc gtgaccatcc ccgacgggccc aagcttcgac	900
tggatcgtgt catacaacag gttcgcggtg ccgatcaggc cgcgctcccc gaaatgctcc	960
tgaatgtcac gggcatccaa accgagctgt cccatctcct gggacatttg ctgggttggc	1020

-continued

ttggccagcg acgtgatcgc gtgcgtcata ttctgggtgg cctgatcagc acccataccc	1080
gactgagtga gcatcgccaa cgacgcatac aggtcattga cgctttcacc aactccggcc	1140
gcgaccggct cgacgttggt caacgcccc gcgaagtctt gcagcgaaac cttcgacaac	1200
cccacggcca cattcaattt cgaggtgacg ttggcggcct gctcgaccgg gatgtggtag	1260
tcgtgcatcg tggctcgtgac cgcgctgac gtctcgccca ggteccggcc ctcttcggac	1320
gcaagctcgg ctgaagccgt caacaccttg atcgcatac tgccccgata gccggccttc	1380
tcgacacca gcgcagcgtc catcagcttt tgaggcgcac aaccgaccac gctcgacagt	1440
ttcagcacgc catcgctgat cgccttcaga ttggccggcg actcctcggc caccgtgtgc	1500
agcctctgca acgactgctg gaaattcccc gccgcatac tcgccgagac a	1551

<210> SEQ ID NO 3

<211> LENGTH: 1847

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 3

acatcgggtg gagtgcgggg agatcgagca cgcgctgcgc ggccaccgc tggtcgcgc	60
cgcgacgggtg gtccccatcc acaactgcac tgcgctgggc gccgggatcg tcgtaaccgg	120
cagcggcgca gagcaatttg acgactccac acccggcgcg ctgcgcgcc atctcgcct	180
ccgactcccg cagtacatga tccccagggt gttcgtctca tgccccgagc tacgctcac	240
cgccaacggc aaggtcgacc ggggcaaat cgcggcgcgc ctggaagcgg ccgcgcgggc	300
accccagccg ctcgacacgt catccacct cactgtggtc gagcggctgg tcgccgaggt	360
ctggtccgat gtgctggggc cgcgcatcac cggccgcgag gacaattct tcgccaggg	420
cggcgatagc ctgcgcgcca ccgaagcggc cccccgactg acgcgcaggg gagtggcgg	480
agcggaggtg ggccagctgc tcagccacca gacgcttggg cagttcagcg cggcgtgtgt	540
gctcgcgcac ccggcatccg aggcatacca gtcggcggcc gatgtcggcg aaccctgac	600
accggggcag gggttccgc tcaccgggt gcagcaggcc tacacgctgg gcgcgccgg	660
actcaatggg agcacctgtg caccacgta ttctcgggtg gtgctggccg ccgcccga	720
gtctcgggt atagacctgg atcggtttgc ccgtgtggtc accagatgcg tcgacgaatt	780
cgcgatgctg cgggtcgcgc tggacgccga caccaccaa cgggtgcagg tcgacgccg	840
gccggtgccc gtccatgacc ttgatataca agacgacccc gacctgttac tgcggcgcac	900
ggcggcgcgc ccgttcgac cgcattcggc tccggtgac cagtgtctcg caccgtcgag	960
gtcaccctcg cacgtcggtc tgetgatcag ctatctgggc ctgatgccc gcagcctgtc	1020
caccgtcgtc accacgatca tcgccgaata ccagtgcga cccccggcgc gccaggtcga	1080
cccgaccgcg gcggtctctg cccggtttgc ctccgaaagc gcttggggcg aaaacgatgt	1140
cgacaacagc gttgccggcc ctccgctgct gccgctgcac gaccagcgac gtgaccctt	1200
cgagcgggtc acctcgcgc ggcgcagctt caccatcga gaacaggctg ccgccacgct	1260
gcgtgagcacc gccgcacacc tcggcgtcac cccaccgcgc ctggtctctg aagccttcgc	1320
acatgcgctg gcgtcgcac gcgcccgtca gcgattcgcg gtgacagtcc ccaagtcga	1380
ccgtcccgcac taeccccccg cggaccgcga ggtgttgggc aacttcaccc gcctggcgtc	1440
gtgcgaggtc gactacggcg ccgtgagacc gggatctgcc gaagcgggtt ccgcccgcga	1500
gccggaaactg tggcgcgcgg tgagccacga cggtgacatc accggcgggc tggcccgaac	1560
gcggaccgcg ggtggctacc ccgtggtgtt caccagcacc ctggggctca ccatcagga	1620

-continued

cgccagcggg	ctgaccaacg	tgcggacatt	gaccagacc	ccggcgctct	ggctggactg	1680
ccagaccgag	gacgaggteg	ccggaattcg	tatgagctgg	gacatagcta	ccaatgtggt	1740
tgcccgcgaa	tcgatctcgg	tggttttttc	ccgattcgag	gaggcggtgc	ggcgccacgc	1800
ggggcaagcc	gagccgcgg	gcacggccgt	tgctccggcg	gtgggcg		1847

<210> SEQ ID NO 4

<211> LENGTH: 4716

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 4

agtggcgag	cgcggtgac	gcccgcgc	tgcgcaactg	ccgaccgag	cagggtgtgc	60
cgcaatacac	catgctggg	cgcgctggg	aggcaactgc	atacgtcccc	tcaggatacg	120
ccgcttcga	cgtegaacg	gcgctcggc	ggttgccgg	catcgteacc	ggagccgtgt	180
caccccgac	gctgatcga	gaccgcagc	tgacgcccga	ggcgctgctg	ctgctgacg	240
accgcatcg	gatggccctc	gacgacctg	ccggggccat	cttcgggac	gcgcgacgc	300
tggggcgcg	gcttcgctc	gtggaagtgc	gctcccggac	cggcctgac	accgaacggt	360
tgaccgagct	ggtggcgctg	gtggctcagg	aataactgtg	cctggagcca	aaccgcagc	420
tcgccggaat	cgccgcggg	cgcgcttcc	cgcccccac	ccgcccagtc	gacgcaaccg	480
acgcgcgctc	ggcgctgac	gtgggtgatc	gctgtgggtc	gctgcatcag	ctgcccagac	540
cggaggcggt	cctcgaggcg	atcacctgt	ccgacgacg	ttggctgtgg	atggtggaga	600
attccgaggc	cacccaagcg	acgctcatca	gcgcgccgt	tctcgacccc	ggcctgctcg	660
cgtccgatcc	gaagacgctg	cgctccggcg	atcggtggtg	gcggtcctac	gccgaccacg	720
gttgcgcaac	gacgcacatg	atccaggacg	gaccggcct	cacgctcacc	gcgcacggcc	780
ccgacaagcc	cgcatgccc	acaccgccc	ccgaacagcg	ccgacgaggt	aggtggtcgc	840
gaccggctgt	gcccggctc	tcgctgccga	ccgacggcac	ggtgggtggc	acgcttgccc	900
agatctggca	ggtcatctc	gccattocaa	caccggcgt	cgacgacgac	ttcttctcgc	960
tcggcgggta	cagcctcgtc	gcgaccggg	tctacggcca	ccttcggggc	gccggtttcg	1020
gccaactcgc	ttctgctgac	ctgttcaacc	actcgacgct	cggtgagctc	gcggcacacg	1080
ccggcccgcg	caccggccc	gaagtgtcgg	tggcggtga	gtcgaccggg	ggcggcaccc	1140
acgacccgaa	ccgattccc	ctcacctcgc	tgacgaacgc	gtatcggggc	ggcgagagaag	1200
gcgcttggat	cctcgggggc	gtcgccgcgc	actgctactt	cgagttcgag	ctcgcgagct	1260
tcgacccggc	gagattcgat	tcggccgcac	gccaaactcg	agcacggcac	gccggactgc	1320
gcaccacggt	gtcaccggcg	ggcaccgacg	cgccctcctc	gggtgaggtc	gccgtcgtgc	1380
acaccgcgcc	gatcgagccc	gtcgtgagag	accacgacga	cggtcgagcc	gcatgagcgc	1440
accagatcat	cgacttgacg	gcccggccc	gcatcgactt	gggggtgcaa	accgcggcg	1500
acgggcgcac	cgctgctggc	atcagcatgg	acaacaccat	gctcgacggc	gccagcatga	1560
tgatcgccct	gtcccgaactc	gatcacctct	atcgcgggca	aaccggtgac	caattgccc	1620
cgctggaaac	gtcttttcg	cactacgtgt	ggaaccaccc	ggagctgctg	cccgaacggc	1680
acgaggcggt	gctgcccggg	ctggccgcca	gccgagacta	ttggcgcgca	cgctgccc	1740
ctttgcccgc	ggcgccgaaa	ttggccgaca	tgtaactgct	gttcgagatc	gaggagccga	1800
ggttcgaacg	ggcaaccggc	accattccc	ccgtcgactg	gtcgaggtga	acgcatcgt	1860
gccgtgccga	ggcgctcacc	gtcgctcat	ttctgctcgc	caactatgca	cggtgctgt	1920

-continued

ctcggtggtc	ggggaccgac	cacttctgca	tcaacgtcac	gctattcgac	cgcgaccccc	1980
atgtcgtggg	gategaaaac	gtcgtcggag	atttcaactc	cctggtgttg	ttggagtgcc	2040
gagtcgatga	gcccgcctcg	atctgggaga	gcgtgcgcgc	tctgcagcgg	caattgatga	2100
ccgacctgcc	gcaccgcggc	gcggacgcgg	tgtggctgca	acgcgaactg	ctgcggtttc	2160
acggcaaccc	gacggcgcgg	ctgtttcccg	tctgtctcac	cagcggactg	ggccttgtcg	2220
acgcctcggc	tccggcggcg	gtccggttcg	ccgaaccggt	attcgcgcc	tgcagacac	2280
cgcagacggt	gttggacttc	caggtgtggg	aaagcgcggg	ggcgtgaag	ctgtcgtggg	2340
acttcgtcag	tcaggcgggtg	tcccccgcc	ccgcgcgcac	tcagctcgag	tcgttggtgg	2400
acggcateac	cggtgtcgcc	acacgcagcc	gccgcacga	acacaagttg	ggcgaggggg	2460
catccaatga	cgagctcctg	caacgtggtt	cgaggatctg	cgctccgcc	ctgggtcagc	2520
cgagggtcga	acctcaegac	aattttcttc	agctcggcgg	cgattcggtc	agcgcgacca	2580
aggtggtcga	acagatcggc	cgtgagctgt	cagcctcggc	caccttcga	ctgctgttcg	2640
ccaatccggt	gatcggcgac	ttcgcgcga	aaatcgcga	cactgacaac	gccgacgaac	2700
ccgacctgac	cgttgaggag	ggcatgttat	gaccgcggcc	gagctcgtcg	accacctgcg	2760
gggtatcggc	gtccaactgt	gggcgcagcg	tgagaatttg	cgctaccgag	caccgcaaca	2820
agtcctcaoc	gcggacctga	aagctcaact	ggcggcggtc	aaaacggacg	tgattacctt	2880
gctggcggaa	gagacgaccc	tgtcgcgcgc	gccgcaggac	cggttcgagc	cgtttcgcct	2940
caccgacgtg	caagccgcat	atctggtcgg	gcgcacgtcg	gcgtttcagt	ggggcggggt	3000
aggctgccac	ggctacgccc	agttcgcggg	cgaccacacc	gtggcaacac	cgagcgcga	3060
gcaatatacg	gaggcgtggc	gcaaggttgc	cgaccgccac	gacatgttgc	gctgcgtcgt	3120
tcataccgag	gggtatcagg	tgatatgccc	cgacgtgccc	gacgacgggc	tggtcatcca	3180
tcagtgtcac	acggtcgaag	acgttgccgg	cgtacgtgcc	ggggtcacgg	aacatctgcg	3240
taaccggata	tatccgctgg	gcgaagcgc	gatgtatgac	ctggtgatca	cgatgggccc	3300
tgacgacacc	gtggtccatc	tgtccgttga	cctgctgac	gccgacttcg	tcagcatctc	3360
catcctgatg	accgactttc	agcagtgett	gcttgacccc	gaatgcgacc	ttgcgcccgt	3420
cgatttcagc	ttccgcgact	acctgtgaa	tctcgtcgc	gagcgaagct	cgccgcgcgg	3480
tagtgcccgc	cgggaacgcg	atctcgcta	ctggcgggat	cggtcgtatc	agctgcccgc	3540
accactgtcg	ttaccggtgc	tgcccagcga	ctagacgccc	tcagcgtcgc	ggaocgcacc	3600
gaccttttca	cggcgttcga	tgcgcctgtc	ggcgcagcgt	ttcgaggtgc	tcagccggcg	3660
cgccgcagag	cacggagcga	ccgtcaacgt	cgccatcgtg	acggcattca	gccgggccc	3720
cgcacgctat	ggcgatcgcg	accatttcct	gctgaccctg	acgaccatgg	accgccacgc	3780
gttcaactccc	gccgtcgggc	agttggtcgg	ggatttcacc	ggcaccagcg	tgctcgaagt	3840
cgatgtccgc	ggacagcga	ccttcgcca	actgctgcac	ggcgtcggtg	atcgtctctt	3900
cgacgacatg	gaccattcga	ccaccggcgg	cgtcaacgtc	gcccggtccc	tggggcagcg	3960
cgacgacgac	cgaggtgagc	agacgcccgt	cgtgttcacc	tcgacactcg	gcgccacgac	4020
ccgaatcgac	agcggcgcga	catccttgtt	gcaccgatt	caaggtcgcg	gtctgagcca	4080
gactcctcag	gtcgtcgtgg	actgtcaggt	cgccgaaatc	gacggcatgc	ttgaggtcaa	4140
ctgggacacc	cgcgatecag	ccgtgcccgc	cgaggtcctc	gaccgcgcgt	tcgcccactt	4200
ccgccacgcc	ctggatctgc	tcagcaccca	cgccctggcc	tgccaccggc	cgctattgcc	4260
ggcccagccc	cccagacca	caccggtcga	ggggccacgc	acacaccatg	aaccggcgct	4320

-continued

gatccatacc gggttcctgc gcaacgtgct ggtgacaccg gatgccgtcg ccatccgcca	4380
cggtgatcgg gccaccacgt acgccgaatt gctcgtgccc gcgaccgcgg tggccgacac	4440
gctggccgcg acgggggtgc ggccgcgtga ctacgtcggg atccggttgc cgcagggccc	4500
cgcccagatc gcggtctccc tggggcatt gctggcccgg gccgctatg tgcggttga	4560
cgtcggctgg cccactcacc gcgtcgacca gatcgcggcc caatgctcgc tggcagcgct	4620
gtgcgagccg gacggagagg tggaccggct gcttgcccgc ccgcagacct ggtcaccgcg	4680
cgccggcggg gtgcccggag cccacagcga ggtgct	4716

<210> SEQ ID NO 5

<211> LENGTH: 2392

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 5

ccgactccgg taaccagcaa ccgatctgg gctcggttga tgatttcacc ccaggacgca	60
agggcgctcc aacctcaaa gtggttcacc tcgggctgat gtggggacaa cacggggccg	120
gcatcggtgg ccttgtcaaa aaccggttcg tgaaaatggt cgatccaaa gatcaggctc	180
gccccatac attcggcttg gatgaccgcg gcacgccagg tccggtagtt cggggtccc	240
gccggctgga tctgaacacc aacggaggag atcattgccc gccaccggc tgtcgatccc	300
gcgagggttg cagtcggtgg gtgatccagt cgaggatgat tgactgggta tctgcccgtg	360
agcccatgtg cccggtgcca aagaaccgcg ccgatttggg ggaccgctgc tcgagcagca	420
ggtagtagtc gctgatcggg aatacgggtg cgtgtagccc gttgatgagc agcgtctcag	480
cgcagggccg gtccaggata ccttggcgtg caagggacag cttcggcgcg tagtccacc	540
attgctcgaa ggtgtcgcgc ccaaatgcat agggcagggg ctcggccagc tcgaaggggt	600
actcgcctga ttgggcctgc tcgatccact cggcgtcaaa gccatggtcg atgcagccgc	660
cttgctgac cacacaggcg agggagttcc ggtgggtgtg ggcgattttg gccgcccagt	720
agccaccggt actaccacc caatagocca caccgcgcg gtccagttcg ggtcggccg	780
cgatccactc caacacgggg gtgaacattc gctcggcgtc ctccgaacca gccagcggag	840
catcaccgac cccgggatg tcgatcgcca atgttgccgac gttccgggcg agcacctgat	900
cggtgtgcat gtcttctctg aagggtgtga tcccgccga gacgagcaat acaggcagcc	960
ggtccgtcgc cttcgggaca cgcagatggg cgatgatcgc gtcaccctcc ccgggcccgc	1020
cggcgaacgg gatctcgaca cgctcgatgg ggatgtcgaa gtaccgggat gcgcgcagca	1080
acatttctg cgacttcggg taggcgctct tcttaccggc cgaattcacc gtcgggttagc	1140
gggcatcgcg gtagtagcca taggcgcgca aatagtttc gtgtgccaca ctggtgtcgc	1200
caatctgttc ggctctctcg gctcgggct gatagcgtgc ggccacggca ctgaaccgcg	1260
ccgccaggc gtcgcatca taggagttca ggccgctcaa cacgccctgc acatcatcgg	1320
cgagcgtgta ctgaaacgga taaattccc cctgggccga gaccgccata gcggccatag	1380
ctcggtcacc ggtcgcgtgt gtcgcttcac acggccaagt tatgaccgct cgcggctgca	1440
gatattggag aaaacggaca tctcacgcgt gccgtgggcg agccgcaccg ggaagtctg	1500
tgggagccgc tccagccagt gccaggcatt cgctgatcag gtgggactga tcggcgtaac	1560
ccgctcgat cgctagatcg gccaccggga ttcggtgat accggcatgg atcaaatcca	1620
gcagccgctg gaaccgaaa atccggccaa gccgcttggg gccgaagccg accgcccgt	1680
cgaaaccgag gcgaagctgc cgctcgctca acccaacaga ggaggcgaca tccgaaacag	1740

-continued

```

agcggggcgg gtggacacgc agcatctcga ttgagtgcgc cactggctga tcgaccacgg 1800
gatcgatctt gaccacatac cgcgccagca ccgcccga taatgcgacc cgttgccgaa 1860
aagcgttggc ctccagcaca tcctcagcca atccggacgc ggatgtcccg aacaccgagt 1920
caaccgggat ctgggtatct cttaactcgc tgacaggatg gccacgaccc gcggcggcag 1980
caccggctcg taaccgcaga ccgatcatcg caccttcggt gcccgcacat tggctgtaaa 2040
aggttgtggc cggaccagaa accatgactg cgccctcggc ggtgacgaac agatcgacgc 2100
aaccttccgg catcaccgcg agcgcgccac ccgcggtatt cgagcgcacc cagccacagt 2160
ccacatacgg cgcgagcgcg cccaaaggtc gacactcaa ataccgacc atcacgctca 2220
cattcagcca ggtctactgt cgctccgcaa caaattgggc cgcaatcctt cgaagccgcc 2280
cggtcagcgc agactcccag gcggcaatac tcgtgaaccg taaactgcgg cgtagaggaa 2340
atgcccgcga gggcgcaaca cccgcgccac ttcggcgaga aagcgtgaca gg 2392

```

```

<210> SEQ ID NO 6
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 6
cggtgccg tagga 15

```

```

<210> SEQ ID NO 7
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 7
cgacctggtc tgcgtcatc 19

```

```

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        probe

```

```

<400> SEQUENCE: 8
ccgcgaggac ctgctgacgc 20

```

```

<210> SEQ ID NO 9
<211> LENGTH: 16
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 9
gcgatccggc gaaacg 16

```

```

<210> SEQ ID NO 10
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 10

gacgagtttag acgcacagat tct 23

<210> SEQ ID NO 11
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 11

acgccttgca gttagctccc cg 22

<210> SEQ ID NO 12
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 12

atcgaccggc tgggtgac 18

<210> SEQ ID NO 13
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 13

gcgagcagca cgttgaac 18

<210> SEQ ID NO 14
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 14

tcgacgcat gaccgcgctc g 21

<210> SEQ ID NO 15
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 15

cagacacctg cagagctact ac 22

<210> SEQ ID NO 16
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 16
 agcatgctcgg cgacggt 17

 <210> SEQ ID NO 17
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 17
 caagcccgac tacgccgcc t 21

 <210> SEQ ID NO 18
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 18
 tcgcggtgcc gatcag 16

 <210> SEQ ID NO 19
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 19
 gtttggatgc ccgtgacatt c 21

 <210> SEQ ID NO 20
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 20
 ccgcgctccc cgaaatgctc 20

 <210> SEQ ID NO 21
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 21
 cggcgaaccc gtgaca 16

 <210> SEQ ID NO 22
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 22

cacaggtgct cccattgagt 20

<210> SEQ ID NO 23
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 23

ccgcgcccag cgtgtaggc 19

<210> SEQ ID NO 24
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 24

gcgcagcttc accatcga 18

<210> SEQ ID NO 25
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 25

gtgcgaaggc ttcgaagac 19

<210> SEQ ID NO 26
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

<400> SEQUENCE: 26

aacaggctgc cgccacgct 19

<210> SEQ ID NO 27
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 27

gccacgacgg tgacatca 18

<210> SEQ ID NO 28
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 28

 ccagggtgct ggtgaaca 18

 <210> SEQ ID NO 29
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 29

 ccgcgttgcg gccagcccg 19

 <210> SEQ ID NO 30
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 30

 ggaatcgatc tcggtggcctt tt 22

 <210> SEQ ID NO 31
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 31

 ccaccgccgg agcaa 15

 <210> SEQ ID NO 32
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 32

 ccgcgtggcg ccgca 15

 <210> SEQ ID NO 33
 <211> LENGTH: 15
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 33

 gccgcgacgg taggt 15

 <210> SEQ ID NO 34
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 34

 ccagatctcg gcaagcgt 18

 <210> SEQ ID NO 35
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 35

 acgacgccgg cacagcc 17

 <210> SEQ ID NO 36
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 36

 cggacggaga ggtggac 17

 <210> SEQ ID NO 37
 <211> LENGTH: 14
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 37

 gggctccggc acca 14

 <210> SEQ ID NO 38
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 38

 ttgccgaccc gcagacctg 19

 <210> SEQ ID NO 39
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 39

 ggtaaccagc aaccgatct g 21

 <210> SEQ ID NO 40
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 40

 cccgaggtga accacttga 20

 <210> SEQ ID NO 41
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 41

 tcggttgatg atttcacccc aggacgc 27

 <210> SEQ ID NO 42
 <211> LENGTH: 16
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 42

 ggcgtcctcc gaacca 16

 <210> SEQ ID NO 43
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 43

 gcaacattgg cgatcgacat c 21

 <210> SEQ ID NO 44
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic probe

 <400> SEQUENCE: 44

 cagcggagca tcaccgaccc cgg 23

 <210> SEQ ID NO 45
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

 <400> SEQUENCE: 45

 ggctcgggcc tgategc 17

 <210> SEQ ID NO 46
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 46

gcggcctgaa ctctatgat c                               21

<210> SEQ ID NO 47
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    probe

<400> SEQUENCE: 47

ccgcgttcag tgccgtggcc                               20

<210> SEQ ID NO 48
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 48

gcaataactcg tgaaccgtaa actg                         24

<210> SEQ ID NO 49
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 49

gtggcgcggg tgttg                                    15

<210> SEQ ID NO 50
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    probe

<400> SEQUENCE: 50

ccctggcggg catttctct acg                            23

```

What is claimed is:

1. A method for differentially detecting *Mycobacterium avium paratuberculosis* (MAP) comprising:

obtaining nucleic acids from a sample;

contacting the nucleic acids from the sample with at least one primer set, having one forward primer and one reverse primer, wherein the at least one primer set is

selected from: SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO:

37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO: 49; and full complements thereof;

amplifying at least one MAP-specific target nucleic acid from the sample to form an amplified nucleic acid,

wherein the at least one MAP-specific target nucleic acid is selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, and full complements thereof, wherein the foregoing target nucleic acids are present in MAP organisms and absent in non-MAP *Mycobacterium Avium Complex* (MAC) organisms, wherein the non-MAP MAC organism is selected from the group consisting of *Mycobacterium*

67

bacterium avium subsp. *Avium* (MAA), *Mycobacterium avium* subsp. *silvaticum* (MAS), and *Mycobacterium* subsp. *hominissuis* (MAH),

wherein detecting the amplified nucleic acid confirms the presence of a MAP organism in the sample.

2. The method of claim 1, wherein the step of detecting an amplified nucleic acid comprises using a probe that can hybridize to and reveal the presence of the amplified nucleic acid.

3. The method of claim 2, wherein the probe is selected from SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, SEQ ID NO: 50, full complements thereof and labeled derivatives thereof, wherein the label is selected from a dye, a radioactive isotope, a chemiluminescent label, an enzyme, a fluorescein dye, a rhodamine dye, a cyanine dye and combinations thereof.

4. The method of claim 1, wherein the sample is an veterinary sample, an animal sample, a food sample, an agricultural sample, a produce sample, a clinical sample, an environmental sample, a biological sample, a water sample or an air sample.

5. The method of claim 4, wherein the animal sample is obtained from a cow, a sheep, a bison, a ruminant, a deer, a rabbit, a fox or a bird.

6. The method of claim 1 further comprising:

detecting the presence of at least two MAP-specific nucleic acids that are absent in non MAP MAC organisms comprising:

amplifying from the sample a first MAP specific nucleic acid by contacting nucleic acids present in the sample with at least a first primer set, comprising one forward primer and one reverse primer, the first primer set designed to amplify the first MAP specific nucleic acid; and amplifying simultaneously from the same sample a second MAP specific nucleic acid by simultaneously contacting nucleic acids present in the sample with at least a second primer set, comprising one forward primer and one reverse primer, the second primer set designed to amplify the second MAP specific nucleic acid, wherein the contacting in steps a) and b) is performed under conditions suitable for a nucleic acid amplification reaction,

68

wherein the first and the second MAP-specific nucleic acids that are absent in non-MAP MAC organisms, are selected from the group consisting of SEQ ID NO:1, SEQ ID. NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, and the full complements thereof; and

detecting at least one amplified nucleic acid amplified by either the amplification reactions of steps a) or b), wherein detection of at least one amplified nucleic acid indicates the presence of a MAP organism in the sample.

7. The method of claim 6, further comprising detecting the first and the second amplified nucleic acids amplified by the amplification reactions steps a) and b).

8. A kit for use in the method of claim 1 comprising:

at least one synthetic primer set selected from SEQ ID NO: 6 and SEQ ID NO: 7; or SEQ ID NO: 9 and SEQ ID NO: 10; or SEQ ID NO: 12 and SEQ ID NO: 13; or SEQ ID NO: 15 and SEQ ID NO: 16; or SEQ ID NO: 18 and SEQ ID NO: 19; or SEQ ID NO: 21 and SEQ ID NO: 22; or SEQ ID NO: 24 and SEQ ID NO: 25; or SEQ ID NO: 27 and SEQ ID NO: 28; or SEQ ID NO: 30 and SEQ ID NO: 31; or SEQ ID NO: 33 and SEQ ID NO: 34; or SEQ ID NO: 36 and SEQ ID NO: 37; or SEQ ID NO: 39 and SEQ ID NO: 40; or SEQ ID NO: 42 and SEQ ID NO: 43; or SEQ ID NO: 45 and SEQ ID NO: 46; or SEQ ID NO: 48 and SEQ ID NO:49; and full complements thereof; and

at least one labeled probe selected from SEQ ID NO: 8, SEQ ID NO: 11, SEQ ID NO: 14, SEQ ID NO: 17, SEQ ID NO: 20, SEQ ID NO: 23, SEQ ID NO: 26, SEQ ID NO: 29, SEQ ID NO: 32, SEQ ID NO: 35, SEQ ID NO: 38, SEQ ID NO: 41, SEQ ID NO: 44, SEQ ID NO: 47, and SEQ ID NO: 50, and full complements thereof, wherein the label is selected from a dye, a radioactive isotope, a chemiluminescent label, an enzyme, a fluorescein dye, a rhodamine dye, and a cyanine dye;

one or more components selected from a group consisting of: at least one enzyme, dNTPs, at least one buffer, at least one salt, at least one control nucleic acid sample; and

an instruction protocol.

9. The kit of claim 8 wherein the probe is labeled with two labels.

* * * * *